

***Helicobacter suis* and the stomach-brain axis: epidemiological, pathogenetic and diagnostic aspects**

Caroline Blaecher

A dissertation submitted in fulfillment of the requirements for the degree of Doctor of Veterinary Science (PhD),
Faculty of Veterinary Medicine, Ghent University, 2017

Promoters:

Prof. dr. F. Haesebrouck

Prof. dr. A. Smet

Dr. S. Dobbs

Dr. J. Dobbs

Ghent University, Faculty of Veterinary Medicine
Department of Pathology, Bacteriology and Avian Diseases

Members of the Examination Committee

Prof. dr. Herman Favoreel
Chairman, Ghent University, Belgium

Prof. dr. Richard Ducatelle
Ghent University, Belgium

Prof. dr. Roosmarijn Vandenbroucke
Ghent University, Belgium

Prof. dr. Steffen Backert
Friedrich-Alexander-University Erlangen-Nürnberg, Germany

Prof. dr. Benedicte De Winter
Antwerp University, Belgium

List of abbreviation	1
Review of the Literature.....	5
1. The rise of the genus <i>Helicobacter</i>	7
2. Gastric <i>Helicobacter</i> species in humans with emphasis on zoonotic pathogens.....	10
2.1 Phylogeny of animal-associated gastric <i>Helicobacter</i> species.....	11
2.2 Clinical significance of gastric non- <i>Helicobacter pylori</i> <i>Helicobacter</i> infections in humans.....	13
2.3 Diagnosis of gastric <i>Helicobacter</i> infections.....	14
2.3.1 Non- or minor invasive tests.....	16
2.3.1.1 The urea breath test.....	16
2.3.1.2 Stool antigen test (SAT).....	17
2.3.2 Invasive tests.....	17
2.3.2.1 Endoscopy and analysis of gastric biopsies.....	17
2.3.2.1.1 Histology.....	18
2.3.2.1.2 Polymerase chain reaction.....	19
2.3.2.1.3 Culture.....	19
2.3.3 Conclusion.....	19
3. <i>H. suis</i> : a porcine pathogen with a zoonotic potential for humans.....	20
3.1 Prevalence of <i>H. suis</i> in pigs.....	20
3.2 Pathogenic significance of <i>H. suis</i> in its natural hosts and experimentally infected laboratory animals.....	20
3.2.1 <i>H. suis</i> infection in pigs.....	20
3.2.2 <i>H. suis</i> infections in non-human primates.....	21
3.2.3 <i>H. suis</i> infections in experimentally infected laboratory animals.....	22
3.2.4 <i>H. suis</i> infections in man.....	23
3.3 Use of genomics and genetics to obtain better insights in the pathogenesis of <i>H. suis</i> infections.....	24
3.3.1 pH homeostasis.....	25
3.3.2 Chemotaxis and motility.....	25
3.3.3 Oxidative stress.....	26
3.3.4 Outer membrane proteins.....	26
3.3.5 Virulence factors contributing to the gastric lesions.....	27
3.4 Host immune response to gastric <i>Helicobacter</i> infections.....	28
3.4.1 Innate immune response to <i>H. pylori</i> and <i>H. suis</i>	28
3.4.2 Acquired immune response to <i>H. pylori</i> and <i>H. suis</i>	30
3.4.2.1 T helper cells.....	30
3.4.2.2 B cells.....	32
4. Extragastric manifestations of <i>Helicobacter</i> infections with emphasis on Parkinson's disease (PD).....	35
4.1 PD: clinical outcome.....	38
4.1.1 History and pathological findings.....	38
4.1.2 Motor and non-motor symptoms.....	40
4.1.3 Incidence and prevalence.....	41
4.1.4 Etiology.....	41
4.2 Pathogenesis of PD: dysfunction of neural barriers.....	42
4.2.1 Blood-brain barrier (BBB).....	42
4.2.2 Blood-cerebrospinal fluid barrier (blood-CSF barrier).....	43
4.3 Involvement of <i>H. pylori</i> infection in PD: what is known so far?.....	45
Scientific Aims.....	75
Experimental Studies.....	79
1. <u>Chapter 1:</u> Significantly higher frequency of <i>Helicobacter suis</i> in patients with idiopathic parkinsonism than in control patients.....	81
2. <u>Chapter 2:</u> The choroid plexus epithelium as a novel player in the stomach-brain axis during <i>Helicobacter</i> infection.....	93
3. <u>Chapter 3:</u> A novel isolation protocol and probe-based RT-PCR for diagnosis of gastric infections with the zoonotic pathogen <i>Helicobacter suis</i>	123
General Discussion.....	149
Summary.....	174
Samenvatting.....	179
Curriculum Vitae.....	185
Bibliography.....	189
Dankwoord.....	193

List of abbreviations

AGS	human gastric adenocarcinoma cells
AhpC	alkyl hydroperoxide reductase
AlpA/B	adherence-associated lipoprotein A/B
AnsB	l-asparaginase II
aPKC	atypical protein kinase C
BabA	blood group antigen binding adhesion A
BBB	blood-brain barrier
Blood-CSF barrier	blood-cerebrospinal fluid barrier
CagA	cytotoxin-associated gene A
CagPAI	cag pathogenicity island
CarR	non-coding gene in the genome of <i>Helicobacter suis</i>
Che	chemotaxis protein
CNS	central nervous system
Cldn	Claudin
COX	cyclo-oxygenase
CPE	choroid plexus epithelium
CSF	cerebrospinal fluid
CXCL	chemokine ligand
DC	dendritic cell
DNA	deoxyribonucleic acid
ECIS	electrical cell impedance system
EGF	epithelial growth factor
EIA	enzyme immune assay
ErbB	epidermal growth factor receptors
ERK	extracellular signal-regulated kinase
FCS	fetal calf serum
FITC	fluorescein isothiocyanate
FlaA/B	flagellin A/B
FldA	flavodoxin protein
FlgD	flagellar cap
FlgE	flagellar hook
FOXP3	forkhead box protein P3
Ggt	gamma-glutamyl transpeptidase
HBSS	Hank's balanced salt solution
HCL	hydrochloric acid
HE	hematoxylin and eosin
Hof	<i>Helicobacter</i> outer membrane protein family

Hom	<i>Helicobacter</i> outer membrane
Hop	<i>Helicobacter</i> outer membrane porin
Hor	hop related protein
HP-NAP	<i>Helicobacter pylori</i> neutrophil-activating protein
Hsp60	heat shock protein 60
HtrA	high temperature requirement A
IBD	inflammatory bowel disease
ICA	immunochromatography assay
IceA	ulcer-associated protein restriction endonuclease
IFN	Interferon
IL	Interleukin
iNos	inducible nitric oxide synthase
IP	idiopathic parkinsonism
KatA	Catalase
KC	keratinocyte-derived chemokine
Levadopa	L-3,4-dihydroxyphenylalanine
LIX	LPS-induced CXC chemokine
LPS	Lipopolysaccharide
LRRK2	leucine-rich repeat kinase 2
MALT	mucosa-associated lymphoid tissue
MAP	mitogen-activating protein
MIC	minimum inhibitory concentration
MIP	macrophage inflammatory protein
MLST	multi-locus sequence typing
MPTP	1-methyl-4-phenyl-1,2,3,6-tetrahydropyridin
mRNA	messenger ribonucleic acid
MRPS6	mitochondrial ribosomal protein S6
Muc	Mucin
NADPH	nicotinamide adenine dinucleotide phosphate
NAP	neutrophil-activating protein
ND	not detected
NF	nuclear factor
NHPH	non- <i>Helicobacter pylori Helicobacter</i>
NLRP	nucleotide-binding oligomerization domain-like receptor family, pyrin domain containing 3
NOD	nucleotide-binding oligomerization domain
NSAID	non-steroidal anti-inflammatory drug
Ocln	Occluding

OHDA	Hydroxydopamine
OipA	outer inflammatory protein A
OMP	outer membrane protein
PAMP	pathogen-associated molecular patterns
PAR	pseudo-autosomal region
PARK	parkin gene
PBS	phosphate-buffered saline
PCR	polymerase chain reaction
PD	Parkinson's disease
PgbA/B	plasminogen-binding protein A/B
PH	measure of acidity or basicity
PINK	phosphatase and tensin homolog (PTEN)-induced putative kinase 1
PorA/B/C/D	pyruvate oxidoreductase complex
PPI	proton pump inhibitor
PRR	pattern recognition receptor
PrtC	gene for the production of collagenase
PTEN	phosphatase and tensin homolog
PutA/B	proline utilization A/B gene
RdxA	oxygen-insensitive NAD(P)H-nitroreductase gene
ROS	reactive oxygen species
rRNA	ribosomal ribonucleic acid
RT	room temperature
RT-PCR	real-time PCR
RUT	rapid urease test
SabA	sialic acid binding adhesion A
SAT	stool antigen test
SodB	superoxide dismutase
SPF	specific pathogen free
Spp.	Species
SSCR	simple sequence cytosine repeat
T4SS	type IV secretion system
TEER	transepithelial electrical resistance
TGF	tissue growth factor
Th	T-helper cell
Tlp	methyl-accepting chemotaxis proteins
TLR	toll-like receptor
TNF	tumour necrosis factor
Treg	regulatory T cell

List of abbreviations

UBT	urea breath test
UreA/B	urease subunit A/B
VacA	vacuolating cytotoxin
VIP	vasoactive intestinal peptide
VIPR	vasoactive intestinal peptide receptor
ZO	zonulae occludentes

REVIEW OF THE LITERATURE

The present thesis deals with *Helicobacter suis* (*H. suis*), a bacterium that naturally colonizes the stomach of pigs and non-human primates. Infections with this microorganism have also been associated with gastric disease in humans.

Here, we first present an overview of the genus *Helicobacter* and the prevalence, phylogeny and clinical significance of gastric *Helicobacter* infections in humans. Then, the different diagnostic methods for detection of gastric *Helicobacter* infections are discussed. Additionally, the prevalence of *H. suis* in pigs and the pathogenic significance of this agent for pigs, non-human primates and experimentally infected lab animals are considered. Subsequently, more information is provided on the genomics and genetics as well as the host immune response against *H. suis* infections. However, since only limited information is available for *H. suis* on these two subjects, other gastric *Helicobacter* spp. and in particular *H. pylori*, are considered here as well. Finally, a brief overview is presented on the association between *H. pylori* infections and extragastric manifestations, with emphasis on Parkinson's disease (PD).

1. The rise of the genus *Helicobacter*

In 1984, Marshall and Warren reported the presence of a typical curved bacterium in patients with active chronic gastritis, duodenal or gastric ulcers and highlighted its importance in the etiology of these diseases (Marshall and Warren, 1984). Initially this microorganism was believed to be a member of the genus *Campylobacter*. However, major differences in ultrastructural features, cellular fatty acids and the 5S and/or 16S rRNA gene sequences suggested that this bacterium did not belong to the genus *Campylobacter* but to a novel genus. This microorganism was therefore designated *Helicobacter pylori* (Goodwin et al., 1989). Since the description of this human pathogen, many other species in the genus *Helicobacter* have been described. Several of those possess the ability to colonize animal hosts and some of them have a zoonotic potential in humans (Haesebrouck et al., 2009; Flahou et al., 2016). The *Helicobacter* bacteria can roughly be divided into gastric and enterohepatic species. Gastric *Helicobacter* species are able to survive the hostile, acidic environment in the stomach by expressing urease at a high level (Weeks et al., 2000; Pot et al., 2007). Enterohepatic helicobacters thrive on the mucosal surfaces of the intestinal tract and/or the liver (Sterzenbach et al., 2007). Currently the *Helicobacter* genus comprises 46 identified species. An overview of the different species, their hosts, disease associations and zoonotic potential is shown in **Table 1**.

Table 1: *Helicobacter* spp. and their pathogenic significance for humans and animals (adapted from Flahou et al., 2016).

Group	Taxon	Natural host	Disease associations	Zoonotic potential
gastric <i>Helicobacter</i> spp.	' <i>Candidatus H. bovis</i> '*	cattle	ND	yes
	' <i>Candidatus H. homininae</i> '*	chimpanzee, gorilla	ND	unknown

	<i>H. acinonychis</i>	cheetah, tiger, lion	severe chronic gastritis	unknown
	<i>H. ailurogastricus</i>	cat	asymptomatic, gastritis, peptic ulcers, mucosa-associated lymphoid tissue lymphoma (MALT)-lymphoma	unknown
	<i>H. baculiformis</i>	cat	ND	unknown
	<i>H. bizzozeronii</i>	cat, dog	asymptomatic, gastritis, peptic ulcers, MALT-lymphoma	yes
	<i>H. cetorum</i>	whale, dolphin	asymptomatic, gastritis, peptic ulcers	unknown
	<i>H. cynogastricus</i>	dog	ND	unknown
	<i>H. felis</i>	dog, cat, cheetah, new guinea wild dog, rabbit	asymptomatic, gastritis, peptic ulcers, MALT-lymphoma	yes
	<i>H. heilmannii</i>	dog, cat, cheetah, bobcat, tiger, lynx, leopard, puma	asymptomatic, gastritis, peptic ulcers, MALT-lymphoma	yes
	<i>H. mustelae</i>	ferret	asymptomatic, gastritis, peptic ulcers, gastric cancer, MALT-lymphoma	unknown
	<i>H. pylori</i>	human	asymptomatic, gastritis, peptic ulcers, gastric cancer, MALT-lymphoma	/
	<i>H. salomonis</i>	dog, cat, rabbit	asymptomatic, gastritis, peptic ulcers, MALT-lymphoma	yes
	<i>H. suis</i>	pig, crab-eating macaque, rhesus macaque, mandrill monkey	asymptomatic, gastritis, peptic ulcers, MALT-lymphoma	yes
enterohepatic <i>Helicobacter</i> spp.	' <i>Candidatus H. colifelis</i> '*	cat	ND	unknown
	<i>H. anseris</i>	goose	ND	unknown
	<i>H. aurati</i>	hamster	asymptomatic, hepatic and intestinal disease	unknown
	<i>H. bilis</i>	mouse, rat, Mongolian gerbil, dog, cat, sheep	asymptomatic, hepatic and intestinal disease	yes
	<i>H. brantae</i>	goose	ND	unknown
	<i>H. callitrichis</i>	marmoset	ND	unknown
	<i>H. canadensis</i>	bird, pig	ND	yes
	<i>H. canis</i>	dog, cat	asymptomatic, diarrhea, hepatitis, gastroenteritis	yes

	<i>H. cholecystus</i>	hamster	asymptomatic, hepatic and intestinal disease	unknown
	<i>H. cinaedi</i>	hamster, cat, dog, rat, rhesus monkey, baboon	asymptomatic, hepatic and intestinal disease	Yes
	<i>H. equorum</i>	horse	ND	Unknown
	<i>H. fennelliae</i>	dog	asymptomatic, enteritis, proctitis, proctocolitis	Yes
	<i>H. ganmani</i>	mouse	asymptomatic, hepatic and intestinal disease	Yes
	<i>H. hepaticus</i>	mouse, Mongolian gerbil	asymptomatic, hepatic and intestinal disease	Yes
	<i>H. macacae</i>	baboon, rhesus monkey	asymptomatic, chronic colitis, intestinal adenocarcinoma	Unknown
	<i>H. marmotae</i>	cat, woodchuck	asymptomatic, hepatitis, typhlocolitis	Unknown
	<i>H. magdeburgensis</i>	mouse	asymptomatic, hepatic and intestinal disease	Unknown
	<i>H. mastomyrinus</i>	rodents	asymptomatic, hepatic and intestinal disease	Unknown
	<i>H. mesocricetorum</i>	hamster	asymptomatic, hepatic and intestinal disease	Unknown
	<i>H. muricola</i>	mouse	asymptomatic, hepatic and intestinal disease	Unknown
	<i>H. muridarum</i>	rat, mouse	asymptomatic, hepatic and intestinal disease	Unknown
	<i>H. pamatensis</i>	pig, cat, bird	ND	Yes
	<i>H. pullorum</i>	poultry	asymptomatic, hepatitis, diarrhea, enteritis	Yes
	<i>H. rappini</i>	sheep, dog, mouse	ND	Yes
	<i>H. rodentium</i>	rat, mouse	asymptomatic, hepatic and intestinal disease	Unknown
	<i>H. saguini</i>	cotton-top tamarin	asymptomatic, ulcerative colitis, typhlocolitis and dysplasia	Unknown
	<i>H. suncus</i>	house musk shrew	ND	Unknown
	<i>H. trogonum</i>	sheep, pig, mouse	asymptomatic, hepatic and intestinal disease	Unknown
	<i>H. typhlonius</i>	rat, mouse	asymptomatic, hepatic and intestinal disease	Unknown

	<i>H. valdiviensis</i>	wild birds	ND	Unknown
	<i>H. westmeadii</i>	human	asymptomatic, inflammatory bowel disease	/
	<i>H. winthamensis</i>	human, wild rodents	asymptomatic, gastroenteritis	Yes

ND: Not described

*: *Helicobacter* species that has not yet been isolated and cultivated *in vitro*

2. Gastric *Helicobacter* species in humans with emphasis on zoonotic pathogens

H. pylori is the best studied and most prevalent of the *Helicobacter* species which colonize the human stomach. Infections with this microorganism have not only been associated with gastritis and peptic ulcer disease as described above, but also with more severe gastric pathologies including mucosa-associated lymphoid tissue (MALT-) lymphoma and adenocarcinoma (Stolte et al., 1993; Kusters et al., 2006). *H. pylori* infections are frequently acquired during early childhood and the disease progresses with age (Suerbaum & Josenhans, 2007). Most of the *H. pylori* infections remain asymptomatic. Only 10-20% of the people carrying *H. pylori* will develop gastric disorders (Kusters et al., 2006; Suerbaum and Josenhans, 2007). **Figure 1** provides an overview of the different *H. pylori*-associated gastric disease outcomes in human patients.




Figure 1: Sequential steps of gastric disease resulting from *H. pylori* infection. Depending on the localization of the *H. pylori*-induced chronic gastritis, no gastric disease, duodenal ulcers, gastric ulcer disease or cancer may develop. In 85% of the patients, an infection with *H. pylori* will not result in the development of gastric diseases. In 10-15% of the *H. pylori*-infected patients, gastritis is mainly localized in the antrum (antrum predominant gastritis), which results in increased acid secretion

and successively in the development of duodenal ulcers. In only 2-5% of the patients, *H. pylori* infection will lead to the development of atrophic gastritis in the corpus of the stomach. This latter pathology is accompanied by decreased acid secretion as a result of atrophy of the acid producing glands, thereby increasing the risk to develop gastric cancer. The precancerous cascade progresses from multifocal atrophic gastritis, through intestinal metaplasia, to dysplasia, the severe grade represents carcinoma *in situ*, then invasive adenocarcinoma. Chronic gastritis is associated with MALT-lymphoma. Gastric ulcer occurs with corpus gastritis, before the stage of achlorhydria. Depending on the virulence of the *H. pylori* strain, the environmental factors and the host immune response, infections with this bacterium will or will not result in the development of gastric adenocarcinoma (Karczewska et al., 2014).

It is estimated that half of the human population in the world is infected with *H. pylori* albeit with large geographical variations. In developing countries, more than 80% of the human population is infected with *H. pylori*, whereas the prevalence rate of *H. pylori* in developed countries varies from 20 to 50% (Kusters et al., 2006; Pounder and Ng, 1995). This drop in prevalence in the Western world may create a niche for colonization of the human stomach by other organisms (Pounder and Ng 1995; Kusters et al., 2006).

Indeed, gastric disease in humans has also been associated with other, long, spiral-shaped bacteria. These gastric non-*H. pylori* *Helicobacter* spp. (NHPH) are similar in morphology to bacteria earlier reported in the stomach of pigs, cats, dogs and non-human primates (Salomon, 1898; Mendes et al., 1990; Queiroz et al., 1990). Generally, they have been found in 0.2 to 6% of gastric biopsies from humans with severe gastric complaints (Heilmann and Borchard, 1991; Stolte et al., 1994; Svec et al., 2000; Solnick et al., 2003). NHPH include *H. suis* from pigs and non-human primates and *H. felis*, *H. bizzozeronii*, *H. salomonis* and *H. heilmannii* from cats and dogs (Haesebrouck et al., 2009; Flahou et al., 2016). Although *H. cynogastricus*, *H. baculiformis* and *H. ailurogastricus* also have pets as natural host, their zoonotic significance currently remains unknown. Besides these long spiral-shaped helicobacters, 3 other animal-associated gastric *Helicobacter* spp., with a slightly-curved morphology more similar to that of *H. pylori*, are known: *H. acinonychis* from wild large felines, *H. cetorum* from marine mammals and *H. mustelae* from ferrets (Paster et al., 1991; Harper et al., 2002; Eaton et al., 1993). Infections of humans with those agents have not been described.

2.1 Phylogeny of animal-associated gastric *Helicobacter* species

Although minor differences in morphology among the porcine, feline and canine gastric NHPH have been described (Haesebrouck et al., 2009; Smet et al., 2011; Joosten et al., 2015), species identification based on microscopic analysis of human biopsy samples is not recommended (Eaton et al., 1996). Therefore, analysis of the *Helicobacter* 16S rRNA gene sequence has frequently been used in the past for differentiation between gastric *Helicobacter* species (O'Rourke et al., 2004b). However, sequencing of the 16S rRNA gene cannot distinguish between the canine and feline helicobacters, including *H. bizzozeronii*, *H. felis*, *H. salomonis*, *H. baculiformis*, *H. cynogastricus* and *H. heilmannii* (Smet et al., 2011; Baele et al., 2008a; Van den Bulck et al., 2006; Dewhirst et al., 2005). For differentiation between those species, analyses of the *ureA* and *ureB* and the *hsp60* gene sequences have been performed. The similarity of their *ureAB* genes is lower than 85%, which is sufficient for discrimination between these species (O'Rourke et al., 2004b). It has recently been shown that the newly described *H. ailurogastricus* species from cats can not be distinguished from its closest relative *H. heilmannii* by means of its 16S rRNA and *ureAB* genes but by comparison of their complete genome sequences

(Joosten et al., 2015). This means that the discriminatory capacity of these gene sequences is not high enough for the distinction of closely related gastric *Helicobacter* species. In this respect, genome sequencing based approaches are superior compared to the traditional 16S rRNA sequence analysis for studying phylogeny because they are based on the complete genome content and have a better resolution for discriminating both distantly and closely related bacteria (Vandamme, 2014). **Figure 2** is an illustration of a phylogenetic tree based on the core genomes of all gastric helicobacters from which the genome sequence is publically available (Joosten et al., 2015).




Figure 2: Phylogenetic tree based on the aligned and concatenated core genes of *Helicobacter* species from which the genomic content is known. The phylogram represents a maximum-likelihood tree of gastric and enterohepatic *Helicobacter* spp. based on 303 aligned and concatenated core genes. The enterohepatic *Helicobacter* spp. (*H. cinaedi*, *H. hepaticus*, *H. winthamensis*, *H. canadensis*, *H. pullorum*, *H. billis*, *H. canis*) were used as an outgroup. Approximate likelihood ratio test values of >99% were obtained for all nodes. Optimization of topology, branch length and parameters of the starting tree were performed (adapted from Joosten et al., 2015).

2.2 Clinical significance of gastric non-*Helicobacter pylori* *Helicobacter* infections in humans

A number of reports have highlighted that people having close contact with dogs, cats and pigs are more likely to be infected with an animal-associated gastric *Helicobacter* species (Chung et al., 2014; De Bock et al., 2007; Joosten et al., 2013; Meining et al., 1998). *Helicobacter* DNA has been detected in saliva from cats, dogs and pigs, indicating that the oral cavity of these animals may act as source of NHPH infections for humans (Ekman et al., 2013; Casagrande Proietti et al., 2010; ShojaeeTabrizi et al., 2010). Fecal-oral transmission has also been suggested as a possible route for infection in cats (Ghil et al., 2009). Besides direct contact with animals, other routes of transmission of NHPH should not be neglected. It has been shown that *H. felis* is able to survive in water for several days highlighting the possible role for water in the transmission of this species as well (Azevedo et al., 2008). *H. suis* can be present on and survive in minced pork indicating that raw or undercooked pork may also constitute a source of *H. suis* infection for humans (De Cooman et al., 2013).

H. suis, *H. heilmannii*, *H. bizzozeronii*, *H. felis* and *H. salomonis* (**Table 1**) have been associated with gastritis, gastric and duodenal ulcers and low grade MALT-lymphoma in humans (Haesebrouck et al., 2009; Matsumoto et al., 2014; Iwanczak et al., 2012; Duquenoy and Le Luyer, 2009). They have not been associated so far with gastric adenocarcinoma. The gastritis induced by NHPH is usually less active and less severe compared to *H. pylori* but the risk to develop MALT-lymphoma is higher (Morgner et al., 2000; Haesebrouck et al., 2009). Clinical signs associated with NHPH infections in humans include: epigastric pain, nausea, hematemesis, irregular defecation frequency and consistency, vomiting, recurrent dyspepsia, heartburn, dysphagia and decreased appetite (Dieterich et al., 1998; Goddard et al., 1997; Heilmann and Borchard, 1991; Kaklikkaya et al., 2002; Mention et al., 1999; Oliva et al., 1993; Schildt et al., 2000; Seo et al., 2003; Sykora et al., 2003; Van Loon et al., 2003; Wegmann et al., 1991; Yang et al., 1998; Yoshimura et al., 2002). In patients undergoing endoscopy, a variety of lesions can be observed ranging from a normal to slightly hyperaemic mucosa, mucosal oedema, nodular inflammation and the presence of ulcerations in the antrum of the stomach or in the duodenum (Haesebrouck et al., 2009; Sykora et al., 2003; Van Loon et al., 2003; Yang et al., 1995; Yoshimura et al., 2002). Histological examination of gastric biopsies generally reveals infiltration of lymphocytes and plasma cells. In some cases, lymphocytic aggregates or intestinal metaplasia have been described (Ierardi et al., 2001; Joosten et al., 2013; Matsumoto et al., 2014; Morgner et al., 2000; Yakoob et al., 2012).

In several studies the prevalence of gastric NHPH was investigated by analyzing human biopsies from patients with gastric complaints (Trebesius et al., 2001; De Groote et al., 2005; Van den Bulck et al., 2005). In general, NHPH have been detected in 0.2–6% of these patients, with *H. suis* being the most prevalent NHPH species (Haesebrouck et al., 2009).

Treatment of gastric *Helicobacter* infections is recommended in case of *Helicobacter*-related symptoms or in order to prevent the evolution to atrophic gastritis and gastric cancer (Cats et al., 1998; Kusters and Kuipers, 2001). Like for *H. pylori* eradication, the treatment of a NHPH infection consists of the administration of a proton pump inhibitor (PPI) in combination with two antibiotics, such as clarithromycin and tetracycline or amoxicillin. However, failure of successful eradication of both *H. pylori* and NHPH infections has been reported

and may be due to an increasing occurrence of antimicrobial resistance (Kivisto et al., 2010; Megraud et al., 2004; Horiki et al., 2008; Graham et al., 2010). An overview of what is known so far for NHPH is given below.

Van den Bulck et al. (2005) studied the susceptibilities of animal-derived *H. felis*, *H. bizzozeronii*, and *H. salomonis* isolates to 10 antimicrobial agents. All species were highly susceptible to ampicillin, clarithromycin, tetracycline, tylosin, enrofloxacin, gentamicin, and neomycin. This was demonstrated by low Minimum Inhibitory Concentrations (MICs). Higher MICs were obtained for lincomycin and spectinomycin. Two *H. felis* strains showed a MIC of 16 µg/ml for metronidazole, suggesting acquired resistance to this antimicrobial agent (Van den Bulck et al., 2005). Another study highlighted that *H. bizzozeronii* and potentially other NHPH easily acquire clinically relevant resistance to metronidazole due to the high mutation rate of the simple sequence cytosine repeat (SSCR) located in the 3' region of the oxygen-insensitive NAD(P)H-nitroreductase (*rdxA*) gene (Kondadi et al., 2013). For this reason, metronidazole, lincomycin and spectinomycin are considered as second line therapy. Vermoote et al. (2011) reported the presence of acquired resistance to fluoroquinolones in a porcine *H. suis* isolate. For all tested *H. suis* strains, the MIC of ampicillin was higher than described for *H. felis*, *H. bizzozeronii*, *H. salomonis* and *H. pylori*, possibly indicating intrinsic reduced susceptibility of *H. suis* to aminopenicillins compared to other gastric *Helicobacter* species. The significance of this finding for the treatment of *H. suis* infections is currently not known. However there is a report of a patient with a spiral gastric *Helicobacter* infection, which relapsed after triple therapy including amoxicillin and clarithromycin, and was then eradicated by quadruple therapy containing tetracycline and bismuth (Dobbs et al 2005). This organism was reported to be *H. heilmannii* by 16S rRNA based PCR, but was later shown to be *H. suis*.

2.3 Diagnosis of gastric *Helicobacter* infections

To date, several non-invasive and invasive diagnostic tools are available for the detection of gastric infections with *Helicobacter* in general and *H. pylori* in particular. An overview is given in **Table 2** and the most frequently used tests are described below in more detail.

Table 2: Overview of the available non- or minor invasive diagnostic tests for the identification of gastric *Helicobacter* spp.

Diagnostic test	Type	Identification	Advantages	Disadvantages	References
urea breath test	non-invasive	<i>H. pylori</i>	- simple and safe - high sensitivity/specificity in adults - initial screening - confirmation eradication	- variable sensitivity/specificity in children - variable sensitivity/specificity in case of partial gastrectomy and peptic ulcer bleeding	Malfertheiner et al., 2012; Guarner et al., 2010 ; Barkun et al., 2010 ; Velayos et al., 2012 ; Wardi et al., 2012
		gastric NHPH species	- simple and safe - initial screening	- false negative results - difficult use in animals - test results depending on colonization, speed of gastric emptying and recent use of antibiotics	Okiyama et al., 2005; Matsui et al., 2014 ; Blaecher et al., 2013 ; Cornetta et al., 1998
stool antigen test (SAT)	non-invasive	<i>H. pylori</i>	- safe, fast and reliable - use in children	- variable accuracy - reduced sensitivity in	Gisbert and Pajares, 2006; Saez et al., 2012;

			- use after distal gastrectomy	gastrointestinal bleeding - reduced accuracy in unformed or watery stool samples	Cardenas et al., 2008; Quieroz et al., 2013 ; Adamopoulos et al., 2009 ; Yan et al., 2010
		gastric NHPH species		- not available	
serology	non-invasive	<i>H. pylori</i>	- use for initial screening, patients using antibiotics, patients suffering from acute bleeding	- variability in sensitivity and specificity - variability in efficiency depending on <i>H. pylori</i> strain (heterogeneity between strains), geographical area and host genetic differences - no discrimination between past and present infection	Rubicz et al., 2011; Mégraud et al., 2012 ; Burucoa et al., 2013; Vale et al., 2008-2009 ; Vitoriano et al., 2011 ; Malfertheiner et al., 2007 ; Hirschl and Rotter, 1996 ; Choi et al., 2011 ; Herbrink and van Doorn, 2000
		gastric NHPH species		- not available	
enzyme-linked immunosorbent assay (ELISA)	non-invasive	<i>H. pylori</i>	- easy and reliable - rather high sensitivity and specificity - use for saliva and dental plaques	- use in young children (development adaptive immune system)	Bode et al., 2002; Raymond et al., 1996 ; Luzza et al., 1997 ; De Pascalis et al., 1999 ; Sunnerstam et al., 1999 ; Malaty et al., 2000
		gastric NHPH species		- not available	
polymerase chain reaction (PCR)	non-invasive	<i>H. pylori</i>	- use for fecal samples - use for dental plaques, saliva	- fecal samples: false negative results as a result of food degradation products, PCR- inhibitory factors and overload of bacterial species	Teoman et al., 2007; Souto and Colombo, 2008
		gastric NHPH species	- fecal samples: decent image on the presence of enterohepatic <i>Helicobacter</i> spp.	- fecal samples: false negative results as a result of food degradation products, PCR- inhibitory factors and overload of bacterial species	Shinozaki et al., 2002
	Invasive	<i>H. pylori</i>	- identification and quantification in gastric biopsies - high sensitivity and specificity - identification of virulence factors	- variation as a result of transport conditions and number of specimens	Qin et al., 2016; Schabereiter-Gurtner et al., 2004; Ndip et al., 2003; Ricci et al., 2007; De Martel et al., 2010
		gastric NHPH species	- identification and quantification in gastric biopsies - high sensitivity and specificity - differentiation between gastric NHPH species - strain differentiation	- variation as a result of transport conditions and number of specimens - aspecific interactions with host DNA and DNA from other NHPH species	Dieterich et al., 1998; De Groote et al., 2001; Hwang et al., 2002; Baele et al., 2004; O'Rourke et al., 2004; Trebesius et al., 2001; Matsumoto et al., 2009; Kivisto et al., 2010; Wuppenhorst et al., 2013; Joosten et al., 2013; Liang et al., 2013; Matsui et al., 2014
	Invasive	<i>H. pylori</i> and gastric NHPH species	- visualization of macroscopic abnormalities in the stomach	- not reliable - low sensitivity and specificity - no differentiation between gastric <i>Helicobacter</i> spp.	Haesebrouck et al., 2009; Kato et al., 2013; Oguri et al., 2007; Watanabe et al., 2012; Kusters et al., 2006;

rapid urease test (RUT)	Invasive	<i>H. pylori</i> and gastric NHPH species	- relatively high specificity and sensitivity - low cost and rapid diagnosis - semi-quantitative	- lower sensitivity in case of: peptic ulcer bleeding, partial gastrectomy and formalin contamination of forceps - false negatives in case of low grade colonization	Tseng et al., 2005; Gisbert et al., 2006; Ozaslan et al., 2010; Choi et al., 2012; Tian et al., 2012; Happonen et al., 1998
histology	Invasive	<i>H. pylori</i>	- evaluation of status gastric mucosa - cheap versus expensive staining	- reliability depends on localization and number of gastric biopsies - variability as a result of use of proton pump inhibitors, leading to false negative results	Wang et al., 2010; Aggarwal et al., 2011; El-Zimaity et al., 2013; Lash and Genta et al., 2013
		gastric NHPH species	- evaluation of status gastric mucosa - cheap versus expensive staining	- detection hampered by focal distribution of bacteria - not recommended for identification up to species level	Wang et al., 2010; Aggarwal et al., 2011; Lash and Genta et al., 2013; Happonen et al., 1998; Stoffel et al., 2000
isolation and cultivation	Invasive	<i>H. pylori</i>	- golden standard - specificity up to 100%	- variability in sensitivity depending on quality, quantity and transport conditions of gastric biopsies.	Ndip et al., 2003; Ricci et al., 2007; Kusters et al., 2006
		gastric NHPH species	- specificity up to 100%	- low sensitivity - low success rate - laborious procedure	Ndip et al., 2003; Ricci et al., 2007; Kusters et al., 2006; Happonen et al., 1998; Javala et al., 1998; Haesebrouck et al., 2009

2.3.1 Non- or minor invasive tests

2.3.1.1 The urea breath test

A frequently used test for the detection of gastric *Helicobacter* species is the urea breath test. This simple and safe non-invasive test has been reported to be highly sensitive (88%-95%) and specific (95%-100%) for the detection of *H. pylori* in adults, whereas in children a large variability in sensitivity and specificity, ranging from 75% up to 100% has been observed (Malfertheiner et al., 2012; Guarner et al., 2010). Because of the rather low sensitivity and specificity, the reliability of the urea breath test has likewise been questioned when applied after emergency endoscopy in peptic ulcer bleeding and in case of partial gastrectomy (Barkun et al., 2010; Velayos et al., 2012; Wardy et al., 2012). The principle of the urea breath test is based on the urease activity of gastric *Helicobacter* species. The presence of urease in the stomach breaks down the orally administered ^{13}C -labeled or radioactive ^{14}C -labeled urea into ammonia and soluble $^{13/14}\text{CO}_2$, which is absorbed by the gastric mucosa, transported by the systemic circulation and excreted in the exhaled breath. Finally the amount of exhaled $^{13/14}\text{C}$ is calculated by measurement of the ratio of $^{13/14}\text{C}$ to ^{12}C in the exhaled breath or the serum of the patient (Logan, 1998; Savarino et al. 1999; Graham and Klein, 2000; Gisbert and Pajares, 2004). An increase over baseline, 30 minutes after urea ingestion, more than a specified value constitutes a positive result. In general, the use of the urea breath test is recommended in case of initial diagnosis of *Helicobacter* and for confirmation of *Helicobacter* eradication after treatment (Malfertheiner et al., 2012). Up to date, limited information is available on the sensitivity and specificity of the urea breath test for the detection of NHPH

infections in humans. Although all gastric NHPH species are characterized by their urease activity (Haesebrouck et al., 2009), false negative results are often obtained when the urea breath test is performed in NHPH-infected human patients (Okiyama et al., 2005; Matsui et al., 2014). Possible explanations might be the patchy distribution of these microorganisms in the human stomach (Haesebrouck et al., 2009) and the relative low colonization density (Matsui et al., 2014).

2.3.1.2 Stool antigen test (SAT)

The stool antigen test (SAT) is a non-invasive diagnostic tool based on the detection of *H. pylori* antigens in human fecal samples (Gisbert and Pajares, 2006). This test is frequently applied for the screening of *H. pylori* in children (Cirak et al., 2007; Choi et al., 2011; Chehter et al., 2013). Although SATs are considered to be safe, fast and reliable for the identification of *H. pylori* infections (Gisbert et Pajares, 2006), the accuracy is influenced by antigenicity of the *H. pylori* strains as well as by the time between sample collection and measurements (Ritchie et al., 2009; Queiroz et al., 2013). A decreased sensitivity is found in patients suffering from gastrointestinal bleeding, whereas unformed or watery stool samples can result in a reduced accuracy because of the diluting effect on the *H. pylori* antigens (Saez et al., 2012). Two types of highly sensitive and specific SATs are available: the immunochromatography assay (ICA) and the enzyme immune assay (EIA). As ICA-tests are easy to apply and do not require any specialized equipment, they are frequently used for rapid diagnosis (Shibayama et al., 2011). They are, however, less reliable than the EIA- tests (Gisbert et al., 2006; Shibayama et al., 2011; Sato et al., 2012; Korkmaz et al., 2013). The ICA tests are often presented as a test-strip (eg. Testmate rapid pylori antigen). The presence of *H. pylori* antigens in a suspended stool sample results in the formation of immune complexes with color-labeled monoclonal antibodies. Migration of these antigen-antibody complexes over the surface of strip results in binding with secondary antibodies, which is visualized by the appearance of a colored line on the test-strip (Suzuki et al., 2002). In EIA tests the optical density is measured after incubation and washing of a mixture of suspended stool and peroxidase-conjugated *H. pylori* antibodies (Suzuki et al., 2002). Except for controversial results in post-eradication assessment, EIAs based on polyclonal antibodies are reliable diagnostic tests for the detection of *H. pylori* infections (Odaka et al., 2002; Veijola et al., 2005). Nevertheless, monoclonal antibody based SATs generally represent a higher specificity compared to polyclonal based SATs (Dominguez et al., 2006; Deguchi et al., 2009) and they are considered more reliable than the urea breath test in children (Cardenas et al., 2008; Queiroz et al., 2013) and in patients who underwent distal gastrectomy (Adamopoulos et al., 2009; Yan et al., 2010). Currently, no antigen assays have been developed for the detection of gastric NHPH infections in human patients.

2.3.2 Invasive tests

2.3.2.1 Endoscopy and analysis of gastric biopsies

Endoscopy is frequently performed to evaluate the gastric mucosa for the presence of gross abnormalities. Endoscopic findings characteristic for an *H. pylori* infection are: diffuse redness, spotted redness, mucosal swelling, erosions and peptic ulcers (Haesebrouck et al., 2009; Kato et al., 2013). Diagnosis of *Helicobacter*

infections based solely on endoscopic findings is not reliable enough because of the low sensitivity and specificity. In order to obtain a more trustworthy diagnosis, gastric biopsy samples need to be further examined using histology, PCR techniques and/or *in vitro* cultivation. Biopsy is, of course, needed for gastric RUT. Here the biopsy is placed in a chamber containing urea and an indicator, which changes colour in presence of ammonia. Several studies have indicated that the corpus greater curve, in general, is the most optimal biopsy site for the diagnosis of *H. pylori* infections (Calvet et al. 2013; El-Zimaity et al., 2013). It may not be appropriate in antral or pan gastritis.

2.3.2.1.1 Histology

In general, histological examination of gastric biopsies is considered as a standard method for the detection of an *H. pylori* infection. However, the reliability depends on the location of biopsy collection in the stomach and the number of biopsies (El-Zimaity et al., 2013). Histology has the advantage that additional information can be obtained on the status of the gastric mucosa, such as the presence of acute or chronic inflammation, lymphoid aggregates, intestinal metaplasia and glandular atrophy (Lash and Genta, 2013).

For identification of *H. pylori*, the haematoxylin eosin (HE) staining is, in terms of cost-effectiveness, the most designated staining (**Figure 3**) (Wang et al., 2010). Other stainings that have been applied for the detection of *H. pylori* in gastric biopsies, brush cytology and impression smear cytology specimens are: Warthin-Starry-staining, Gram-staining, Genta-staining, modified Giemsa-staining, toluidine-blue-staining and Romanowski-staining (Warren and Marshall, 1983; Mendoza et al. 1993; Mirsa et al., 1993). Although *H. pylori* visualization is also possible using immunohistochemical stainings, application of these specialized, more expensive stainings is only justified for analysis of biopsies derived from patients with chronic active and inactive gastritis and for biopsies collected to confirm *Helicobacter* eradication therapy (Wang et al., 2010; Aggarwal et al., 2011). Like for *H. pylori*, diagnosis of infections with other gastric *Helicobacter* species in human biopsies is also routinely performed by histology and cytology (Debongnie et al. 1995). However, detection of NHPH infections is hampered by the focal distribution of these bacteria and therefore infection with these agents is often missed (Haesebrouck et al., 2009).




Figure 3: HE staining of the gastric mucosa of *Helicobacter*-infected human patients. (A) HE staining of the gastric mucosa of a *H. pylori*-infected human patient. The slightly curved *H. pylori* bacteria, visualized within the black circle, are mainly

located at the surface of the gastric epithelial cells or in the mucus layer in the stomach (adapted from Batts et al., 2013), (B) HE staining of the gastric mucosa of a *H. suis*-infected human patient. These spiral-shaped bacteria (within the black circle) are more frequently found in the gastric pits (adapted from Joosten et al., 2013).

2.3.2.1.2 Polymerase chain reaction

The polymerase chain reaction (PCR) can be used as a diagnostic tool to screen gastric biopsies for the presence of *Helicobacter* DNA. Moreover, using real-time (RT)-PCR, quantitation of bacteria in gastric biopsies is possible as well. In *H. pylori*-positive patients with severe precancerous lesions, detection of the vacuolating cytotoxin (*VacA*) gene in gastric biopsies using PCR has been proven really valuable (Schabereiter-Gurtner et al., 2004). PCR assays for the detection of gastric NHPH species are mainly based on the amplification of a part of the *16S rRNA* gene or the *ureA/B* genes (De Groote et al., 2001; Hwang et al., 2002; Baele et al., 2004; O'Rourke et al., 2004). Recently, a study was published describing the development of a PCR based on the *H. suis*-specific *carR* gene for identification of *H. suis* infections (Matsui et al., 2014). Despite the development of different PCR assays for the detection of NHPH in general and *H. suis* in particular, no information on their sensitivity and specificity is available.

2.3.2.1.3 Culture

Another method for the identification of gastric *Helicobacter* infections by analyses of gastric biopsies is isolation and cultivation of *Helicobacter*. Generally, *Helicobacter* grows under micro-aerobic conditions (5% O₂, 10% CO₂ and 85% N₂) at 37°C and at a pH ranging between 5.5 and 8 (Scott et al., 2002; Stingl et al., 2002; Kusters et al., 2006; Haesebrouck et al., 2009). The specificity for the detection of *H. pylori* infections by cultivation reaches up to 100% (Ndip et al., 2003), whereas significant variability in sensitivity has been described, depending on the quality, the quantity and the transportation conditions of the gastric biopsies (Ricci et al., 2007). Because of the high specificity and relative high sensitivity, cultivation is considered the golden standard for the diagnosis of *H. pylori* infections (Kusters et al., 2006). Compared to *H. pylori*, isolation and cultivation of gastric NHPH species has a low success rate due to the very fastidious nature of these bacteria (Happonen et al., 1998; Javala et al., 1998). So far, only *H. felis* (Wüppenhorst et al., 2012) and *H. bizzozeronii* (Andersen et al., 1999; Kivistö et al., 2010; Schott et al., 2011) have successfully been isolated from the human stomach.

2.3.3 Conclusion

Apart from histology, no commercial tests are currently available for the diagnosis of infections with NHPH in general and *H. suis* in particular. Therefore the development of accurate and reliable diagnostic techniques for the identification of gastric NHPH infections is required to obtain a credible picture on the prevalence of NHPH in humans.

3. *H. suis*: a porcine pathogen with a zoonotic potential for humans

3.1 Prevalence of *H. suis* in pigs

In general, a prevalence rate of 60% has been described for *H. suis* infections in pigs at slaughter age in Europe, Asia and South- and North-America (Barbosa et al., 1995; Grasso et al., 1996; Cantet et al., 1999; Roosendaal et al., 2000; Choi et al., 2001; Hellemans et al., 2007; Kopta et al., 2010). It has been suggested that *H. suis* is transmitted between pigs via the oral-oral route through saliva or via the gastric-oral route through vomiting or regurgitation, but this requires further investigation (Haesebrouck et al., 2009; Flahou et al., 2016).

A very high prevalence has been reported in adult pigs (> 80%). Lower infection rates have been shown in younger animals ranging from 2% in suckling piglets (Hellemans et al., 2007), 25 % in piglets after weaning (Hellemans et al., 2007) and 76 % in slaughter pigs of approximately 6 months old (Bosschem et al., 2016). This remarkable increase in prevalence of *H. suis* after weaning may suggest the presence of protective antibodies in the sow milk (Melnichouk et al., 1999; Hellemans et al., 2007). The protective effect of maternal milk against the colonization of *H. pylori* in the human and mouse stomach has also been described (Corthésy-Theulaz et al., 2003; Bhuiyan et al., 2010). In young piglets and animals at slaughter age, *H. suis* is mainly localized in the antrum and the pylorus region of the stomach, whereas this bacterium is mainly located in the fundus of the stomach in adult pigs (Hellemans et al., 2007; De Bruyne et al., 2012).

3.2 Pathogenic significance of *H. suis* in its natural hosts and experimentally infected laboratory animals

3.2.1 *H. suis* infections in pigs

Experimental and natural *H. suis* infections in pigs have been associated with chronic gastritis in the antrum of the stomach. Inflammation is mainly characterized by the infiltration of lymphoid cells and the development of lymphoid follicles (Mendes et al., 1991, Grasso et al., 1996, Queiroz et al., 1996; Park et al., 2000; Hellemans et al., 2007; De Bruyne et al., 2012). Besides the strong association with gastritis, infection with *H. suis* has also been associated with ulceration of pars oesophagea of the stomach (**Figure 4**) (Barbosa et al., 1995; Queiroz et al., 1996; Roosendaal et al., 2000; Choi et al., 2001), although *H. suis* does not colonize this area of the stomach. The exact role of *H. suis* in the development of this pathology remains to be further elucidated. *H. suis* infection in pigs has been associated with an increased number of gastrin producing cells and a decreased number of somatostatin producing cells in the antrum of the stomach (Sapierzynski et al., 2007). Since gastrin stimulates and somatostatin inhibits the secretion of hydrochloric acid by parietal cells, this may influence gastric acid production, which may also be altered due to the tropism of this bacterium for parietal cells (Hellemans et al., 2007). It has been suggested that such an altered gastric acid secretion might be involved in the development of ulceration of the pars oesophagea (Sapierzynski et al., 2007; Haesebrouck et al., 2009). Other research groups could not find a link between a *H. suis* infection and ulceration of the pars oesophagea (Grasso et al., 1996; Melnichouk et al., 1999; Park et al., 2000; Szeredi et al., 2005). Differences in virulence between *H. suis* strains, different sampling and laboratory techniques might explain these contradictory results.

Besides *H. suis*, other factors, such as particle size of the feed, the presence of short fatty acids in the stomach, the occurrence of concurrent diseases or changes in the microbiome, may also contribute to the development of gastric ulceration (**Figure 4**) (Hessing et al., 1992; Argenzio et al., 1996; Ayles et al., 1996; Krakowka et al., 1998; Lindberg et al., 2001; Robertson et al., 2002; Amory et al., 2006; Miller et al., 2012). The presence of ulcers in the porcine non-glandular mucosa of the stomach may result in decreased feed intake, a decrease in daily weight gain and even sudden death due to fatal hemorrhage (Ayles et al. 1996; Haesebrouck et al. 2009), thus leading to significant economic losses (**Figure 4**). A decrease in daily weight gain of up to 10 % has been observed in weaned piglets (6-8 weeks old) after 4 and 6 weeks of experimental infection with *H. suis*, although without a clear association with the development of ulceration in the non-glandular part (De Bruyne et al., 2012).




Figure 4: Schematic overview of the contributing factors and the possible successive effects of ulcerations in the stomach of pigs. Apart from an infection with *H. suis*, the development of gastric ulceration has also been associated with a broad range of factors, such as particle size of the feed, the presence of short fatty acids in the stomach, concurrent diseases and changes in the microbiome. The presence of ulcers in the stomach of pigs may result in decreased feed intake, decreased daily weight gain and fatal hemorrhage. This figure was adapted from Baele et al., 2008 and Haesebrouck et al., 2009.

3.2.2 *H. suis* infections in non-human primates:

Besides in pigs, *H. suis* is also found in the stomach of non-human primates, including cynomolgus monkeys, mandrill monkeys and rhesus macaques (Haesebrouck et al., 2009; Flahou et al., 2016; O'Rourke et al., 2004; Nakamura et al., 2007; Martin et al., 2013; Matsui et al., 2014). This microorganism is mainly localized at the

surface epithelium, the gastric pits and superficial glands of the non-human primate stomach (O'Rourke et al., 2004). Further research is required to determine the exact pathogenic significance of *H. suis* infections in non-human primates (Haesebrouck et al., 2009; Flahou et al., 2016).

3.2.3 *H. suis* infections in experimentally infected laboratory animals

Specific pathogen free (SPF; free of *Helicobacter* spp.) inbred C57BL/6 and BALB/c mice have been shown to be useful models for the study of *Helicobacter*-related gastric disease (O'Rourke and Lee, 2003; Rogers and Fox, 2004). Generally, in experimentally infected mice, inflammation caused by *H. suis* is characterized by the infiltration of mononuclear and polymorphonuclear cells in the lamina propria mucosae and/or the tunica submucosa (**Figure 5A-B**) (Flahou et al., 2010; Bosschem et al., 2016). Infiltration of macrophages can be visualized from 3 weeks post-infection onwards. At 9 weeks post-infection, an augmentation in the number of neutrophils and (CD3⁺) T-cells can be observed as well (Flahou et al., 2010). At 8 months post-infection large lymphoid aggregates were observed in the stomach of *H. suis*-infected rodents (**Figure 5C**) (Flahou et al., 2010). Apart from inflammation, *H. suis* infections in mice have also been shown to cause necrosis of parietal cells and an increase in mucosal epithelial cell proliferation (Flahou et al., 2010; O'Rourke et al., 2004; Nakamura et al., 2007). Furthermore, gastric lymphoid nodules, hyperplasia of the gastric mucosa and mucus metaplasia, have been described as well after long term (>18 months) infection of mice with *H. suis* (O'Rourke et al., 2004; Park et al., 2008; Matsui et al., 2014). C57BL/6 and BALB/c mice genetically are predominant Th1 and Th2 responders, respectively (Flahou et al., 2010). Differences in gastric inflammation and colonization after *H. suis* infection have been found in these mouse strains. A more pronounced gastric inflammation has been described in the BALB/c mice compared to C57BL/6 mice after infection with *H. suis*, whereas a higher colonization density (both in antrum and fundus) has been noted in *H. suis*-infected C57BL/6 mice compared to *H. suis*-infected BALB/c mice.

Mongolian gerbils are yet another useful rodent model to study *Helicobacter*-related gastric pathology. In these animals, inflammation and colonization is more confined to the antrum and a narrow zone at the forestomach-stomach transition zone, which resembles the antrum predominant colonization by NHPH in humans. In addition, lesions observed in Mongolian gerbils infected with *H. suis* develop faster and are more pronounced than those in mice (Flahou et al., 2010; Bosschem et al., 2016). This includes the relatively fast development of severe inflammation-related pathologies, such as gastric MALT-lymphoma-like lesions. They already appear from 9 weeks post-infection onwards (**Figure 5D**) (Flahou et al., 2010; Bosschem et al., 2016). Humans suffering from a NHPH gastritis have also been shown to develop gastric MALT-lymphoma more frequently than those suffering from *H. pylori* gastritis (Stolte et al., 2002). These findings indicate that the outbred Mongolian gerbil may be more suitable than mice to study interactions of *H. suis* with its human host. Unfortunately, less molecular tools are available for the study of *Helicobacter*-related gastric pathology in gerbils compared to mice.




Figure 5: Images of HE stainings of the stomach of mice (A-C) and a Mongolian gerbil (D). **(A)** HE staining of the fundus of C57BL/6 mouse 4 weeks after inoculation of the growth medium of *H. suis* (control mouse). The arrow points to the normal composition and architecture of the mucosa of the fundus, characterized by no or negligible infiltration of mononuclear and polymorphonuclear cells in the lamina propria mucosae and tunica submucosa; **(B)** HE staining of the fundus of C57BL/6 mouse 4 weeks after infection with a porcine *H. suis* strain. The arrow appoints the infiltration of mononuclear and polymorphonuclear cells in the lamina propria mucosae and tunica submucosa; **(C)** HE staining of the fundus of BALB/c mouse 8 month after infection with a porcine *H. suis* strain. The black arrows show the two large lymphoid aggregates in the lamina propria mucosae as a result of major infiltration of mononuclear and polymorphonuclear cells in the lamina propria mucosae in combination with a loss of the normal mucosal architecture. **(D)** HE staining of the antrum of a *H. suis*-infected Mongolian gerbil at 8 months post-infection. The arrow shows the infiltration of mononuclear cells in the tunica muscularis (adapted from Flahou et al., 2010).

3.2.3 *H. suis* infections in man

H. suis is considered to be the most prevalent gastric NHPH species in humans (Haesebrouck et al., 2009). Infections with this zoonotic bacterium have been shown to comprise 13.9-78.5% of all diagnosed human gastric NHPH infections (Trebesius et al., 2001; Van den Bulck et al., 2005; Haesebrouck et al., 2009). It is not exactly known how *H. suis* is transmitted from pigs to humans, but direct and indirect contact with pigs and pork carcasses, as well as the consumption of raw or undercooked pig meat have been considered risk factors (Meining et al., 1998; Van den Bulck et al., 2005; Haesebrouck et al., 2009; Joosten et al., 2013; De Cooman et al., 2013). Similar to other gastric NHPH infections, infections with *H. suis* in humans have been associated with gastric disease and clinical signs may include epigastric pain, nausea, hematemesis, vomiting, irregular defecation frequency and consistency, recurrent dyspepsia, heartburn and decreased appetite (Dieterich et al.,

1998; Goddard et al., 1997; Heilmann and Borchard, 1991; Kaklikkaya et al., 2002; Mention et al., 1999; Oliva et al., 1993; Schildt et al., 2000; Seo et al., 2003; Sykora et al., 2003; Van Loon et al., 2003; Wegmann et al., 1991; Yang et al., 1998; Yoshimura et al., 2002; Dobbs et al., 2005). A recent case report described an *H. suis* infection in a pig veterinarian suffering from reflux esophagitis and general dyspeptic symptoms (Joosten et al., 2013). Diagnosis in this patient was performed using *ureA*-based PCR on gastric biopsy samples and sequencing of the PCR amplicons. Additionally, histological examination revealed the presence of spiral-shaped bacteria in the lumen of the gastric glands as well as the presence of chronic, diffuse gastritis, characterized by a pronounced infiltration of lymphocytes and macrophages (Joosten et al., 2013). Apart from gastritis, *H. suis* infections in humans have also been associated with gastric and duodenal ulcers and MALT-lymphoma (Haesebrouck et al., 2009; Matsumoto et al., 2014). Treatment of human *H. suis* infections generally consists of the administration of amoxicillin, clarithromycin and a proton pump inhibitor (Joosten et al., 2013). However in case of therapy failure, the administration of tetracycline might be recommended (Dobbs et al., 2005).

Due to the extremely difficult diagnosis, the knowledge on human *H. suis* infections is rather limited. Further research is recommended in order to obtain more insights on the pathogenesis and clinical significance of *H. suis* infections in human patients.

3.3 Use of genomics and genetics to obtain better insights in the pathogenesis of *H. suis* infections

Comparison of the recent available whole genome sequences of NHPH, including *H. suis*, with that of *H. pylori* has revealed the presence of genes encoding homologues of *H. pylori* virulence factors involved in acid acclimation, motility, chemotaxis, oxidative stress resistance and gastric epithelial cell damage (Arnold et al., 2011; Schott et al., 2011; Vermoote et al., 2011; Smet et al., 2013; Joosten et al., 2015). An overview is shown in **Table 3**.

Table 3: Overview of NHPH genes encoding homologues of *H. pylori* virulence factors and their role in the pathogenesis of *Helicobacter* infections.

NHPH gene homologues encoding <i>H. pylori</i> virulence factors	Proteins	Function
<i>ureA/B/I/E/F/G/H</i>	- UreA and UreB - urease accessory proteins (UreEFGH) - Urel	- acid resistance - regulation of urease activity - acid activated urea channel
<i>che, tlp</i>	- chemotaxis proteins (<i>che</i>) - methyl-accepting chemotaxis proteins (<i>tlp</i>)	Chemotaxis
<i>katA, sodB, Ahpc</i>	- catalase (KatA) - superoxide dismutase (SodB) - alkyl hydroperoxide reductase (HhpC)	oxidative stress resistance
<i>napA</i>	- neutrophil-activating protein (NAP)	oxidative stress
<i>Ggt</i>	- γ -glutamyl transpeptidase	induction of gastric lesions (apoptosis-inducing and immunosuppressive)
<i>fldA</i>	- flavodoxin protein (FldA)	electron acceptor of the pyruvate oxidoreductase enzyme complex
<i>porA/B/C/D</i>	- pyruvate oxidoreductase complex (Por)	electron donor for FldA
<i>pgbA/B</i>	- plasminogen-binding protein (Pgb)	induction of gastric lesions
<i>ansB</i>	- L-asparaginase II	cell-cycle inhibition of fibroblasts and gastric cell lines

3.3.1 pH homeostasis

Like all gastric *Helicobacter* spp., the genome of *H. suis* contains genes encoding the urease enzyme, which consists of two subunits (*UreA* and *UreB*) and plays a role in acid resistance by converting urea into ammonia and carbon dioxide (Eaton et al., 1991; Burn and Chen, 2000). By neutralizing the hydrochloric acid, the ammonia induces an elevated pH in the stomach (Eaton et al., 1991). Additionally, ammonia reduces the viability of the gastric epithelial cells (Smoot et al., 1990). The second urease gene clusters consists of the urease accessory *ureEFGH* genes and the *ureI* gene, encoding for the UreEFGH and the Urel proteins, respectively (Akada et al., 2000). The Urel protein forms an acid-activated urea channel, thereby controlling the transport of urea into the bacterial cell (Scott et al., 2000). In order to be functional, two nickel molecules (Ni^{2+}) should be present within the metallocenter of the *H. pylori* urease (Ge et al., 2013). Together the UreEFGH proteins are responsible for the incorporation of nickel, thereby regulating the urease activity (Cussac et al., 1992; Mobley et al., 1995).

3.3.2 Chemotaxis and motility

Genes coding for proteins involved in the chemotactic behavior (*che* and *tlp* genes) are typical for gastric helicobacters (Arnold et al., 2011; Schott et al., 2011; Vermoote et al., 2011; Smet et al., 2013; Joosten et al., 2015). They allow bacterial movement towards urea, cholesterol, arginine and bicarbonate and away from hydrochloric acid (Foyne et al., 2000; McGee et al., 2005; Terry et al., 2005; Lertsethtakarn et al., 2011). Moreover, the presence of the TlpB receptor, which is required for pH-taxis, is also needed for the colonization of *Helicobacter* spp. (O'Toole et al., 1994; Kim et al., 1999).

All gastric *Helicobacter* spp. are able to migrate to the epithelial cell surface of the stomach and towards a more neutral pH using flagellae (Salama et al., 2013; Arnorld et al., 2011; Schott et al., 2011; Vermoote et al., 2011, Smet et al., 2011). The flagellae are composed of a body and flagellar filaments, consisting of two flagellins A (FlaA) and B (FlaB), a flagellar hook (FlgE) and a flagellar cap (FlgD) (Josenhans et al., 1995; Andrutis et al., 1997; O'Toole et al., 1994; Kim et al., 1999).

Both flagellum dependent motility and chemotaxis are important for successful colonization of the stomach (Moens and Vanderleyden, 1996; Otteman et al 1997).

3.3.3 Oxidative stress

Catalase (*KatA*), superoxide dismutase (*SodB*) and alkyl hydroperoxide reductase (*AhpC*) allow gastric helicobacters to counteract oxidative stress induced by the host immune response (Seyler et al., 2001; Harris et al., 2002; Olczak et al., 2003; Ernst et al., 2005, Schott et al., 2011; Smet et al., 2011; Arnold et al., 2011; Vermoote et al., 2011). The neutrophil-activating protein (NAP) is another virulence factor present in all gastric *Helicobacter* spp., which is involved in oxidative stress resistance (Arnold et al., 2011; Schott et al., 2011; Vermoote et al., 2011; Smet et al., 2013; Joosten et al., 2015). Additionally it induces the production of reactive oxygen radicals (ROS) (Olczak et al., 2003; de Bernard et al., 2010; Evans et al., 1995) and it is responsible for chemotaxis of neutrophils and monocytes, increased adhesion of neutrophils, activation of monocytes, mast cells and neutrophils as well as increasing the lifespan of monocytes and neutrophils (Satin et al., 2000; Montemurro et al., 2001; 2002; Amedei et al., 2006; Polenghi et al., 2007; Cappon et al., 2010).

3.3.4 Outer membrane proteins

Adhesion plays a crucial role in the initial colonization of the stomach by *Helicobacter* (McGuckin et al., 2011). Approximately 64 outer membrane proteins (OMPs) have been identified in the genome of *H. pylori*. Their role in the colonization of the stomach has been studied thoroughly (Alm et al., 2000; Odenbreit et al., 2009). These OMPs can be divided into 5 families. The first family consists of the *Helicobacter* outer membrane porin (Hop) and the Hop related (Hor) proteins. The second is the *Helicobacter* OMP family (Hof) protein family. The third is the *Helicobacter* outer membrane (Hom) protein family. The fourth is the iron-regulated OMPs family. The fifth is the efflux pump OMPs family (Oleastro and Ménard, 2013). Hop proteins include adhesins and porins, which play an important role in the adhesion of *H. pylori* to the gastric mucosa (Alm et al., 2000; Aspholm et al., 2006; Odenbreit, 2005; Gerhard et al., 2001). The two most important adhesins in *H. pylori* are the blood group antigen binding adhesin A (BabA) and the sialic acid binding adhesin A (SabA). BabA binds to human fucosylated Lewis b group antigen or related terminal fucose residues in blood group A, B and O antigens, expressed on Mucin 1 (MUC1) and Mucin 5AC (MUC5AC), whereas SabA binds to sialylated carbohydrate structures upregulated in the inflamed gastric mucosa (Boren et al., 1993; Ilver et al., 1998; Mahdavi et al., 2002).

Although ca. 55-60 OMPs have been identified in the genome of the gastric NHPH (Joosten et al., 2015; Alm et al., 2000), comparative genomics revealed that none of the genes encoding the *H. pylori* adhesins were present

within the NHPH genomes (Joosten et al., 2015). Genes encoding homologs of all members of the *H. pylori* Hof proteins appear to be present in all NHPH genomes, except for *H. mustelae* (Smet et al., 2014). A recent study showed that the *H. heilmannii* HofE and HofF play a role in the adhesion of this bacterium to gastric mucins and epithelial cells (Liu et al., 2016). Further research, however, is recommended in order to determine which other OMPs contribute to the adhesion process.

3.3.5 Virulence factors contributing to the gastric lesions

Major virulence factors of *H. pylori* involved in the induction of gastric lesions are the cytotoxin-associated gene pathogenicity island (*cagPAI*) and the vacuolating cytotoxin A (*VacA*). Both virulence factors are, however, absent in all NHPH with the exception of *H. cetorum* carrying a functional *VacA* toxin (Vermoote et al., 2011). *CagPAI* consists of 30 genes encoding a type IV secretion system (T4SS), which penetrates the epithelial cells of the stomach, allowing injection of the CagA protein into the host cells. Phosphorylation of this protein induces morphological changes in the epithelial cells by changes in the cytoskeleton and indirect activation of the pro-inflammatory nuclear factor (NF)- κ B (Akopyants et al., 1998; Segal et al., 1999; Asahi et al., 2000; Christie and Vogel, 2000; Odenbreit et al., 2000; Fisher, 2001; Yamazaki et al., 2003; Brandt et al., 2005; Mimuro et al., 2007; Backert et al., 2008). *VacA*, on the other hand, induces vacuolization of the epithelial cells and mitochondrial-mediated apoptosis. The toxicity of this protein depends on the formation of membrane channels, which are induced by this protein (Cover and Blaser, 1992; Szabo et al., 1999; Kimura et al., 1999; Willhite and Blanke, 2004).

Another virulence factor, which on the contrary is present in all gastric *Helicobacter* species is the gamma-glutamyltranspeptidase (GGT) (Arnold et al., 2011; Vermoote et al., 2011; Schott et al., 2011; Smet et al., 2013). For *H. pylori*, it has been shown that GGT causes damage to epithelial cells by inducing a depletion of glutamine and glutathione, both important for the maintenance of healthy gastrointestinal tissue (Shibayama et al., 2003; 2007). Additionally, it inhibits T cell proliferation and induces an upregulation of cyclo-oxygenase (COX)-2 and epithelial growth factor (EGF)-related peptide as well as apoptosis of human gastric adenocarcinoma (AGS) cells (Busiello et al., 2004; Schmees et al., 2007; Kim et al., 2007; 2010; Zhang et al., 2015). The contribution of the *H. suis* GGT in apoptotic and necrotic cell death of gastric epithelial cells has been described as well. By the disintegration of glutathione into pro-oxidant degradation products, *H. suis* GGT induces increased oxidative stress, resulting in lipid peroxidation and finally gastric epithelial cell death (Flahou et al., 2011).

Besides *VacA*, *CagPAI* and GGT, other virulence factors have been associated with gastric lesions. These include PutA, PutP, AnsB, FldA, PgbA, PgbB, HtrA, PrtC and IceA. Most of them are present in all gastric *Helicobacter* spp. (Arnold et al., 2011; Schott et al., 2011; Vermoote et al., 2011; Kersulyte et al., 2013; Smet et al., 2013; Joosten et al., 2015). These factors have been studied in *H. pylori*. L-Asparaginase II (AnsB) plays a role in the cell-cycle inhibition of fibroblasts and gastric cell lines (Scotti et al., 2010). FldA functions as an electron acceptor for the pyruvate oxidoreductase enzyme complex, whereas pyruvate oxidoreductase (Por) proteins function as electron donors for FldA. The presence of the FldA protein has been associated with MALT-lymphoma (Chang et al., 1999). The plasminogen-binding proteins (PgbA and PgbB) will allow *Helicobacter* to

bind to host plasminogen, which may contribute to the obstruction of the natural healing process of gastric ulcers (Vermooten et al., 2011). Furthermore, HtrA is involved in cleavage of E-cadherin, and thus contributes to the disruption of the gastric epithelial barrier (Hoy et al., 2010; Lower et al., 2011). By the production of collagenase (PrnC), gastric helicobacters are able to induce the degradation of collagen (Zhang et al., 2015). Finally, IceA has been associated with peptic ulcer disease and seems to be induced by contact between epithelial cells and *Helicobacter* (Peek et al., 1998; 2000; Donahue et al., 2000).

Although a substantial number of genes encoding putative virulence factors has been described in the NHPH genomes in general and the *H. suis* genome in particular, for the majority of these, their exact role in pathogenesis still remains unclear.

3.4 Host immune response to gastric *Helicobacter* infections

In general, *Helicobacter* causes a life-long gastric infection since the evoked host immune response is not able to eliminate *Helicobacter* from the stomach (Kusters et al., 2006; Flahou et al., 2012). Several studies describing the host immune response (both innate and adaptive host responses) against *H. pylori* are available, whereas only few reports have been published on the host immune responses against *H. suis* infections (Baele et al., 2008; Flahou et al., 2010; Flahou et al., 2012; Bosschem et al., 2016). The next section is therefore particularly focused on the innate and adaptive host immune response induced by *H. pylori* and where possible, information about *H. suis*-induced immune responses is provided as well.

3.4.1 Innate immune response to *H. pylori* and *H. suis*

Dendritic cells (DC), macrophages and gastric epithelial cells form the first barrier against infections with *H. pylori* (Dubois and Borén, 2007). Like other invading pathogens, *H. pylori* bacteria are recognized by the cells of the innate immune system via distinct conserved structures, also referred to as pathogen-associated molecular patterns (PAMPs). These PAMPs are not expressed by the host cells, allowing recognition of pathogens by the host immune response (Takeda et al., 2004; Peek et al., 2010). The host cells, on their account, feature pattern recognition receptors (PRRs) on the cytoplasmic and endosomal membranes as well as in the cytosol. These PRRs, including the surface-expressed Toll-like receptors (TLRs), are able to recognize the bacterial PAMPs. Specific interactions between PRRs and PAMPs result in the activation of intracellular signaling pathways inducing the transcription of various genes that are responsible for the production of inflammatory cytokines, chemokines, antigen-presenting molecules, etc (Peek et al., 2010). Gastric epithelial cells from *H. pylori*-infected human patients have been shown to express TLR4, TLR5 and TLR9 (**Figure 6A**) (Schmausser et al., 2004). TLR2 is generally expressed on the surface of intestinal and gastric epithelial cells (Cario et al., 2002; Smith et al., 2003) and in contrast to TLR4, TLR5 and TLR9, it is responsible for the recognition of multiple *H. pylori* PAMPs, e.g. lipopolysaccharide (LPS) and the *H. pylori* neutrophil-activating protein (HP-NAP) (Peek et al., 2010) (**Figure 6A**). Interaction of HP-NAP with TLR2 leads to the activation of NF- κ B, followed by the upregulation of pro-inflammatory cytokines interleukin (IL)-12, IL-23 and tumor necrosis factor (TNF) (Amedei et al., 2006). (**Figure 6A**).

Since *H. pylori* possesses 4 to 6 monopolar sheated flagella and TLR5 is known to recognize bacterial flagellin, TLR5 was assumed to be involved in the recognition of *H. pylori* flagellin (Eaton et al., 1996). However, full TLR5 activation requires the presence of conserved amino acids. Because of the lack of these molecules, it has been reported that *H. pylori* flagellin cannot be recognized by TLR5 (Andersen-Nissen et al., 2005) (**Figure 6A**).

Additionally to TLRs, gastric epithelial cells may express the cytosolic nucleotide-binding oligomerization domain (NOD), which is a cytosolic pathogen recognition molecule. By its involvement in the intracellular recognition of bacterial mucopeptides, derived from bacterial peptidoglycan, NOD1 plays an important role in the innate immunity against *H. pylori*. Using the type 4 secretion system (T4SS), *H. pylori* secretes its peptidoglycan into the cytoplasm of the gastric epithelial cells, resulting in the activation of NOD1 (Viala et al., 2004). *H. pylori*-induced NOD1 activation results in the activation of NF- κ B, leading to the production of pro-inflammatory cytokines and chemokines, eg CXCL-8 (IL-8), CXCL-2 (murine homologue of IL-8) and type 1 interferon (IFN) (Viala et al., 2004; Watanabe et al., 2011) (**Figure 6A**).

Furthermore, IL-8 production by gastric epithelial cells can also be mediated in a TLR/NOD1 independent manner by interaction with *H. pylori* OMPs (Oleastro & Menard, 2013) (**Figure 6A**). Via BabA and SabA adhesins, *H. pylori* is also able to bind to gastric mucins, such as MUC1, which also results in the secretion of IL-8 by the gastric epithelial cells (**Figure 6A**) (Sheng et al., 2013). Additionally, interaction between the gastric epithelial cells and other *H. pylori* adhesins, including HopH, outer inflammatory protein A (OipA), HomB and adherence-associated lipoprotein A/B (AlpA/AlpB), may also induce gastric epithelial IL-8 secretion (Oleastro & Menard, 2013; Sheng et al., 2013) (**Figure 6A**).

The production of chemokines by the epithelial cells of the stomach results in the recruitment of innate immune cells, eg. dendritic cells, monocytes, macrophages and neutrophils to the site of infection. They are considered to be the second stage of the innate immune response (Allen et al., 2000; 2001). Interaction between *H. pylori* and monocytes/macrophages results in the activation of both TLR2 and TLR4 signaling in these cells, leading to the secretion of inflammatory cytokines such as IL-1 β , IL-6, IL-8, IL-10, IL-12 and TNF (Maeda et al., 2001; Gobert et al., 2004; Mandell et al., 2004). On the other hand, DC signaling in response to an *H. pylori* infection is mainly mediated by TLR2 and to a lesser extent by TLR9 and TLR4, inducing the secretion of IL-1 β , IL-6, IL-10, IL-12 and TNF (Rad et al., 2007; 2009; Kim et al., 2013). The production of these pro-inflammatory cytokines by DC's and macrophages leads to the recruitment, differentiation and activation of CD4⁺ Th cells, which organize the adaptive immune response (Wilson & Crabtree, 2007). Furthermore, macrophages and neutrophils are responsible for phagocytosis, which is an important antibacterial innate defense mechanism. However, by its ability to survive within the phagosomes, *H. pylori* partially evades phagocyte-mediated killing (Allen et al., 2000; 2001). In addition, *H. pylori* induces the extracellular release of ROS by phagocytes, but the bacterium is capable of surviving this response through catalase and superoxide dismutase activity (Ramaraio et al., 2000).

Little data is available on the innate immune response against an infection with *H. suis*. A recent *in vivo* study reported infiltration of neutrophils and mononuclear cells in the mucosa and submucosa of *H. suis*-infected BALB/c mice (Bosschem et al., 2016) (**Figure 6B**). The same study described enhanced expression of porcine DC

activation markers (CD25, CD40 and CD80/86), decreased expression of major histocompatibility complex (MHC)-II presentation at the surface of porcine DCs and DC-triggered expansion of regulatory T cells (Tregs) upon stimulation with *H. suis* lysate. Finally, *H. suis* stimulation of murine DCs resulted in an increased secretion of IL-6 and IL-23, inducing the differentiation from naïve T cells to T helper 17 (Th17) effector cells (Bosschem et al., 2016) (**Figure 6B**). Since no information is currently available on *H. suis* PAMPs (**Figure 6B**), further research is required to identify the *H. suis* genes that contribute in the activation of the host immune response.

3.4.2 Acquired immune response to *H. pylori* and *H. suis*

In response to the cytokine production by the innate immune cells, naïve T CD4⁺ helper (T CD4⁺) cells will differentiate towards either T helper 1 (Th1) cells, T helper 2 (Th2) cells, Th17 or regulatory T (Treg) cells. An overview of the cytokines responsible for the differentiation of naïve T CD4⁺ cells into the different effector cells in combination with the secretion of cytokines by the different T effector cells is presented in **Table 4**.

Table 4: Overview of the cytokines responsible for the differentiation of naïve T CD4⁺ cells into the different T effector cells and their secreted cytokines.

	Th1 effector cells	Th2 effector cells	Th17 effector cells	Treg effector cells
cytokines for differentiation	- IL-12	- IL-6/ IL-4	- IL-6/ IL-23	- TGF-β
secreted cytokines	- IFN-γ/ IL-2/ TNF	- IL-4/ IL-5/ IL-6/ IL-9/ IL-10/ IL-13	- IL-17/ IL-21/ IL-22	- IL-10/ TGF-β
references	Romagnani et al., 2000; Larussa et al., 2015 ; Banchereau et al., 2000	Romagnani et al., 2000; Larussa et al., 2015 ; Banchereau et al., 2000	Larussa et al., 2015	Larussa et al., 2015

3.4.2.1 T helper cells

The T cell response evoked by *H. pylori* infections consists of the IFN-γ producing CD4⁺ T cell phenotype, indicating a Th1 response (D'Elia et al., 1997; Bamford et al., 1998; Peek et al., 2010; Larussa et al., 2015). Differentiation of the naïve T CD4⁺ helper cells into Th1 helper cells is induced by IL-12 (**Table 4**), produced and secreted by activated monocytes, macrophages and DCs (Guiney et al., 2003; Hafsi et al., 2004) (**Figure 6A**). The severity of *H. pylori*-associated gastric pathology mainly depends on the Th1 responses and the Th1 produced cytokines (Mohammadi et al., 1996; Smythies et al., 2000; Sommer et al., 2001). These findings have been confirmed by the reduced gastric pathology in mouse models deficient in Th1 cell development (Sommer et al., 2001) and IFN-γ^{-/-} mouse models (Smythies et al., 2000). Several *H. pylori* virulence factors have been associated with the stimulation of the Th1 cell response, including CagPAI and HP-NAP. *H. pylori* infections have also been associated with a Th2 response (Serrano et al., 2007; Robinson and Atherton, 2010), characterized by the secretion of multiple cytokines including IL4, IL6 and IL10 (**Table 4**)(Kusters et al., 2006; Smythies et al., 2000; Odenbreit et al., 2006). *H. pylori* strains inducing a Th2 immune response typically lack genes encoding

major virulence factors such as the CagPAI and the VacA and the BabA and SabA adhesins (Liu et al., 2014). Apart from the production of multiple cytokines (**Table 4**), activated Th2 cells are also responsible for the inhibition of phagocytic cells (Romagnani et al., 2000). Indeed, several reports suggest the protective role of Th2 cells against severe lesions by suppressing the *H. pylori*-induced inflammation (Fox et al, 2000; Smythies et al., 2000). Other reports could not confirm these findings (Saldinger et al., 1998; Garhart et al., 2003) and further research is required to elucidate the exact role of the Th2 response in *H. pylori* infections.

A Th2 host immune response has also been associated with an *H. suis* infection in mice (Park et al., 2008). This type of response was characterized by elevated expression levels of IL-10 and IL-4 (Flahou et al., 2012; Zhang et al., 2015) (**Figure 6b**), as well as by the formation of gastric MALT-lymphoma-like lesions (Greiner et al., 1997; Flahou et al., 2012). Gastric MALT-lymphoma is characterized by a strong proliferation of B lymphocytes which may be dependent on Th2-type cytokines. Since infection with NHPH in general and *H. suis* in particular results indeed in a higher risk for the development of MALT-lymphoma, it is more likely to induce a Th2 response rather than a Th1 response (Greiner et al., 1997; Knörr et al., 1999; Haesebrouck et al., 2009; Flahou et al., 2010; Flahou et al., 2012). On the other hand, infections with *H. suis* in gerbils result in increased expression of IFN- γ in the stomach, suggesting the induction of a Th1 response by this pathogen as well (Vermoote et al., 2012; 2013; Bosschem et al., 2015; Zhang et al., 2015) (**Figure 6B**). These results indicate that the *H. suis*-evoked immune response might be host dependent. However, further investigation is recommended.

Another important immune mechanism contributing to the development of *H. pylori*-induced gastritis is the Th17 response (Shiomi et al., 2008; Shi et al., 2010; Hitzler et al., 2012; Raghavan and Quiding-Järbrink, 2012; Gray et al., 2013; Larussa et al., 2015). Th17 cells secrete IL-17, which plays a role in the recruitment of neutrophils to the site of infection, as well as in the stimulation of IL-8 secretion by the gastric epithelial cells (Luzza et al., 2000; Mizuno et al., 2005) (**Figure 6A**).

An infection with *H. suis* has also been associated with a Th17 response. The presence of Th17 cells and/or increased levels of IL-17 in the stomach of *H. suis*-infected mice, Mongolian gerbils and humans has been reported in several studies (Luzza et al., 2000; Mizuno et al., 2005, Flahou et al., 2012; Vermoote et al., 2012; 2013; Zhang et al., 2015; Bosschem et al., 2015; 2016) (**Figure 6B**). However, the Th17 response upon *H. suis*-infected mice appeared to be significantly higher in BALB/c mice than in C57BL/6 mice (Flahou et al., 2012). This finding can be explained by the fact that the overall inflammatory response in BALB/c mice is generally higher compared to C57BL/6 mice (Flahou et al., 2012). Additionally, a remarkable negative correlation between *H. suis* colonization and the expression levels of IL-17 mRNA has been reported (Flahou et al., 2012; Vermoote et al., 2012; 2013; Bosschem et al., 2015). Indeed, the higher inflammatory response in BALB/c mice might induce a less favorable environment for *H. suis*, thereby resulting in a lower colonization density (Flahou et al., 2012).

An increased number of Treg cells in the gastric mucosa of *H. pylori*-infected human patients has been described, suggesting that *H. pylori* colonization could evoke proliferation and migration of Treg cells as well (Lundgren et al., 2005). FOXP3, a gene contributing to the development of Tregs, is expressed by the CD4⁺ CD25⁺T cells in the gastric and duodenal mucosa of *H. pylori*-infected patients (Lundgren et al., 2005) (**Figure**

6A). By the production of anti-inflammatory cytokines (IL-10 and TGF- β), Treg cells are able to limit the proliferation and function of other immune cells, which may result in decreased inflammation and tissue damage (Eaton et al., 2001; Raghavan and Quiding-Järbrink, 2012). Indeed, Treg responses in *H. pylori*-infected children resulted in the inhibition of the Th1 and Th17 responses leading to a less severe inflammation and ulceration of the gastric mucosa (Harris et al., 2008). Moreover, the number of Treg cells was considerably higher in *H. pylori*-infected patients without peptic ulceration compared to patients suffering from peptic ulcers (Robinson et al., 2008).

H. suis infections in mice and gerbils have also been associated with the induction of a Treg immune response and increased expression levels of IL-10 in the gastric mucosa (Vermoote et al., 2012; 2013; Zhang et al., 2015; Bosschem et al., 2015; 2016). Elevated expression of IL-10 and the FOXP3-gene were observed in *H. suis*-infected mice (Zhang et al., 2015) (**Figure 6B**). Additionally, a positive correlation between IL-10 expression levels and *H. suis* colonization density has been described in BALB/c mice (Vermoote et al., 2012; 2013; Flahou et al., 2012).

3.4.2.2 B-cells

It has been suggested that B cells may also contribute to the immunopathogenesis of *H. pylori* infections. Uncontrolled proliferation of B lymphocytes has been reported in response to the production of certain cytokines, like IL-2, after interaction between T cells and *H. pylori* antigens (Wotherspoon et al., 1991). A minority of *H. pylori*-infected patients develop gastric MALT-lymphoma, which mainly arise by the proliferation of B cells (Kusters et al., 2006). Moreover, complete MALT-lymphoma regression has been described in more than 80% of the patients after successful *H. pylori* eradication (Parsonnet and Isaacson, 2004; Peek et al., 2010). Compared to wild-type mice, a reduced colonization and more severe gastritis combined with the increased infiltration of CD4⁺ cells was detected in B cell deficient mice at 8 and 16 weeks post-infection, indicating that B cells may temper the effect of the CD4⁺ cells (Akhiani et al., 2004). Further research is recommended to uncover the exact role of B cells as well as the role of T cell-B cell interactions in the pathology of the *H. pylori*-evoked immune response (Peek et al., 2010).

Although the exact role of B cells in *H. suis* infections remains to be elucidated, previous studies suggested that they might play a role in the pathogenesis as well (Vermoote et al., 2012; 2013; Zhang et al., 2015) (**Figure 6B**). Increased numbers of B cells have been reported after a 6 month-infection of mice and Mongolian gerbils with *H. pylori* and *H. suis* (Zhang et al., 2015). Inoculation of C57BL/6 mice with an *H. suis* strain evoked a host immune response characterized by activation and infiltration of B cells and CD4⁺ cells, which were mainly located in the surrounding of gastric lymphoid follicles. Moreover, their numbers were positively correlated with the duration of *H. suis* infection (Yamamoto et al., 2011). Additionally, a pronounced upregulation of chemokine ligand 13 (CXCL13), also called B cell attracting chemokine 1 or B lymphocyte chemoattractant, along with infiltration and proliferation of B cells in the centers of lymphocytic aggregates/follicles has been reported in the stomach of *H. suis*-infected, but not *H. pylori*-infected mice and Mongolian gerbils (Zhang et al., 2015) (**Figure 6B**). In contrast, another study showed that *H. suis*-induced lymphoid follicles were not only

composed of B cells but also of CD4⁺ T helper cells and that the development of gastric lymphoid follicles could not be detected in *H. suis*-infected IFN- γ deficient mice, indicating that IFN- γ might play a role in the pathogenesis of *H. suis*-induced gastric lymphoid follicles (Mimura et al., 2011).




Figure 6: Comparison of *H. pylori*-induced and *H. suis*-induced innate and adaptive immune response. **(A)** Overview of the *H. pylori*-evoked host innate and adaptive immune response. Contact between epithelial cells and *H. pylori* or its bacterial products may lead to the activation of NF- κ B, which results in the secretion of pro-inflammatory chemokines and cytokines, including IL-8. IL-8 induces the recruitment of immune cells to the lamina propria of the gastric mucosa. Depending on their effector functions, activated immune cells secrete pro-inflammatory cytokines (including IL-1 β , TNF, IL-6, IL-12, and IFN- γ). Based on these cytokines, naïve T CD4⁺ cells will differentiate towards either Th1 cells, Th2 cells, Th17 or Treg cells. Depending on the type of T-helper cell, certain cytokines will be secreted. **(B)** Overview of the *H. suis*-evoked host innate and adaptive immune response. Contact between epithelial cells and *H. suis* or its bacterial products may lead to the activation of NF- κ B, which results in the secretion of pro-inflammatory chemokines and cytokines, including IL-8 and IL-1 β . In contrast to *H. pylori*, the virulence factors of *H. suis* that evoke the innate immune response remain unknown. Subsequently, the pro-inflammatory cytokines induce the recruitment of immune cells to the lamina propria of the gastric mucosa. Depending on their effector functions, activated immune cells secrete pro-inflammatory cytokines, such as IL-23 and IL-6. Based on these cytokines, naïve T CD4⁺ cells will differentiate towards either Th1 cells, Th2 cells, Th17 or Treg cells. Depending on the type of T-helper cell, certain cytokines will be secreted. In conclusion, only limited information on the *H. suis*-induced innate and adaptive host immune response is currently available and thus further research is required. IL: interleukin; TNF: tumor necrosis factor; TLR: Toll-like receptor; IFN: interferon; Th: T-helper cell; NOD: nucleotide-binding oligomerization domain; NF: nuclear factor; CXCL-13: chemokine ligand 13; MUC1: mucin 1; T4SS: Type 4 secretion system; LPS: lipopolysaccharide; TGF: Tissue growth factor; NAP: Neutrophil-activating protein.

4. Extragastric manifestations of *Helicobacter* infections with emphasis on Parkinson's disease (PD)

The bidirectional communication between the gastrointestinal tract and the brain via the gut-brain axis has been considered (Cryan et al., 2012; Mayer et al., 2014). Anatomically, the gut-brain axis consists of the central nervous system (CNS), the autonomic nervous system (ANS) and the peripheral nervous system (PNS) (also known as the enteric nervous system or “little brain”) (Caso et al., 2008; Konturek et al., 2011; Budzynski et al., 2014). Via the gut-brain axis, changes in stress and emotions are able to influence the gastrointestinal immune system, the mucosal inflammation and the composition of gut microbiota and thereby the functionality of the gastrointestinal tract (Budzynski et al., 2014). On the other hand, alterations in the gut microbiota and the function and/or coordination of the gastrointestinal tract have been suggested to play a role in the overall physical and emotional state of the host (Cryan et al., 2012; Budzynski et al., 2014; Mayer et al., 2014).

Infections with *H. pylori* have not only been associated with changes in the composition of the gastrointestinal microbiota (Engstrand et al., 2013; Budzynski et al., 2014), but also with disturbed gastric emptying (Miyaji et al., 1999). Additionally, *Helicobacter* spp. in general and *H. suis* in particular are able to cause damage to the gastric epithelial cell lining by inducing epithelial cell apoptosis as described above (De Bock et al., 2006; Kusters et al., 2006; Flahou et al., 2010). Furthermore, *Helicobacter* antigens were also detected in macrophages in the mesenteric lymph nodes during chronic infection in mice (unpublished results). Together, these findings suggest that the inflammatory effects caused by gastric helicobacters are not restricted to the stomach. At the start of this PhD, only one clinical association had been made between *H. suis* infections and extragastric disease (Dobbs et al., 2005). In contrast, *H. pylori* infections had been associated with several

extragastric manifestations. A concise overview is shown in **Table 5**. In this part of the thesis, only the possible link between *H. pylori* infections and Parkinson's disease (PD) will be described in more detail.

Table 5: Concise overview of *H. pylori*-associated extragastric disorders. (MS): multiple sclerosis; (AD): Alzheimer's disease; (CNS): central nervous system; (PD): Parkinson's disease; (IBD): inflammatory bowel disease.

Group	Disease	Concise disease description	Association with <i>H. pylori</i>	References
neurological manifestations	ischemic stroke	stroke mainly caused by occlusion of the carotic or cerebral vessels	positive	Yang et al., 2011; Alvarez-Arellano et al., 2014 ; De Bastiani et al., 2008; Zhang et al., 2008; Wang et al., 2012; Wong et al., 2014
	migraine headaches	severe headaches	positive	Deleu et al., 1998; Gervil et al., 1999; Gasbarrini et al., 1998; Yiannopoulou et al., 2007; Tunca et al., 2004 ; Hong et al., 2007
	Guillain-Barré syndrome	acute, inflammatory autoimmune neuropathy characterized by progressive weakening of the motoric functions, mainly commencing in the legs	positive	Suzuki et al., 2006; Alvarez-Arellano et al., 2014 ; Kountouras et al., 2005; Chiba et al., 2002
	multiple sclerosis (MS)	most prevalent inflammatory demyelinating disorder of the central nervous system (CNS)	negative	Wender, 2003; Li et al., 2007; Mohebi et al., 2013
	neuromyelitis optica	disease characterized by inflammation of the optic nerves and the spinal cord	positive	Li et al., 2009; Long et al., 2013
	Alzheimer's disease (AD)	neurodegenerative disorder characterized by synaptic loss and neuronal cell death as a result of the intra- and extracellular accumulations of β -amyloid deposits in the brain	positive	Selkoe, 2002; Kountouras et al., 2006; Honjo et al., 2009; Chang et al., 2013; Alvarez-Arellano et al., 2014; Testerman et al., 2014; Wong et al., 2014; Malaguarnera et al., 2004; Kountouras et al., 2009a; Roubaud-Baudron et al., 2012 ; Kountouras et al., 2010
	Parkinson's disease (PD)	neurodegenerative disorder characterized by progressive loss of dopaminergic neurons, mainly in the substantia nigra of the brain	positive	Altschuler, 1996; Tsolaki et al., 2015; Bjarnason et al., 2005; Dobbs et al., 2010; Alvarez-Arellano et al., 2014 ; for review see Dobbs et al., 2016
endocrine manifestations	diabetes	endocrine disorder characterized by a disturbance in the glucose metabolism	positive	Simon et al., 1989, Oldenburg et al., 1996; Quadri et al., 2000; Gulcelik et al., 2005; Bener et al., 2007; Longo-Mbenza et al., 2007; Devrajani et al., 2010; Zhou et al., 2013; Jeon et al., 2012
	dyslipidemia	manifestation characterized by disturbances in the production and clearance of plasma lipoproteins	positive	Laurila et al., 1999; Hoffmeister et al., 2001; Majka et al., 2002; Takashima et al., 2002; Chimienti et al., 2003; Papamichael et al., 2009; He et al., 2014; Feingold et al., 1992; Chimienti et al., 2003
	autoimmune thyroid disorders	autoimmune mediated impairment of the thyroid gland	positive	Papamichael et al., 2009; Smyk et al., 2014; Testerman et al., 2014; Bassi et al., 2010; Bassi et al., 2012; Shi et al., 2013; Tomb et al., 1997; Bassi et al., 2010; Bertalot et al., 2004

hematological manifestations	iron deficiency anemia	anemia as a result of a lack of iron often caused by gastrointestinal bleeding.	positive	Monzon et al., 2013; Blecker et al., 1991; Bruel et al., 1993; Dufour et al., 1993 ; Milman et al., 1998; Peach et al., 1998; Collett et al., 1999 ; Yip et al., 1997; Kang et al., 2011; Musumba et al., 2012
	idiopathic thrombocytopenic purpura	autoimmune hematological disease characterized by low platelet counts and mucocutaneous bleedings caused by the autoimmune destruction of platelets	positive	Gasbarrini et al., 1998; Hino et al., 2003; Huber et a., 2003; Stasi et al., 2005; Suzuki et al., 2005; Campuzano-Maya, 2014 ; Franceschi et al., 2004; Takahashi et al., 2004
cardiopulmonary manifestations	cardiovascular manifestations, such as atherosclerosis and coronary artery disease	narrowing of the lumen of the blood vessels	positive	Mendall et al., 1994; Huang et al., 2011; Park et al., 2011; Vafaeimanesh et al., 2014; Kowalski et al., 2001; 2002; Oshima et al., 2005; Tamer et al., 2009; Chimienti et al., 2003; Kucukazman et al., 2009; Franceschi et al., 2002
	asthma and allergies	hypersensitivity reaction associated with a chronic Th2 response	negative	Amedei et al., 2010; Testerman et al., 2014; Wong et al., 2014; Chen et al., 2008; Zhou et al., 2013; Wang et al., 2013; Strachan, 1989; 2000
hepatic and biliary manifestations	non-alcoholic fatty liver disease	a group of liver disorders ranging from non-alcoholic fatty liver to the development of non-alcoholic steatohepatitis in which insulin resistance plays a role	positive	Rabelo-Gonçalves et al., 2015; Waluga et al., 2015; Cindoruk et al., 2008; Pirouz et al., 2009; Dogan et al., 2013; Abenavoli et al., 2013
	hepatic fibrosis, cirrhosis	disease characterized by replacement of liver cells by extracellular matrix proteins, including collagen, as a result of chronic liver diseases	positive	Rabelo-Gonçalves et al., 2015; Waluga et al., 2015; Goo et al., 2009; Ki et al., 2010; Ponzetto et al., 2000a; b; Feng et al., 2014; Esmat et al., 2012; Stalke et al., 2001
	hepatic encephalopathy	complication of liver cirrhosis characterized by the occurrence of neuropsychiatric symptoms and the excessive ammonia levels	positive	Amodio et al., 2004; Schulz et al., 2014
	hepatocellular adenocarcinoma	primary malignant liver tumor	positive	Rabelo-Gonçalves et al., 2015; Waluga et al., 2015; Avenaud et al., 2000; Nilsson et al., 2001; Dore et al., 2002; Verhoef et al., 2003; Pellicano et al., 2004; Huang et al., 2004
intestinal manifestations	inflammatory bowel disease (IBD)	IBD, an intestinal disorder of unknown pathogenesis, is considered a multifactorial manifestation mainly resulting from a complex interaction between environmental factors and gut microbiota in genetically susceptible humans	negative	Buchner et al., 2011; Papamichael et al., 2014; Ruuska et al., 1994; Thia et al., 2008; Luther et al., 2010; Rokkas et al., 2015; Tursi, 2006; Guiney et al., 2003; Chen et al., 2008
dermatologic manifestations	chronic spontaneous urticaria	dermatological manifestation of unknown etiology characterized by the spontaneous occurrence of itchy wheals lasting for at least 6 weeks	positive	Zuberbier et al., 2009; Magen et al., 2014; Smyk et al., 2014; Testerman et al., 2014, Wong et al., 2014; Federman et al., 2003; Fukuda et al., 2004; Magen et al., 2007; Akashi et al., 2011; Chiu et al., 2013
	rosacea	chronic skin disorder characterized by persistent	positive	Rebora et al., 1994; Güreter et al., 2002; Powell, 2005 ; Argenziano et al., 2003;

		central facial erythema		Bakar et al., 2007; Mashimo et al., 2006
	psoriasis	autoimmune skin disorder characterized by increased proliferation and less differentiation of the keratinocytes in the epidermis	positive	Magen et al., 2014; Smyk et al., 2014; Testerman et al., 2014; Qayoom et al., 2003; Ali et al., 2008; Martin Hübner et al., 2008, Onsun et al., 2012
	Behçet's disease	autoimmune multisystemic inflammatory vascular disorder featured by uveitis, skin lesions, genital ulcers and recurrent oral aphthous ulcers	positive	Magen et al., 2014; Avci et al., 1999; Pay et al., 2007
ocular manifestations	glaucoma	progressive ocular neuropathy is caused by apoptosis of retinal ganglion cells, leading to modifications of the optical nerve and vision loss	positive	Kountouras et al., 2001; 2002; 2003; Zavos et al., 2012; Tsolaki et al., 2015; Ritch, 2007; Sacca et al., 2014; Davies et al., 1994
	central serous chorioretinopathy	this visual impairment is caused by the leakage of fluid in the subretinal space as a result of the break-down of blood-retina barrier consisting of retinal pigment epithelium	positive	Wang et al., 2008; Casella et al., 2012; Giusti, 2004; Cotticelli et al., 2006; Casella et al., 2012; Dang et al., 2013; Rahbani-Nobar et al., 2011 ; Sacca et al., 2014
autoimmune systemic rheumatic manifestations	Sjögren syndrome	chronic inflammatory autoimmune disorder characterized by lymphoid cell infiltration and destruction of endocrine glands, specifically lacrimal and salivary glands	positive	Hasni et al., 2011; Radic et al., 2014; Aragona et al., 1999; Showji et al., 1996
	rheumatoid arthritis	autoimmune inflammatory disorder with symmetric destructive polyarthritis	negative	Hasni et al., 2011; Radic et al., 2014; Showji et al., 1996; Tanaka et al., 2005; Meron et al., 2010; Janssens et al., 1992
	systemic lupus erythematosus	autoimmune multisystemic inflammatory disorder affecting multiple organ systems such as skin, joints, kidneys, brain, etc	negative	Showji et al., 1996; Sawalha et al., 2004; Hasni et al. 2011, Radic et al., 2014; Yamanishi et al., 2006

4.1 PD: clinical outcome

4.1.1. History and pathological findings

PD was first described in 1817 by James Parkinson as the “shaking palsy”, a disorder characterized by tremor, muscle rigidity and poverty and slowness of movement (hypo- and bradykinesia) (Parkinson, 1817) and is the second most common neurodegenerative disorder in the current human population (Wirdefeldt et al., 2011; Mhyre et al., 2012). This neurological manifestation is characterized by a degeneration of the dopamine producing neurons in the pars compacta of the substantia nigra (in the basal ganglia of the brain) (**Figure 7**), thereby inducing dystrophic striatal projections (Wirdefeldt et al., 2011; Mhyre et al., 2012; Kannarkat et al., 2013). Inflammatory processes, characterized by oxidative stress, accumulation of cytokines and activation of microglia cells, have been suggested to contribute to the degeneration of the nigrostriatal pathway and

accelerate the disease progression, especially in case of idiopathic PD, a syndrome of unknown etiology (Tansey et al., 2007; Dobbs et al., 2008). Indeed, neuro-inflammation was more pronounced in the pons, striatum, substantia nigra, and frontal and temporal cortical regions in patients with early PD compared to age-matched control patients (Gerhard et al., 2006), and chronic use of non-steroidal anti-inflammatory drugs (NSAID) has been suggested to lower the risk of PD development (Tansey et al., 2007). Additionally, PD is considered as a chronic neuro-inflammatory disorder characterized by the occurrence of Lewy bodies, which are mainly localized in the surviving neurons. Lewy bodies are formed by the accumulation of intracellular protein aggregates, comprised mainly of misfolded α -synuclein. They are identified as eosinophilic inclusion bodies in neurons (Gibb et al., 1988; McNaught et al., 2003; Wirdefeldt et al., 2011; Mhyre et al., 2012; Kannarkat et al., 2013). Furthermore, an abnormal mitochondrial morphology, characterized by a long, thin appearance, has been observed within the enterocytes of the duodenum of PD patients, whereas impaired functionality has been reported in nigral and platelet mitochondria (Dobbs et al., 2008; Dobbs et al., 2016). Mitochondria are responsible for the production of ROS as a byproduct of molecular oxygen consumption in the electron transport chain. Additionally, these subcellular organelles are targets for cytokines, such as TNF, resulting in the overproduction of ROS by the mitochondria. Finally, cell death may occur as a result of oxidative stress, caused by the intracellular accumulation of ROS (Tansey et al., 2007).




Figure 7: Comparison between a healthy substantia nigra and a diminished substantia nigra in PD, characterized by a loss of dopamine producing neurons.

4.1.2 Motor and non-motor symptoms

This progressive neurodegenerative disorder leads to a broad spectrum of motor and non-motor symptoms. In general, PD-associated motor symptoms, such as resting tremor, muscle rigidity, brady- and hypokinesia and stooped posture are commonly known (Parkinson, 1817; Wirdefelt et al., 2011; Kannarkat et al., 2013). However, degeneration of the majority of the dopamine producing neurons (up to 80%) in the substantia nigra is a prerequisite for the development of motoric symptoms in PD patients (Kordower et al., 2013). In the majority of the patients (60-70%), resting tremor is the first indicative symptom for PD that appears. Generally the tremor has an unilateral and distal character (*e.g.* hand, fingers, foot, forearm), but over time it tends to expand to the contralateral upper limb or ipsilateral lower limb (Wirdefelt et al., 2011; Mhyre et al., 2012). Rigidity has a rather unilateral character and is generally more pronounced in the flexor muscles than in the extensor muscles. Bradykinesia is characterized by decreased moving speed in combination with slower response in movement changes and reduced fine motor skills. Hypokinesia is defined as a decreased frequency and amplitude of the movements, such as reduced stride length, reduced arm swing etc. The typical stooped posture of PD patients is still absent in early PD and is caused by changes in body position and gait as a result of the increased rigidity (Mhyre et al., 2012).

This age-related disease has also been associated with other, non-motor symptoms: 30-40% of the PD patients are diagnosed with dementia, apathy, anxiety, psychosis and depression, 80-90% suffer from sleeping disorders and up to 90% of the PD patients display an altered olfactory perception (Mhyre et al., 2012). Autonomic disorders, including cardiovascular, gastrointestinal and genitourinary disorders, have also frequently been reported in PD patients. In 30-58% of the PD patients orthostatic hypotension can be detected as a result of cardiovascular autonomic dysfunction (Goldstein, 2006). Genitourinary disorders, such as bladder dysfunction, decreased libido and impaired sexual performances, have been described in 40-80% of the PD patients (Winge et al., 2006; Sakakibara et al., 2008; Mhyre et al., 2012). Upper gastrointestinal tract disorders, like oropharyngeal dysphagia and esophageal motility impairment, are generally present in up to 70% to 80% of the PD patients. A major complication of upper gastrointestinal tract dysfunction is food aspiration (25-30% of the patients), possibly resulting in aspiration pneumonia. Bloating, nausea and altered absorption of medication in PD patients is related to a delayed gastric emptying. Furthermore, compared to control patients the occurrence of infrequent bowel contractions, potentially leading to obstipation, megacolon and volvulus, is clearly higher in patients suffering from PD. James Parkinson noted constipation as a feature in his essay (Parkinson 1817). In PD, frequency of defaecation diverges from that of controls three decades before the median age of neurological diagnosis (Charlett et al., 1997). Morphological and neurochemical changes of PD are found throughout the enteric nervous system and in dorsal vagal nuclei (Dobbs et al., 2008). Indeed, the presence of infrequent bowel movements is considered a risk factor for the development of PD (Abbott et al., 2001; Pfeiffer et al., 2003; 2011; Mhyre et al., 2012).

4.1.3 Incidence and prevalence

Studies in the US and Europe have reported a PD incidence between 12-19 cases/100.000 inhabitants per year (Twelves et al., 2003; Hirtz et al., 2007). Pringsheim and colleagues showed that the prevalence of PD in patients between 70 and 79 years old differed strongly depending on the geographical region: in North-America, Europe and Australia a prevalence of 1601 cases/100000 inhabitants has been reported, whereas a prevalence of 646 cases/100000 inhabitants has been reported in Asia. Furthermore, the same study observed that the prevalence of PD increased with age (Pringsheim et al., 2014). Another study reported that PD affects up to 1% of the population at 65-70 years of age, increasing to 4-5% of the population at the age of 85 (Fahn, 2003). Since age is considered the single greatest risk factor for the development of PD and the life expectancy of Western population is generally increasing, the number of PD cases is expected to increase (Dorsey et al., 2007). Compared to women, the prevalence of PD is 1.4 times higher in men (Twelves et al., 2003; Hirtz et al., 2007; Wirdefelt et al., 2011).

4.1.4 Etiology

Up to date the exact etiology of PD remains undetermined, however it is considered a multifactorial disease in which both genetic and environmental factors can play a role (Wirdefelt et al., 2011; Mhyre et al., 2012; Kannarkat et al., 2013). In 5-10% of the PD patients, the disease is the result of specific gene mutations, such as mutations in the α -synuclein-encoding gene (*PAK1/4*, *SNCA*), in the Leucin-Rich Repeat Kinase 2 (*LRRK2*)-encoding gene (*PARK8*, *LRRK2*), in the *parkin* (*PARK2*) locus and the Phosphatase and Tensin Homolog (*PTEN*)-induced Putative Kinase 1 (*PINK1*) encoding gene (Wirdefelt et al., 2011; Mhyre et al., 2012; Kannarkat et al., 2013). Mutations in the α -synuclein encoding gene and the *LRRK2* gene have been associated with dominantly inheritable parkinsonism, whereas mutations in *PINK* and *parkin* encoding genes have been associated with recessively inheritable parkinsonism (Tansey et al., 2007).

The majority of the PD cases are classified as sporadic/idiopathic Parkinsonism (IP), since the etiology remains unknown. However, different studies have identified several lifestyle and environmental risk factors for the development of IP. Generally, a negative association between smoking and coffee drinking on the one hand and the development of PD on the other hand has been reported (Wirdefelt et al., 2011; Mhyre et al., 2012). Additionally, a negative correlation between the occurrence of PD and physical activity, alcohol consumption and dietary intake of antioxidants has also been suggested. However, further research is recommended to confirm these associations. Besides, protracted pesticide exposure and the consumption of dairy products are considered as risk factors for the development of PD (Wirdefelt et al., 2011). Interestingly, some studies have mentioned the positive association between PD and rural living, well water consumption and farming. However, these findings were not confirmed by other researchers (Wirdefelt et al., 2011). This association was initially declared by the repeated protracted exposure to agrochemicals (Wirdefelt et al., 2011). However, a study in 2002 showed that the mortality from PD among farmers from 26 US states was clearly elevated among live-stock farmers, but was lower than in non-farmers and arable farmers (Lee et al., 2002).

4.2 Pathogenesis of PD: dysfunction of neural barriers

In PD, neuro-inflammation is believed to contribute to neurodegeneration with activation of microglia, the residential macrophages of the central nervous system (Mhyre et al., 2012). Peripheral inflammation can activate microglia, in particular when the brain barriers are disturbed (Alvarez-Arellano et al., 2014). The brain is protected by four specialized brain barriers: the blood-brain barrier (BBB), the blood-cerebrospinal fluid barrier (blood-CSF barrier), the blood-retina barrier and the arachnoid barrier. These barriers determine what will (and what will not) enter the brain, and account for many physiological properties of their specific neural environments (De Bock et al., 2014). The focus in this PhD thesis lays on the BBB and the blood-CSF barrier. In the next section, information will be provided on these two neural barriers (BBB and the blood-CSF barrier) and their role in PD.

4.2.1 Blood-brain barrier

The BBB is located at the level of the capillaries of the CNS and is responsible for the regulation of the fluid exchange between the systemic blood circulation and the CNS. This neural barrier is formed by capillary endothelial cells and supported by surrounding astrocytes and pericytes (**Figure 8**), also referred to as the neurogliovascular unit (Ramsauer et al., 2002). Beside their supporting function of the endothelial cells, astrocytes play a role in the regulation of the expression of several tight junctions (Wolburg et al., 1994).




Figure 8 : Illustration of the BBB, consisting of capillary endothelial cells, astrocytes and pericytes (adapted from De Bock et al., 2014).

In order to prevent paracellular diffusion of different molecules, including toxic agents and cytokines from the systemic circulation to the brain, capillary endothelial cells are connected by tight junctions and adherens junctions (**Figure 9**) (Ek et al., 2006). Tight junctions, also referred to as zonulae occludentes (ZO), are formed by multiple transmembrane claudins in which occludin is incorporated (Furuse et al., 1998a, Furuse et al., 1998b). Claudin-3, -5 and -12 play an important role in the connection between the capillary endothelial cells at the BBB (Nitta et al., 2003; Wolburg et al., 1994; Coisne et al., 2011). In contrast, occludin, which mainly has a supportive function, is not strictly necessary for the formation of the tight junctions (Furuse et al., 1998b; De

Bock et al., 2014). The tight junctions are mainly anchored to the actin (**Figure 9**) of the cytoskeleton of the endothelial cells by ZO proteins, ZO-1 and ZO-2 (De Bock et al., 2014). The adherence junctions are mainly formed by transmembrane cadherin proteins. This calcium-dependent adhesion molecule is responsible for cell-cell contact and it can be associated with other γ -catenines, such as β -catenine. In turn these molecules are connected with the cytoplasm of the cell by α -catenin (Vleminckx et al., 1999; Corada et al., 1999; Ek et al., 2006).




Figure 9 : Illustration of the tight junctions and adherens junctions of the blood-brain barrier. The tight junctions are formed by multiple claudins in which occludin is incorporated. In turn, occludin and claudin interact with the actin cytoskeleton via adaptor proteins, including the zonulae occludens proteins. The adherence junctions are formed by transmembrane proteins belonging to the cadherin family and are linked to actin via the catenins. ZO: zonulae occludens; TJ: tight junction; AJ: adherence junction (adapted from De Bock et al., 2014).

Disturbance of the BBB has been reported in patients suffering from PD (Stolp et al., 2009; Nielsen et al., 2012; Lee et al., 2014). Both local and systemic inflammation have been suggested to play a role in the increased permeability of the BBB. As a result of signaling through the endothelial cells, several locally or systemically produced cytokines, such as $IL-1\beta$, $IL-6$ and $TNF-\alpha$, may cause alterations in the tight junction structure, thereby inducing an increased BBB permeability. The increased BBB permeability along with elevated levels of cytokines and inflammatory mediators, such as ROS, may cause neuro-inflammation and oxidative stress, leading to further BBB disruption and brain damage (Stolp et al., 2009; Lee et al., 2014). Furthermore, migration of white blood cells through the BBB as a result of the central inflammatory response has been suggested as well (Stolp et al., 2009).

4.2.2 Blood-cerebrospinal fluid barrier

The blood-CSF barrier is established by the single epithelial cell layer in the choroid plexus and is present in all four brain ventricles (Chodobski et al., 2001). The blood-CSF barrier is located between the CSF and the fenestrated capillaries of the choroid plexus (De Bock et al., 2014; Mortazovi et al., 2014). The cubical choroid plexus epithelial cells (CPEs) possess multiple microvilli and cilia at their apical side and a well perfused extracellular matrix, consisting of connective tissue in combination with collagen, fibroblasts, macrophages and

dendritic cells, at their basal side (**Figure 10**) (Dohrmann et al., 1970; Mortazavi et al., 2014). They are the main producers of CSF (Cushing, 1914) and the remaining 10-30% of the CSF volume originates from the interstitial fluid in the brain (Cserr et al., 1986). The CSF is responsible for the transportation of necessary nutrients and other components in the brain by migrating from the lateral ventricles through the 3th and 4th ventricle to the subarachnoid space of the brain (Chodobski et al., 2001). Additionally, the choroid plexus participates in the elimination of harmful substances and degradation products in the CSF (Coisne et al., 2011) and plays an important role in the communication between the peripheral immune system and the CNS. Indeed, after leakage from the fenestrated capillaries, B cells, T cells and monocytes are able to migrate from the choroid plexus into the brain via specific transport mechanisms (Provencio et al., 2005; Meeker et al., 2012; Demeestere et al., 2015). Inflammation has already been linked with increases in the transport of these immune cells, resulting in a higher number of immune cells in the CSF (Vercellino et al., 2008). Furthermore, an upregulation of the expression of cytokines and inflammatory mediators has been reported in case of systemic inflammation (Kim et al., 2012; Vandenbroucke et al., 2012). Indeed, there is evidence that peripheral/systemic inflammation influences the pathophysiology of PD. In PD patients, there are gradients of objective measures of PD facets on serum cortisol and TNF- α , and of clinical motor scores on peripheral blood mononuclear cell production of cytokines and NF- κ B expression (Charlett et al. 1998; Dobbs et al. 1999; Reale et al, 2009). Serum cortisol is elevated (17%) in PD, as is IL-6. Moreover, in a prospective study, a higher concentration of IL-6, in blood collected 4 years previously, predicted incident PD (Chen et al. 2008). There are also gradients of objective measures of facets of PD on blood leucocyte subtypes counts (Dobbs et al, 2012). Both brady/hypokinesia and flexor rigidity are worse the higher the NK count.




Figure 10 : Illustration of the blood-cerebrospinal fluid barrier formed by the choroid plexus epithelial cells. These cells form a barrier between the CSF and the fenestrated capillaries of the stroma of the choroid plexus and they are localized in all four brain ventricles. CP: choroid plexus; ISF: interstitial fluid; CSF: cerebrospinal fluid; BBB: blood-brain barrier (adapted from De Bock et al., 2014).

Like the endothelial cells in the BBB, the CPEs are inter connected by a network of tight junctions. Together they allow the strict regulation of transport through the blood-CSF barrier (De Bock et al., 2014). Whereas claudin-3, -5 and -12 are important in the BBB, the main tight junction proteins expressed in the blood-CSF barrier are claudin-1, -2, -3 and -11. Claudin-2 is necessary for the formation of paracellular water channels (Rosenthal et al., 2010; Kratzer et al., 2012). As for the BBB, the transmembrane claudins are connected to the cytoskeleton of the cells via the ZO-1 and ZO-2 proteins (De Bock et al., 2014). Since the barrier formed by the choroid plexus cells is less tight compared to the BBB, the blood-CSF barrier is more susceptible to systemic alterations and inflammation and thereby this neural barrier is strongly subjected to immunologic influences (De Bock et al., 2014).

PD has also been associated with an increased permeability of the blood-CSF barrier (Vawter et al., 1996; Pisani et al., 2012). Significantly elevated levels of transforming growth factor beta (TGF- β) were demonstrated in the CSF of PD patients compared to controls. The same study suggested that some of the TGF- β in the CSF emanated from the plasma as a result of leakage (Vawter et al., 1996). Another study reported a significant increase in the albumin ratio (CSF albumin/serum albumin) in advanced-stage PD patients, compared to early-stage PD patients and healthy controls (Pisani et al., 2012).

4.3 Involvement of a *H. pylori* infection in PD: what is known so far?

As already mentioned above, the association between PD and gastrointestinal complaints has been suggested. Constipation, resulting from infrequent bowel movements and slow gut-transit (Abbott et al., 2001; Dobbs et al., 2016), has been described as one of the most common co-morbidities in PD patients (Parkinson, 1817; Dobbs et al., 2016). Schwab and colleagues were the first to propose the association between PD and the presence of gastric and duodenal ulcers (Schwab et al., 1961). In 1965, this was confirmed by Strang who reported an excess of previously diagnosed peptic ulcers in PD (Strang, 1965). Subsequently, the speculation that an infectious agent might be involved in both PD and peptic ulcers was suggested (Szabo, 1979). After the discovery of *H. pylori* in the stomach of human patients suffering from gastritis and peptic ulcers in 1984 (Warren and Marshall, 1984), Altschuler and colleagues were the first to explicitly suggest the causal link between PD and *H. pylori* (Altschuler et al., 1996). Ever since, the association between IP and an infection with *H. pylori* has been described by multiple other authors (Dukowicz et al., 2007; Nielsen et al., 2012; Dobbs et al., 2008; Dobbs et al., 2012; Tan et al., 2015; Csoti et al., 2016; Dobbs et al., 2016; Tan et al., 2015).

Up to date, a high prevalence of *H. pylori* infections has been reported in PD patients (Rees et al., 2011; Nielsen et al., 2012). Additionally, *H. pylori* seropositivity was 3 times more likely to occur among PD patients compared to control patients (Brady et al., 1998; Charlett et al., 1999). The same study showed that, compared to a control group, siblings of PD patients were more likely to be *H. pylori* seropositive and present Parkinson-like symptoms, such as brady/hypokinesia of gait, bradykinesia of hands, abnormal posture and rigidity (Brady et al., 1998; Charlett et al., 1999). Furthermore, compared to age-matched control patients, an increase of the level of *H. pylori* antibodies was reported in a group of patients (younger than 80 years) displaying Parkinson-

like symptoms (Dobbs et al., 2000). Another study described that the odds of parkinsonism increased fivefold (up to the age of 80) in case of CagA-positivity in combination with VacA negativity and urease B negativity (Weller et al., 2005). Moreover, increased numbers of natural killer cells have recently been associated with shorter stride, slower gait and elevated flexor rigidity in patients suffering from PD (Dobbs et al., 2012). Interestingly, the same study described higher numbers of natural killer cells in case of urea-breath test positivity as well as additional reduction in stride and speed in *H. pylori*-positive PD patients (Dobbs et al., 2012).

H. pylori eradication therapy, consisting of 20 mg omeprazole, 500 mg clarithromycin and 1 g amoxicillin, resulted in the improvement of the stride length in PD patients (Bjarnason et al., 2005; Dobbs et al., 2010). Despite a worsening of the upper limb flexor rigidity, an overall improvement of PD-related symptoms, irrespective of anti-parkinsonian medication, could be observed after biopsy-proven successful eradication of *H. pylori* in PD patients (Dobbs et al., 2010). Clarithromycin and/or amoxicillin were replaced by administration of 500 mg tetracycline and/or 400 mg metronidazole in case of *in vitro* resistance of *H. pylori* isolates (Dobbs et al., 2010; Dobbs et al., 2013).

Furthermore, *H. pylori* eradication therapy in patients suffering from PD resulted in an improved response to the administration of L-3,4-dihydroxyphenylalanine (levodopa) in combination with an amelioration of motor symptoms and life quality. Levodopa, which is mainly absorbed in the duodenum, is used as treatment of PD, as it is a precursor of dopamine. Since *H. pylori* infections may induce a disruption of the gastrointestinal mucosa (Hashim et al., 2014; Caron et al., 2015), an infection with this bacterium may thus interfere with the absorption of levodopa, resulting in decreased plasma levels of levodopa and thereby in a faster progression of PD-related symptoms (Pierantozzi et al., 2001; Fasano et al., 2013; Hashim et al., 2014). However, Dobbs et al. (2005) did not find any major differences in hypokinesia improvement following biopsy-proven *H. pylori* eradication between PD patients receiving levodopa and PD patients who were never treated with levodopa, suggesting that the effect on hypokinesia was not due to improved absorption of levodopa as a result of *H. pylori* eradication.

Several hypotheses have been proposed to declare the association between PD and *H. pylori*: it has been suggested that a chronic infection with *H. pylori* results in acquired immune suppression by the continuous antigenic stimulation from *H. pylori* infection in combination with increased circulating IL-6, thereby leading to autoimmunity (Dobbs et al., 2000). Another hypothesis is that there might be molecular mimicry between antigens from humans and *H. pylori*. Possible candidates are the heat shock proteins (hsp), which appear to be highly conserved from bacteria to mammals. Indeed, hsp60 antibodies have been detected in both *H. pylori*-positive patients and PD patients (Dobbs et al., 2000).

In conclusion, these data suggest that gastric infections with *H. pylori* might be associated with PD. In contrast, no information is currently available on whether there might be a possible link between PD and *H. suis* infections in human patients. Further research is required to obtain more information on the role of gastric *Helicobacter* infections in PD.

References

- Abenavoli L, Milic N, Masarone M, Persico M. 2013. Association between non-alcoholic fatty liver disease, insulin resistance and *Helicobacter pylori*. *Med Hypotheses* 81: 913-915.
- Abbott RD, Petrovitch H, White LR, Masaki KH, Tanner CM, Curb JD, Grandinetti A, Blanchette PL, Popper JS, Ross GW. 2001. Frequency of bowel movements and the future risk of Parkinson's disease. *Neurology* 5: 456-462.
- Adamopoulos AB, Stergiou GS, Sakizlis GN, Tiniakos DG, Nasothimiou EG, Sioutis DK, Achimastos AD. 2009. Diagnostic value of rapid urease test and urea breath test for *Helicobacter pylori* detection in patients with Billroth II gastrectomy: a prospective controlled trial. *Dig. Liver Dis.* 41: 4-8.
- Aggarwal N, Snyder P, Owens SR. 2011. Unusual *Helicobacter pylori* in gastric resection specimens: an old friend with a new look. *Int J Surg Pathol* 19: 297-302.
- Akada JK, Shirai M, Takeuchi H, Tsuda M, Nakazawa T. 2000. Identification of the urease operon in *Helicobacter pylori* and its control by mRNA decay in response to pH. *Mol. Microbio* 136: 1071-1084.
- Akashi R, Ishiguro N, Shimizu S, Kawashima M. 2011. Clinical study of the relationship between *Helicobacter pylori* and chronic urticarial and prurigo chronica multiformis: effectiveness of eradication therapy for *Helicobacter pylori*. *J Dermatol.* 38: 761-766.
- Akhiani AA, Schon K, Franzen LE, Poppo J, Lycke N. 2004. *Helicobacter pylori*-specific antibodies impair the development of gastritis, facilitate bacterial colonization and counteract resistance against infection. *J Immunol* 172: 5024-5033.
- Akopyants NS, Clifton SW, Kersulyte D, Crabtree JE, Youree BE, Reece CA, Bukanov NO, Drazek ES, Roe BA, Berg DE. 1998. Analysis of the cag pathogenicity island of *Helicobacter pylori*. *Mol Microbiol* 28: 37-53.
- Ali M, Whitehead M. 2008. Clearance of chronic psoriasis after eradication therapy for *Helicobacter pylori* infection. *J Eur Acad Dermatol Venereol* 22: 753-754.
- Allen LA, Schlesinger LS, Kang B. 2000. Virulent strains of *Helicobacter pylori* demonstrate delayed phagocytosis and stimulate homotypic phagosome fusion in macrophages. *J Exp Med* 191: 115-128.
- Allen LA. 2001. The role of neutrophils and phagocytosis in infection caused by *Helicobacter pylori*. *Curr Opin Infect Dis* 14: 273-277.
- Alm RA, Bina J, Andrews BM, Doig P, Hancock RE, Trust TJ. 2000. Comparative genomics of *Helicobacter pylori*: analysis of the outer membrane protein families. *Infect Immun* 68: 4155-4168.
- Altschuler E. 1996. Gastric *Helicobacter pylori* infection as a cause of idiopathic Parkinson disease and non-arteric anterior optic ischemic neuropathy. *Med Hypotheses* 47: 413-414.
- Alvarez-Arellano L, Maldonado-Bernal C. 2014. *Helicobacter pylori* and neurological diseases: married by the laws of inflammation. *World J Gastroenterol* 5(4): 400-404.
- Amedei A, Cappon A, Codolo G, Cabrelle A, Polenghi A, Benagiano M, Tasca E, Azzurri A, D'Elios MM, Del Prete G, de Bernard M. 2006. The neutrophil-activating protein of *Helicobacter pylori* promotes Th1 immune responses. *J Clin Invest.* 116: 1092-1101.
- Amedei A, Codolo G, Del Prete G, de Bernard M, D'Elios MM. 2010. The effect of *Helicobacter pylori* on asthma and allergy. *Journal of Asthma and Allergy* 3: 139-147.
- Amodio P, Montagnese S, Gatta A, Morgan MY. 2004. Characteristics of minimal hepatic encephalopathy. *Metab Brain Dis* 19: 253-267.
- Amory JR, Mackenzie AM, Pearce GP. 2006. Factors in the housing environment of finisher pigs associated with the development of gastric ulcers. *Vet Rec* 158(8): 260-264.
- Andersen LP, Boye K, Blom J, Holck S, Nørgaard A, Elsborg L. 1999. Characterization of a culturable "*Gastrosprillum hominis*" (*Helicobacter heilmannii*) strain isolated from human gastric mucosa. *J Clin Microbiol.* 37(4): 1069-1076.
- Andersen-Nissen E, Smith KD, Strobe KL, Barrett SL, Cookson BT, Logan SM, Aderem A., 2005. Evasion of Toll-like receptor 5 by flagellated bacteria. *Proc Natl Acad Sci USA.* 102: 9247-9252.
- Andrutis KA, Fox JG, Schauer DB, Marini RP, Li X, Yan L, Josenhans C, Suerbaum S. 1997. Infection of the ferret stomach by isogenic flagellar mutant strains of *Helicobacter mustellae*. *Infect Immun* 65: 1962-1966.

- Aragona P, Magazzù G, Macchia G, Bartolone S, Di Pasquale G, Vitali C, Ferreri G. 1999. Presence of antibodies against *Helicobacter pylori* and its heat shock protein 60 in the serum of patients with Sjögren's disease. *J Rheumatol* 26: 1306-1311.
- Argenzio RA and Eisemann J. 1996. Mechanisms of acid injury in porcine gastroesophageal mucosa. *Am J Vet Res* 57(4): 564-573.
- Argenziano G, Donnarumma G, Iovene MR, Arnese P, Baldassarre MA, Baroni A. 2003. Incidence of anti-*Helicobacter pylori* and anti-CagA antibodies in rosacea patients. *Int J Dermat* 42(8): 601-604.
- Arnold IC, Zigova Z, Holden M, Lawley TD, Rad R, Dougan G, Falkow S, Bentley DS, Müller A. 2011. Comparative whole genome sequence analysis of the carcinogenic bacterial model pathogen *Helicobacter felis*. *Genome Biol Evol* 3: 302-308.
- Asahi M, Azuma T, Ito S, Ito Y, Suto H, Nagai Y, Tsubokawa M, Tohyama Y, Maeda S, Omata M, Suzuki T, Sasakawa C. 2000. *Helicobacter pylori* CagA protein can be tyrosine phosphorylated in gastric epithelial cells. *J Exp Med* 191(4): 593-602.
- Aspholm M, Kalia A, Ruhl S, Schedin S, Arnqvist A, Linden S, Sjöström R, Gerhard M, Semino-Mora C, Dubois A, Unemo M, Danielsson D, Teneberg S, Lee WK, Berg DE, Boren T. 2006. *Helicobacter pylori* adhesion to carbohydrates. *Method Enzymol* 417: 293-339.
- Avci O, Ellidokuz E, Simşek I, Büyükgöbüz B, Güneş AT. 1999. *Helicobacter pylori* and Behçet's disease. *Dermatology* 199(2): 140-143.
- Avenaud P, Marais A, Monteiro L, Le Bail B, Bioulac Sage P, Balabaud C, Mégraud F. 2000. Detection of *Helicobacter* species in the liver of patients with and without primary liver carcinoma. *Cancer* 89(7): 1431-1439.
- Ayles HL, Friendship RM, Ball RO. 1996. Effects of dietary particle size on gastric ulcers, assessed by endoscopic examination, and relationship between ulcer severity and growth performance of individually fed pigs. *Swine Health Prod.* 4(5): 211-216.
- Azevedo NF, Almeida C, Fernandes I, Cerqueira L, Dias S, Keevil CW, Vieira MJ. 2008. Survival of gastric and enterohepatic *Helicobacter* species in water: implications for transmission. *Appl. Environ. Microbiol.* 74(6): 1805-1811.
- Backert S, Selbach M. 2008. Role of type IV secretion in *Helicobacter pylori* pathogenesis. *Cell Microbiol* 10(8): 1573-1581.
- Baele M, Van den Bulck K, Decostere A, Vandamme P, Hänninen ML, Ducatelle R, Haesebrouck F. 2004. Multiplex PCR assay for differentiation of *Helicobacter felis*, *H. bizzozeronii* and *H. salomonis*. *J Clin Microbiol.* 42(3): 1115-1122.
- Baele M, Decostere A, Vandamme P, Ceelen L, Hellemans A, Mast J, Chiers K, Ducatelle R, Haesebrouck F. 2008. Isolation and characterization of *Helicobacter suis* sp.nov. from pig stomach. *Int J Syst Evol Microbiol.* 58(6): 1350-1358.
- Bakar O, Demirçay Z, Yuksel M, Haklar G, Sanisoglu Y. 2007. The effect of azithromycin on reactive oxygen species in rosacea. *Clin Exp Dermatol* 32(2): 197-200.
- Bamford KB, Fan X, Cowe SE, Leary JF, Gourley WK, Luthra GK, Brooks EG, Graham DY, Reyes VE, Ernst PB. 1998. Lymphocytes in the human gastric mucosa during *Helicobacter pylori* have a T helper cell 1 phenotype. *Gastroenterol* 114(3): 482-492.
- Banchereau J, Briere F, Caux C, Davoust J, Lebecque S, Liu YJ, Pulendran B, Palucka K. 2000. Immunobiology of dendritic cells. *Annu Rev Immunol.* 18: 767-811.
- Barbosa AJ, Silva JC, Nogueira AM, Paulino Júnior E, Miranda CR. 1995. Higher incidence of *Gastrosprillum* sp. in swine with gastric ulcer of the pars oesophagea. *Vet Pathol* 32(2): 134-139.
- Barkun AN, Bardou M, Kuipers EJ, Sung J, Hunt RH, Martel M, Sinclair P, International Consensus Upper Gastrointestinal Bleeding Conference Group. 2010. International consensus upper gastrointestinal bleeding conference group: international consensus recommendations on the management of patients with nonvariceal upper gastrointestinal bleeding. *Ann. Intern. Med* 152(2): 101-113.
- Bassi V, Santinelli C, Iengo A, Romano C. 2010. Identification of a correlation between *Helicobacter pylori* infection and Grave's disease. *Helicobacter* 15(6): 558-562.
- Bassi V, Marino G, Iengo A, Fattoruso O, Santinelli C. 2012. Autoimmune thyroid diseases and *Helicobacter pylori*: the correlation is only present in Grave's disease. *World J Gastroenterol* 18(10): 1093-1097.
- Batts KP, Ketover S, Kakar S, Krasinskas AM, Mitchell KA, Wilcox R, Westerhoff M, Rank J, Gibson J, Mattia AR, Cummings OW, Davison JM, Naini BV, Dry SM, Yantiss RK, Rodger C Haggitt Gastrointestinal Pathology Society. 2013. Appropriate use of special stains for identifying *Helicobacter pylori*: Recommendations from the Rodger C. Haggitt Gastrointestinal Pathology Society. *Am J Surg Pathol.* 37(11): e12-22.
- Bener A, Micallef R, Afifi M, Derbala M, Al-Mulla HM, Usmani MA. 2007. Association between type 2 diabetes mellitus and *Helicobacter pylori* infection. *Turk J Gastroenterol* 18(4): 225-229.

- Bertalot G, Montresor G, Tampieri M, Spasiano A, Pedroni M, Milanese B, Favret M, Manca N, Negrini R. 2004. Decrease in thyroid autoantibodies after eradication of *Helicobacter pylori* infection. Clin Endocrinol (Oxf) 61(5): 650-652.
- Bhuiyan TR, Sahal A, Lundgren A, Qadri F, Svennerholm AM. 2010. Immune responses to *Helicobacter pylori* infections in Bangladeshi children during their first two years of life and the association between maternal antibodies and onset of infection. J Infect Dis 202(11): 1676-1684.
- Bjarnason IT, Charlett A, Dobbs RJ, Dobbs SM, Ibrahim MA, Kerwin RW, Mahler RF, Oxlade NL, Peterson DW, Plant JM, Price AB, Weller C. 2005. Role of chronic infection and inflammation in the gastrointestinal tract in the etiology and pathogenesis of idiopathic parkinsonism. Part 2: response of facets of clinical idiopathic parkinsonism to *Helicobacter pylori* eradication. A randomized, double-blind, placebo-controlled efficacy study. Helicobacter 10(4): 276-287.
- Blecker U, Renders F, Lanciers S, Vandenplas Y. 1991. Syncopes leading to the diagnosis of a *Helicobacter pylori* positive chronic active haemorrhagic gastritis. Eur J Pediatr 150(8): 560-561.
- Bode G, Marchildon P, Peacock J, Brenner H, and Rothenbacher D. 2002. Diagnosis of *Helicobacter pylori* infection in children: comparison of a salivary immunoglobulin G antibody test with the [(13)C] urea breath test. Clin Diagn Lab Immunol. 9(2): 493-495.
- Boren T, Falk P, Roth KA, Larson G, Normark S. 1993. Attachment of *Helicobacter pylori* to human gastric epithelium mediated by blood group antigens. Science 262(5141): 1892-1895.
- Bosschem I, Bayry J, De bruyne E, Van Deun K, Smet A, Vercauteren G, Ducatelle R, Haesebrouck F, Flahou B. 2015. Effect of different adjuvants on protection and side-effects induced by *Helicobacter suis* whole-cell lysate vaccination. PLoS ONE 10(6): e131364.
- Bosschem I, Flahou B, Bakker J, Heuvelman E, Langermans JAM, De Bruyne E, Joosten M, Smet A, Ducatelle R, Haesebrouck F. 2016. Comparative virulence of *in vitro* cultures primate- and pig-associated *Helicobacter suis* strains in a BALB/c mouse and Mongolian gerbil model. Helicobacter doi: 10.1111/hel.12349, in press.
- Bosschem I, Flahou B, Van Deun K, De Koker S, Volf J, Smet A, Ducatelle R, Devriendt B, Haesebrouck F. 2016. Species-specific immunity to *Helicobacter suis*. Submitted.
- Brady P, Charlett A, Dobbs RJ, Peterson DW, Wilks M, Weller C, Dobbs SM. 1998. Increased prevalence of *Helicobacter pylori* seropositivity in sufferers from idiopathic parkinsonism and their siblings. British Journal of Clinical Pharmacology 46, 302-303P.
- Brandt S, Kwok T, Hartig R, König W, Blackert S. 2005. NF-KappaB activation and potentiation of proinflammatory responses by the *Helicobacter pylori* CagA protein. Proc Natl Acad Sci Usa 102(26): 9300-9305.
- Bruel H, Dabadie A, Pouedras P, Gambert C, Le Gall E, Jezequel C. 1993. *Helicobacter pylori* infection manifested by acute anemia. Ann Pediatr (Paris) 40(6): 364-367.
- Buchner AM, Blonski W, Lichtenstein GR. 2011. Update on the management of Crohn's disease. Curr Gastroenterol Rep. 13(5): 465-474.
- Budzynski J, Klopocka M. 2014. Brain-gut axis in the pathogenesis of *Helicobacter pylori* infection. World Journal Gastroenterol 20(18): 5212-5225.
- Burne RA and Chen YY. 2000. Bacterial ureases in infectious diseases. Microbes Infect 2(5): 533-542.
- Burucoa C, Delchier JC, Courillon-Mallet A, de Korwin JD, Mégraud F, Zerbib F, Raymond J, Fauchère JL. 2013. Comparative evaluation of 29 commercial *Helicobacter pylori* serological kits. Helicobacter 18(3): 169-179.
- Busiello I, Acquaviva R, Di Popolo A, Blanchard TG, Ricci V, Romano M, Zarrilli R. 2004. *Helicobacter pylori* gamma-glutamyltranspeptidase upregulates COX-2 and EGF-related peptide expression in human gastric cells. Cell Microbiol 6(3): 255-267.
- Calvet X, Ramírez Lázaro MJ, Lehours P, Mégraud F. 2013. Diagnosis and epidemiology of *Helicobacter pylori* infection. Helicobacter 18(1): 5-11.
- Campuzona-Maya G. 2014. Hematologic manifestations of *Helicobacter pylori* infection. World Journal of Gastroenterology 20(36): 12818-12838.
- Cantet F, Magras C, Marais A, Federighi M, Mégraud F. 1999. *Helicobacter* species colonizing pig stomach: molecular characterization and determination of prevalence. Appl Environ Microbiol 65(10): 4672-4676.
- Cappon A, Babolin C, Segat D, Cancian L, Amadei A, Calzetti F, Cassatella MA, D'Elis MM, Bernard M. 2010. *Helicobacter pylori*-derived neutrophil activating protein increases the lifespan of monocytes and neutrophils. Cell Microbiol.12(6): 754-64.

- Cardenas VM, Dominguez DC, Puentes FA, Aragaki CC, Goodman KJ, Graham DY, Fukuda Y. 2008. Evaluation of a novel stool native catalase antigen test for *Helicobacter pylori* infection in asymptomatic North American children. *J. Pediatr. Gastroenterol. Nutr* 46(4): 399-402.
- Cario E, Brown D, McKee M, Lynch-Devaney K, Gerken G, Podolsky DK. 2002. Commensal-associated molecular patterns induce selective Toll-like receptor-trafficking from apical membrane to cytoplasmic compartments in polarized intestinal epithelium. *Am J Pathol.* 160(1): 165-173.
- Caron TJ, Scott KE, Fox JG, Hagen SJ. 2015. Tight junction disruption: *Helicobacter pylori* and dysregulation of the gastric mucosal barrier. *World J Gastroenterol.* 21(40): 11411-11427.
- Caso JR, Leza JC, Menchen L. 2008. The effects of physical and physiological stress on the gastrointestinal tract: lessons from animal models. *Curr Mol Med* 8: 299-312.
- Casagrande Proietti P, Bietta A, Brachelente C, Lepri E, Davidson I, Franciosini MP. 2010. Detection of *Helicobacter* spp. in gastric, fecal and saliva samples from swine affected by gastric ulceration. *J Vet Sci.* 11(3): 221-225.
- Casella AM, Berdel RF, Bressanim GL, Malaguido MR, Cardillo JA. 2012. *Helicobacter pylori* as a potential target for the treatment of central serous chorioretinopathy. *Clinics (Sao Paulo)* 67(9): 1047-1052.
- Cats A, Meuwissen SG, Forman D, Craanen ME, Kuipers EJ. 1998. *Helicobacter pylori*: a true carcinogen? *European Journal of Gastroenterology and Hepatology* 10(6): 447-450.
- Chang YP, Chiu GF, Kuo FC, Lai CL, Yang YH, Hu HM, Chang PY, Chen CY, Wu DC, Yu FJ. 2013. Eradication of *Helicobacter pylori* is associated with the progression of dementia: A population-based study. *Gastroenterol Res Pract.* 175729.
- Chang CS, Chen LT, Yang JC, Lin JT, Chang KC, Wang JT. 1999. Isolation of a *Helicobacter pylori* protein, FldA, associated with mucosa-associated lymphoid tissue lymphoma of the stomach. *Gastroenterology* 117(1): 82-88.
- Charlett A, Dobbs RJ, Weller C, Dobbs SM. 1997. Stasis in the gut: The source of xenobiotic in idiopathic parkinsonism. *Eur J Clin Pharmacol* 52 (suppl): 168.
- Charlett A, Dobbs RJ, Purkiss AG, Wright, DJ, Peterson DW, Weller C, Dobbs SM. 1998. Cortisol is higher in parkinsonism and associated with gait deficit. *Acta Neurol Scand* 97: 77-85.
- Charlett A, Dobbs RJ, Dobbs SM, Weller C, Brady P, Peterson DW. 1999. Parkinsonism: siblings share *Helicobacter pylori* seropositivity and facets of syndrome. *Acta Neurol Scand* 99(1): 26-35.
- Chehter EZ, Bacci MR, Fonseca FL, Gonçalves JA, Buchalla G, Shiraichi SA, Mariano RC. 2013. Diagnosis of the infection by the *Helicobacter pylori* through stool examination: method standardization in adults. *Clin. Biochem* 46(15): 1622-1624.
- Chen Y, Blaser MJ. 2008. *Helicobacter pylori* colonisation is inversely associated with childhood asthma. *J Inf Dis.* 198(4): 553-560.
- Chen H, O'Reilly EJ, Schwarzschild MA, Ascherio A. 2008. Peripheral Inflammatory Biomarkers and Risk of Parkinson's Disease. *Am J Epidemiol* 167: 90-95.
- Chiba S, Sugiyama T, Yonekura K, Tanaka S, Matsumoto H, Fujii N, Ebisu S, Sekiguchi K. 2002. An antibody to VacA of *Helicobacter pylori* in cerebrospinal fluid from patients with Guillain-Barre syndrome. *J Neurol Neurosurg Psychiatry.* 73(1): 76-78.
- Chimienti G, Russo F, Lamanuzzi B, Nardulli M, Messa C, Di Leo A, Correale M, Giannuzzi V, Pepe G. 2003. *Helicobacter pylori* is associated with modified lipid profile: impact on lipoprotein (a). *Clin biochem* 36(5): 359-365.
- Chiu YC, Tai WC, Chuah SK, Hsu PI, Wu DC, Wu KL, Huang CC, Ho JC, Ring J, Chen WC. 2013. The clinical correlations of *H. pylori* virulence factors and chronic spontaneous urticaria. *Gastroenterol Res Pract* 2013: 456727.
- Chodobski A, Szmydynger, Chodobska J. 2001. Choroid plexus: target for polypeptides and site of their synthesis. *Microsc Res Techniq* 52(1): 65-82.
- Choi KY, Han JH, Joo HS. 2001a. Identification of a novel *Helicobacter* species in the pig stomachs by PCR and partial sequencing. *J Clin Microbiol* 39(9): 3311-3315.
- Choi J, Kim CH, Kim D, Chung SJ, Song JH, Kang JM, Yang JI, Park MJ, Kim YS, Yim JY, Lim SH, Kim JS, Jung HC, Song IS. 2011. Prospective evaluation of the new stool antigen test for the detection of *Helicobacter pylori*, in comparison with histology, rapid urea breath test, (13)C-urea breath test and serology. *J. Gastroenterol. Hepatol.* 26(6): 1053-1059.

- Choi JJ. 2012. Are there great savings with rapid urease test by one-plus-one? Clin Endosc 45(4): 341-342
- Christie PJ and Vogel JP. 2000. Bacterial type IV secretion: conjugation systems adapted to deliver effector molecules to host cells. Trends Microbiol. 8(3): 354-360.
- Chung TH, Kim HD, Lee YS, Hwang CY. 2014. Determination of the prevalence of *Helicobacter heilmannii*-like organisms type 2 (HHLO-2) infection in humans and dogs using non-invasive genus/species-specific PCR in Korea. J Vet Med Sci.76(1): 73–9.
- Cindoruk M, Cirak MY, Unal S, Karakan T, Erkan G, Engin D, Dumlu S, Turet S. 2008. Identification of *Helicobacter* species by 16S rDNA PCR and sequence analysis in human liver samples from patients with various etiologies of benign liver diseases. Eur J Gastroenterol Hepatol 20(1): 33-36.
- Cirak MY, Akyön Y, Mégraud F. 2007. Diagnosis of *Helicobacter pylori*. Helicobacter 12(1): 4-9.
- Coisne C, Engelhardt B. 2011. Tight junctions in brain barriers during central nervous system inflammation. Antioxidants and Redox Signaling 15(5): 1285-1303.
- Collett JA, Burt MJ, Frampton CM, Yeo KH, Chapman TM, Buttimore RC, Cook HB, Chapman BA. 1999. Seroprevalence of *Helicobacter pylori* in the adult population of Christchurch: risk factors and relationship to dyspeptic symptoms and iron studies. N Z Med J 112(1093): 292-295.
- Cornetta AM, Simpson KW, Strauss-Ayali D, McDonough PL, Gleed RD. 1998. Use of the [13C]urea breath test for the detection of gastric infection with *Helicobacter* spp. in dogs. Am J Vet Res 59(11): 1364-1369.
- Corada M, Mariotti M, Thurston G, Smith K, Kunkel R, Brockhaus M, Lampugnani MG, Martin-Padura I, Stoppacciaro A, Ruco L, McDonald DM, Ward PA, Dejana E. 1999. Vascular endothelial-cadherin is an important determinant of microvascular integrity *in vivo*. P Natl Acad Sci USA 96(17): 9815-9820.
- Corthésy-Theulaz IE, Corthésy B, Bachmann D, Velin D, Kraehenbuhl JP. 2003. Passive immunity in *Helicobacter* challenged neonatal mice conferred by immunized dams lasts until weaning. Inf Immun 7(4): 2226-2229.
- Cotticelli L, Borrelli M, D'Alessio AC, Menzione M, Villani A, Piccolo G, Montella F, Iovene MR, Romano M. 2006. Central serous chorioretinopathy and *Helicobacter pylori*. Eur J ophthalmol 16(2): 274-278.
- Cover TL and Blaser MJ. 1992: Purification and characterization of the vacuolating toxin from *Helicobacter pylori*. J Biol Chem 267(15): 10570-10575.
- Cryan JF, Dinan TG. 2012. Mind altering microorganisms: the impact of the gut microbiota on the brain and behavior. Nat Rev Neurosci. 13(10): 701-12.
- Cserr FH, Depasquale M, Palak CS, Pullen RG. 1986. Convection of cerebral interstitial fluid and its role in brain volume regulation. Annals of the New York Academy of Sciences 481: 123-134.
- Csoti I, Jost WH, Reichmann H. 2016. Parkinson's disease between internal medicine and neurology. J Neural Transm 123(1): 3-17.
- Cushing H. 1914. Studies on the cerebrospinal fluid. I. Introduction. J Med Res 26: 1-19.
- Cussac V, Ferrero RL, Labigne A. 1992. Expression of *Helicobacter pylori* urease genes in *Escherichia coli* grown under nitrogen-limiting conditions. J Bacteriol 174(8): 2466-2473.
- D'Elios MM, Manghetti M, De Carli M, Costa F, Baldari CT, Burrioni D, Telford JL, Romagnani S, Del Prete G. 1997. T helper 1 effector cells specific for *Helicobacter pylori* in the gastric antrum of patients with peptic ulcer disease. J Immunol 158(2): 962-967.
- Dang Y, Mu Y, Zhao M, Li L, Guo Y, Zhu Y. 2013. The effect of eradicating *Helicobacter pylori* on idiopathic central serous chorioretinopathy patients. Ther Clin Risk Manag 9: 355-360.
- Davies GR, Simmonds NJ, Stevens TR, Sheaff MT, Banatvala N, Laurenson IF, Blake DR, Rampton DS. 1994. *Helicobacter pylori* stimulates antral mucosal reactive oxygen metabolite production *in vivo*. Gut 35(2): 179-185.
- De Bastiani R, Gabrielli M, Ubaldi E, Benedetto E, Sanna G, Cottone C, Candelli M, Zocco MA, Saulnier N, Santoliquido A, Papaleo P, Gasbarrini G, Gasbarrini A. 2008. High prevalence of CagA positive *H. pylori* strains in ischemic stroke: a primary care multicenter study. Helicobacter 13(4): 274-277.

- De Bernard M, D'Elios MM. 2010. The immune modulating activity of the *Helicobacter pylori* HP-NAP: Friend or foe? *Toxicon* 56(7): 1186-1192.
- De Block M, D'Herde K, Duchateau L, Decostere A, Haesebrouck F, Ducatelle R. 2006. The effect of *Helicobacter felis* and *Helicobacter bizzozeronii* on the gastric mucosa in Mongolian gerbils: a sequential pathological study. *J Comp Pathol* 135(4): 226-236.
- De Bock M, Van den Bulck K, Hellemans A, Daminet S, Coche JC, Debongnie JC, Decostere A, Haesebrouck F, Ducatelle R. 2007. Peptic ulcer disease associated with *Helicobacter felis* in a dog owner. *Eur. J. Gastroenterol. Hepatol.* 19(1): 79-82.
- De Bock M, Vandenbroucke RE, Decrock E, Culot M, Cecchelli R, Leybaert L. 2014. A new angle on blood-CNS interfaces: a role for connexins? *FEBS Lett* 588(8): 1259-1270.
- Debongnie JC, Donnay M, Mairesse J. 1995. *Gastrospirillum hominis* ("*Helicobacter heilmannii*"): a cause of gastritis, sometimes transient, better diagnosed by touch of cytology? *Am J Gastroenterol.* 90(3): 411-416.
- De Bruyne E, Flahou B, Chiers K, Meyns T, Kumar S, Vermoote M, Pasmans F, Millet S, Dewulf J, Haesebrouck F, Ducatelle R. 2012. An experimental *Helicobacter suis* infection causes gastritis and reduced daily weight gain in pigs. *Veterinary Microbiology* 160(3-4): 449-454.
- De Cooman L, Flahou B, Houf K, Smet A, Ducatelle R, Pasmans F, Haesebrouck F. 2013. Survival of *Helicobacter suis* bacteria in retail pig meat. *Int. J. Food Microbiol.* 166(1): 164-167.
- Deguchi R, Matsushima M, Suzuki T, Mine T, Fukuda R, Nishina M, Ozawa H, Takagi A. 2009. Comparison of a monoclonal with a polyclonal antibody-based enzyme immunoassay stool test in diagnosing *Helicobacter pylori* infection after eradication therapy. *J. Gastroenterol* 44(7): 713-716.
- De Groote D, Haesebrouck F, van Doorn LJ, Vandamme P, Ducatelle R. 2001. Evaluation of a group-specific 16S ribosomal DNA-based PCR for detection of *Helicobacter bizzozeronii*, *Helicobacter felis* and *Helicobacter salomonis* in fresh and paraffin-embedded gastric biopsy specimens. *J Clin Microbiol* 39(3): 1197-1199.
- De Groote D, Van Doorn LJ, Van den Bulck K, Vandamme P, Vieth M, Stolte M, Debongnie JC, Burette A, Haesebrouck F, Ducatelle R. 2005. Detection of non-pylori *Helicobacter* species in "*Helicobacter heilmannii*"-infected humans. *Helicobacter* 10(5): 398-406.
- Deleu D, Hanssens Y, Worthing EA. 1998. Systematic and prophylactic treatment of migraine: a critical reappraisal. *Clin Neuropharmacol* 21(5): 267-279.
- De Martel C, Plummer M, van Doorn LJ, Vivas J, Lopez G, Carillo E, Peraza S, Muñoz N, Franceschi S. 2010. Comparison of polymerase chain reaction and histopathology for the detection of *Helicobacter pylori* in gastric biopsies. *Int J Cancer* 126(8): 1992-1996.
- Demeestere D, Libert C, Vandenbroucke RE. 2015. Clinical implications of leukocyte infiltration at the choroid plexus in (neuro)inflammatory disorders. *Drug Discovery Today* 20(8): 928-941.
- De Pascalis R, Del Pezzo M, Nardone G, Budillon G, A. Lavitola A. 1999. Performance characteristics of an enzyme-linked immunosorbent assay for determining salivary immunoglobulin G response to *Helicobacter pylori*. *J Clin Microbiol* 37(2): 430-432.
- Devrajani BR, Shah SZ, Soomro AA, Devrajani T. 2010. Type 2 diabetes mellitus: A risk factor for *Helicobacter pylori* infection: A hospital based case-control study. *Int J Diabetes Dev Ctries* 30(1): 22-26.
- Dewhirst FE, Shen Z, Scimeca MS, Stokes LN, Boumenna T, Chen T, Paster BJ, Fox JG. 2005. Discordant 16S and 23S rRNA gene phylogenies for the genus *Helicobacter*: implications for phylogenetic inference and systematics. *J Bacteriol* 187(17): 6106-6118.
- Dieterich C, Wiesel P, Neiger R, Blum A, Corthésy-Theulaz I. 1998. Presence of multiple "*Helicobacter heilmannii*" strains in an individual suffering from ulcers and his two cats. *J. Clin. Microbiol.* 36: 1366-1370.
- Dobbs RJ, Charlett A, Purkiss AG, Dobbs SM, Weller C, Peterson DW. 1999. Association of circulating TNF- α and IL-6 with ageing and parkinsonism. *Acta Neurol Scand* 100: 34-41.
- Dobbs RJ, Charlett A, Dobbs SM, Weller C, Peterson DW. 2000. Parkinsonism: differential age-trend in *Helicobacter pylori* antibody. *Alimentary Pharmacology and Therapeutics* 14 (9): 1199-1205.
- Dobbs SM, Dobbs RJ, Weller C, Charlett A. 2000. Link between *Helicobacter pylori* and idiopathic parkinsonism. *Med Hypoth* 55 (2): 93-98.
- Dobbs RJ, Dobbs SM, Weller C, Bjarnason IT, Oxlade NL, Charlett A, Al-Janabi MA, Kerwin RW, Mahler RF, Price AB. 2005. Role of Chronic Infection and Inflammation in the Gastrointestinal Tract in the Aetiology and Pathogenesis of Idiopathic Parkinsonism. Part 1: eradication of *Helicobacter* in the Cachexia of Idiopathic Parkinsonism. *Helicobacter* 10: 267-275.

- Dobbs RJ, Dobbs SM, Weller C, Charlett A, Bjarnason IT, Curry A, Ellis DS, Ibrahim MA, McCrossan MV, O'Donohue J, Owen RJ, Oxlade NL, Price AB, Sanderson JD, Sudhanva M, Williams J. 2008 *Helicobacter* hypothesis for IP: before and beyond. *Helicobacter* 13(5): 309-322.
- Dobbs SM, Dobbs RJ, Weller C, Charlett A, Bjarnason IT, Lawson AJ, Letley D, Harbin L, Price AB, Ibrahim MA, Oxlade NL, Bowthorpe J, Leckstroem D, Smee C, Plant JM, Peterson DW. 2010. Differential effect of *Helicobacter pylori* eradication on time trends in brady/hypokinesia and rigidity in Idiopathic Parkinsonism. *Helicobacter* 15(4): 279-294.
- Dobbs RJ, Charlett A, Dobbs SM, Weller C, A Ibrahim MA, Iguodala O, Smee C, Plant JM, Lawson AJ, Taylor D, Bjarnason I. 2012. Leukocyte-subset counts in idiopathic parkinsonism provide clues to a pathogenic pathway involving small interstitial bacterial overgrowth. A surveillance study. *Gut Pathog* 4(1): 12.
- Dobbs SM, Charlett A, Dobbs RJ, Weller C, Iguodala O, Smee C, Lawson AJ, Taylor D, Bjarnason I. 2013. Antimicrobial surveillance in idiopathic parkinsonism: indication-specific improvement in hypokinesia following *Helicobacter pylori* eradication and non-specific effect of antimicrobials for other indications in worsening rigidity. *Helicobacter* 18(3): 187-196.
- Dobbs SM, Dobbs RJ, Weller C, Charlett A, Augustin A, Taylor D, Ibrahim MA, Bjarnason I. 2016. Peripheral aetiopathogenic drivers and mediators of Parkinson's disease and co-morbidities: role of gastrointestinal microbiota. *J. Neuroviol* 22(1): 22-32.
- Doğan Z, Filik L, Ergül B, Sarikaya M, Akbal E. 2013. Association between *Helicobacter pylori* and liver-to-spleen ratio: a randomized-controlled single-blind study. *Eur J Gastroenterol Hepatol* 25(1): 107-110.
- Dohrmann GJ. 1970. The choroid plexus: a historical review. *Brain research* 18(2): 197-218.
- Domínguez J, Forné M, Blanco S, Prat C, Galí N, Latorre I, Viver JM, Ausina V. 2006. Comparison of a monoclonal with a polyclonal antibody-based enzyme immunoassay stool test in diagnosing *Helicobacter pylori* infection before and after eradication therapy. *Alim. Pharmacol. Ther.* 23(12): 1735-1740.
- Donahue JP, Peek RM, Van Doorn LJ, Thompson SA, Xu Q, Blaser MJ, Miller GG. 2000. Analysis of iceA1 transcription in *Helicobacter pylori*. *Helicobacter* 5(1): 1-12.
- Dore MP, Realdi G, Mura D, Graham DY, Sepulveda AR. 2002. *Helicobacter* infection in patients with HCV-related chronic hepatitis, cirrhosis, and hepatocellular carcinoma. *Dig Dis Sci* 47(7): 1638-1643.
- Dorsey ER, Constantinescu R, Thompson JP, Biglan KM, Holloway RG, Kieburtz K, Marshall FJ, Ravina BM, Schifitto G, Siderowf A, Tanner CM. 2007. Projected number of people with Parkinson disease in the most populous nations, 2005 through 2030. *Neurology* 68(5): 384-386.
- Dubois A and Borén T, 2007: *Helicobacter pylori* is invasive and it may be a facultative intracellular organisms. *Cell. Microbiol* 9(5): 1108-1116.
- Dukowicz AC, Lacy BE, Levine GM. 2007. Small intestinal bacterial overgrowth: a comprehensive review. *Gastroenterol Hepatol* 3(2): 112-122.
- Dufour C, Brisigotti M, Fabretti G, Luxardo P, Mori PG, Barabino A. 1993. *Helicobacter pylori* gastric infection and sideropenic refractory anemia. *J Pediatr Gastroenterol Nutr* 17(2): 225-227.
- Duquenois A and Le Luyer B. 2009. Gastritis caused by *Helicobacter heilmannii* probably transmitted from dog to child. *Arch Pediatr* 16(5): 426-429.
- Eaton KA, Brooks CL, Morgan DR, Krakowka S. 1991. Essential role of urease in pathogenesis of gastritis induced by *Helicobacter pylori* in gnotobiotic piglets. *Infect Immun* 59(7): 2470-2475.
- Eaton KA, Dewhirst FE, Radin MJ, Fox JG, Paster BJ, Krakowka S, Morgan DR. 1993. *Helicobacter acinonyx* sp. nov., isolated from cheetahs with gastritis. *Int J Syst Bacteriol* 43(1): 99-106.
- Eaton KA, Dewhirst FE, Paster BJ, Tzellas N, Coleman BE, Paola J, Sherding R. 1996. Prevalence and varieties of *Helicobacter* species in dogs from random sources and pet dogs: animal and public health implications. *J Clin Microbiol* 34(12): 3165-3170.
- Eaton KA, Mefford M, Thevenot T. 2001. The role of T cell subsets and cytokines in the pathogenesis of *Helicobacter pylori* gastritis in mice. *J Immunol* 166(12): 7456-7461.
- Ek CJ, Dziegielewska KM, Stolp H, Saunders NR. 2006. Functional effectiveness of the blood-brain barrier to small water soluble molecules in developing and adult opossum (*Monodelphis domestica*). *The Journal of Comparative Neurology* 496(1): 13-26.

- Ekman E, Fredriksson M, Trowald-Wigh G 2013. *Helicobacter* spp. in the saliva, stomach, duodenum and faeces of colony dogs. *Vet J* 195(1): 127-129.
- El-Zimaity H, Serra S, Szentgyorgyi E, Vajpeyi R, Samani A. 2013. Gastric biopsies: a gap between evidence-based medicine and daily practice in the management of gastric *Helicobacter pylori* infection. *Can J Gastroenterol* 27(10): 25-30.
- Engstrand L, Lindberg M. 2013. *Helicobacter pylori* and the gastric microbiota. *Best Pract Res Clin Gastroenterol*. 27: 39–45.
- Ernst FD, Homuth G, Stoof J, Mäder U, Waidner B, Kuipers EJ, Kist M, Kusters JG, Bereswill S, van Vliet AH. 2005. Iron-responsive regulation of the *Helicobacter pylori* iron-cofactored superoxide dismutase SodB is mediated by Fur. *J Bacteriol* 187(11): 3687-3692.
- Esmat G, El-Bendary M, Zakarya S, Ela MA, Zalata K. 2012. Role of *Helicobacter pylori* in patients with HCV-related chronic hepatitis and cirrhosis with or without hepatocellular carcinoma: possible association with disease progression. *J Viral Hepat* 19(7): 473-479.
- Evans DJ, Evans DG, Takemura T, Nakano H, Lampert HC, Graham DY, Granger ND, Kvietys PR. 1995. Characterization of a *Helicobacter pylori* neutrophil-activating protein. *Infect Immun*. 63(6): 2213–2220.
- Fahn S. 2003. Description of Parkinson's disease as a clinical syndrome. *Ann N Y Acad Sci* 991: 1-14.
- Fasano A, Bove F, Gabrielli M, Petracca M, Zocco MA, Ragazzoni E, Barbaro F, Piano C, Fortuna S, Tortora A, Di Giacopo R, Campanale M, Gigante G, Lauritano EC, Navarra P, Marconi S, Gasbarrini A, Bentivoglio AR. 2013. The role of small intestinal bacterial overgrowth in Parkinson's disease. *Mov Disord* 28(9): 1241-1249.
- Federman DG, Kirsner RS, Moriarty JP, Concato J. 2003. The effect of antibiotic therapy for patients infected with *Helicobacter pylori* who have chronic urticaria. *J Am Acad Dermatol*. 49(5): 861-864.
- Feingold KR, Grunfeld C. 1992. Role of cytokines in inducing hyperlipidemia. *Diabetes* 41(2): 97-101.
- Feng H, Zhou X, Zhang G. 2014. Association between cirrhosis and *Helicobacter pylori* infection: a meta-analysis. *Eur J Gastroenterol Hepatol* 26(12): 1309-1319.
- Fischer W, Püls J, Buhrdorf R, Gebert B, Odenbreit S, Haas R. 2001. Systematic mutagenesis of the *Helicobacter pylori* *cag* pathogenicity island: essential genes for CagA translocation in host cells and induction of interleukin-8. *Mol Microbiol*. 42(5): 1337-1348.
- Flahou B, Haesebrouck F, Pasmans F, D'Herde K, Driessen A, Van Deun K, Smet A, Duchateau L, Chiers K, Ducatelle R. 2010. *Helicobacter suis* causes severe gastric pathology in mouse and Mongolian gerbil models of human gastric disease. *Plos one*. 5 (11): e14083.
- Flahou B, Haesebrouck F, Chiers K, Van Deun K, De Smet L, Devreese B, Vandenberghe I, Favoreel H, Smet A, Pasmans F, D'Herde K, Ducatelle R. 2011. Gastric epithelial cell death caused by *Helicobacter suis* and *Helicobacter pylori* γ -glutamyl transpeptidase is mainly glutathione degradation-dependent. *Cell Microbiol* 13(12): 1933-1955.
- Flahou B, Van Deun K, Pasmans F, Smet A, Volf J, Rychlik I, Ducatelle R, Haesebrouck F. 2012. The local immune response of mice after *Helicobacter suis* infection: strain differences and distinction with *H. pylori*. *Vet Research* 43(1): 75.
- Flahou B, Haesebrouck F, Smet A. 2016. Non-*Helicobacter pylori* *Helicobacter* infections in humans and animals. Book: *Helicobacter pylori* research: from bench to bedside, Springer Japan, in press.
- Foyes S, Dorrell N, Ward SJ, Stabler RA, McColm AA, Rycroft AN, Wren BW. 2000. *Helicobacter pylori* possesses two CheY response regulators and a histidine kinase sensor, Che A, which are essential for chemotaxis and colonization of the gastric mucosa. *Infect Immun* 68(4): 2016-2023.
- Fox JG, Beck P, Dangler CA, Whary MT, Wang CT, Shi HN, Nagler-Anderson C. 2000. Concurrent enteric helminth infection modulates inflammation and gastric immune responses and reduces *Helicobacter*-induced gastric atrophy. *Nat Met* 6(5): 536-542.
- Franceschi F, Sepulveda AR, Gasbarrini A, Pola P, Silveri NG, Gasbarrini G, Graham DY, Genta RM. 2002. Cross-reactivity of anti CagA antibodies with vascular wall antigens: possible pathogenic link between *Helicobacter pylori* infection and atherosclerosis. *Circulation* 106(4): 430-434.
- Franceschi F, Christodoulides N, Kroll MH, Genta RM. 2004. *Helicobacter pylori* and idiopathic thrombocytopenic purpura. *Ann Intern Med*. 140: 766-767.
- Fukuda S, Shimoyama T, Umegaki N, Mikami T, Nakano H, Munakata A. 2004. Effect of *Helicobacter pylori* eradication in the treatment of Japanese patients with chronic idiopathic urticarial. *J Gastroenterol*. 39(9): 827-830.

- Furuse M, Fujita K, Hiragi T, Fujimoto K, Tsukita S. 1998a. Claudin -1 and -2: Novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J Cell Biol* 141(7): 1539-1550.
- Furese M, Sasaki H, Fujimoto K, Tsukita S. 1998b. A single gene product, claudin-1 or -2, reconstitutes tight junction strands and recruits occludin in fibroblasts. *J Cell Biol*. 143(2): 391-401.
- Garhart CA, Nedrud JG, Heinzel FP, Sigmund NE, Czinn SJ. 2003. "Vaccine-induced protection against *Helicobacter pylori* in mice lacking both antibodies and interleukin-4." *Infection and Immunity* 71(6): 3628-3633.
- Gasbarrini A, De Luca A, Fiore G, Franceschi F, Ojetto VV, Torre ES, Di Campli C, Candelli M, Pola R, Serricchio M, Tondi P, Gasbarrini G, Pola P, Giacobovazzo M. 1998. Primary headache and *Helicobacter pylori*. *Int J Angiol* 7(4): 310-312.
- Gasbarrini A, Franceschi F, Tartaglione R, Landolfi R, Pola P, Gasbarrini G. 1998. Regression of autoimmune thrombocytopenia after eradication of *Helicobacter pylori*. *Lancet* 352(9131): 878.
- Ge RG, Wang DX, Hao MC, Sun XS. 2013. Nickel trafficking system responsible for urease maturation in *Helicobacter pylori*. *World J Gastroenterol*. 19(45): 8211-8218.
- Gerhard M, Hirno S, Wadström T, Miller-Podraza H, Teneberg S, Karlsson KA, Appelmelk B, Odenbreit S, Haas R, Arnqvist A, Boren T. 2001. *Helicobacter pylori*, an adherent pain in the stomach, p. 185-206. In Achtmann SSM (ed.), *Helicobacter pylori*, molecular and cellular biology, 1st Ed. Horizon Scientific Press, Wymondham, United Kingdom.
- Gerhard A, Pavese N, Hotton G, Turkheimer F, Es M, Hammers A, Eggert K, Oertel W, Banati RB, Brooks DJ. 2006. *In vivo* imaging of microglial activation with [11C](R)-PK11195 PET in idiopathic Parkinson's disease. *Neurobiol Dis* 21(2): 404-412.
- Gervil M, Ulrich V, Kaprio J, Olesen J, Russell MB. 1999. The relative role of genetic and environmental factors in migraine without aura. *Neurology* 53(5): 995-999.
- Ghil HM, Yoo JH, Jung WS, Chung TH, Youn HY, Hwang CY. 2009. Survey of *Helicobacter* infection in domestic and feral cats in Korea. *J Vet Sci*. 10(1): 67-72.
- Gibb WR, Lees AJ. 1988. The relevance of Lewy body to the pathogenesis of idiopathic Parkinson's disease. *J Neurol Neurosurg Psychiatry*. 51(6): 745-752.
- Gisbert JP and Pajares JM. 2004. Review article: 13C-urea breath test in the diagnosis of *Helicobacter pylori* infection-a critical review. *Aliment. Pharmacolog. Ther.* 20(10): 1001-1017.
- Gisbert JP, de la Morena F, Abaira V. 2006. Accuracy of monoclonal stool antigen test for the diagnosis of *H. pylori* infection: a systematic review and meta-analysis. *Am. J. Gastroenterol*. 101(8): 1921-30.
- Giusti C. 2004. Association of *Helicobacter pylori* with central serous chorioretinopathy: hypotheses regarding pathogenesis. *Medical Hypotheses* 63(3): 5242-527.
- Gobert AP, Bambou JC, Werts C, Balloy V, Chignard M, Moran AP, Ferrero RL. 2004. *Helicobacter pylori* heat shock protein 60 mediates interleukin-6 production by macrophages via a toll-like receptor (TLR)-2, TLR-4, and myeloid differentiation factor 88-dependent mechanism. *J Biol Chem* 279: 245-250.
- Goddard AF, Logan RP, Atherton JC, Jenkins D, Spiller RC. 1997. Healing of duodenal ulcer after eradication of *Helicobacter heilmannii*. *Lancet* 349(9068): 1815-1816.
- Goldstein DS. 2006. Orthostatic hypotension as an early finding in Parkinson's disease. *Clin Auton Res* 16(1): 46-54.
- Goo MJ, Ki MR, Lee HR, Yang HJ, Yuan DW, Hong IH, Park JK, Hong KS, Han JY, Hwang OK, Kim DH, Do SH, Cohn RD, Jeong KS. 2009. *Helicobacter pylori* promotes hepatic fibrosis in the animal model. *Lab Invest* 89(11): 1291-1303.
- Goodwin SC, Armstrong JA, Chilvers T, Peters M, Collins DM, Sly L, McConnell W, Harper WES. 1989. Transfer of *Campylobacter pylori* and *Campylobacter mustelae* to *Helicobacter* gen. nov. as *Helicobacter pylori* comb. Nov. and *Helicobacter mustellae* comb. Nov. respectively. *Int J Syst Bact* 39(4): 397-405.
- Graham DY and Klein PD. 2000. Accurate diagnosis of *Helicobacter pylori* 13C-urea breath test. *Gastroenterol. Clin. North Am.* 29(4): 885-893.
- Graham DY, Fischbach L. 2010. *Helicobacter pylori* treatment in era of increasing antibiotic resistance. *Gut* 59(8): 1143-1153.

- Grasso GM, Ripabelli G, Sammarco ML, Ruberto A, Iannitto G. 1996. Prevalence of *Helicobacter*-like organisms in porcine gastric mucosa: a study of swine slaughtered in Italy. *Comp Immunol Microbiol Infect Dis* 19(3): 213-217.
- Gray BM, Fontaine CA, Poe SA, Eaton KA. 2013. Complex T cell interactions contribute to *Helicobacter pylori* gastritis in mice. *Infection and Immunity*. 81 (3): 740-752.
- Greiner A, Knörr C, Qin Y, Sebald W, Schilpl A, Banchereau J, Müller-Hermelink HK. 1997. Low-grade B cell lymphoma's of mucosa-associated lymphoid tissue (MALT-type) require CD40-mediated signaling and Th2-type cytokines for *in vitro* growth and differentiation. *Am J Pathol* 150(5): 1583-1593.
- Guarner J, Kalach N, Elitsur Y, Koletzko S. 2010. *Helicobacter pylori* diagnostic tests in children: review of the literature from 1999-2009. *Eur J. Pediatr* 169(1): 15-25.
- Guiney DG, Hasegawa P, Cole SP. 2003. *Helicobacter pylori* preferentially induces interleukin 12 (IL-12) rather than IL-6 or IL-10 in human dendritic cells. *Infect Immun* 71(7): 4163-4166.
- Gulcelik NE, Kaya E, Demirbas B, Culha C, Koc G, Ozkaya M, Cakal E, Serter R, Aral Y. 2005. *Helicobacter pylori* prevalence in diabetic patients and its relationship with dyspepsia and autonomic neuropathy. *J Endocrinol Invest* 28(3): 214-217.
- Gürer MA, Erel A, Erbas D, Caglar K, Atahan C. 2002. The seroprevalence of *Helicobacter pylori* and nitric oxide in acne rosacea. *Int j Dermatol* 41(11): 768-770.
- Haesebrouck F, Pasmans F, Flahou B, Chiers K, Baele M, Meyns T, Decostere A, Ducatelle R. 2009. Gastric helicobacters in domesticated animals and non-human primates and their significance for human health. *Clin Microbiol Rev* 22(2): 202-223.
- Hafsi N, Volland P, Schwendy S, Rad R, Reindl W, Gerhard M, Prinz C. 2004. Human dendritic cells respond to *Helicobacter pylori*, promoting NK cell and Th1-effector responses *in vitro*. *J Immunol* 173: 1249-1257.
- Happonen I, Linden J, Saari S, Karjalainen M, Hänninen ML, Jalava K, Westermarck E. 1998. Detection and effects of helicobacters in healthy dogs and dogs with signs of gastritis. *J Am Vet Med Assoc* 213(12): 1767-1774.
- Harper CG, Feng Y, Xu S, Taylor NS, Kinsel M, Dewhirst FE, Paster BJ, Greenwell M, Levine G, Rogers A, Fox JG. 2002. *Helicobacter cetorum* sp. nov., a urease-positive *Helicobacter* species isolated from dolphins and whales. *J Clin Microbiol* 40(12): 4536-4543.
- Harris AG, Hinds FE, Beckhouse AG, Kolesnikow T, Hazell SL. 2002. Resistance to hydrogen peroxide in *Helicobacter pylori*: role of catalase (KatA) and Fur, and functional analysis of a novel gene product designated 'KatA-associated protein', KapA (HP 0874). *Microbiology* 148(12): 3813-3825.
- Harris PR, Wright SW, Serrano C, Riera F, Duarte I, Torres J, Pena A, Rollàn A, Viviane P, Guiraldes E, Schmitz JM, Lorenz RG, Novak L, Smythies LE, Smith PD. 2008. *Helicobacter pylori* gastritis in children is associated with a regulatory T cell response. *Gastroenterology* 134(2): 491-499.
- Hashim H, Azmin S, Razlan H, Yahya NW, Tan HJ, Manaf MR, Ibrahim NM. 2014. Eradication of *Helicobacter pylori* infection improves levodopa action, clinical symptoms and quality of life in patients with Parkinson's disease. *PLoS One* 9(11): e112330.
- Hasni S, Ippolito A, Illei GG. 2011. *Helicobacter pylori* and autoimmune diseases. *Oral Dis* 17(7): 621-627.
- He C, Yang Z, Lu NH. 2014. *Helicobacter pylori* infection and diabetes: Is it a myth or fact? *World Gastroenterol* 20(16): 4607-4617.
- Heilmann KL and Borchard F. 1991. Gastritis due to spiral shaped bacteria other than *Helicobacter pylori*: clinical, histological and ultrastructural findings. *Gut*. 32(2): 137-140.
- Hellemans A, Chiers K, Decostere A, De Bock M, Haesebrouck F, Ducatelle R. 2007. Experimental infection of pigs with "*Candidatus Helicobacter suis*". *Vet Res Commun* 31(4): 385-395.
- Hellemans A, Chiers K, De Bock M, Decostere A, Haesebrouck F, Ducatelle R, Maes D. 2007. Prevalence of "*Candidatus Helicobacter suis*" in pigs of different ages. *Vet Rec* 161(6): 182-192.
- Herbrink P and van Doorn LJ. 2000. Serological methods for diagnosis of *Helicobacter pylori* infection and monitoring of eradication therapy. *Eur J Clin Microbiol Infect Dis* 19(3): 164-173.
- Hessing MJ, Geudeke MJ, Scheepens CJ, Tielen MJ, Schouten WG, Wiepkema PR. 1992. Mucosal lesions in the pars esophagea in swine: prevalence and the effect of stress. *Tijdschr Diergeneeskde* 117(15-16): 445-450.

- Hino M, Yamane T, Park K, Takubo T, Ohta K, Kitagawa S, Higuchi K, Arakawa T. 2003. Platelet recovery after eradication of *Helicobacter pylori* in patients with idiopathic thrombocytopenic purpura. *Ann Hematol* 82(1): 30-32.
- Hirschl AM and Rotter ML. 1996. Serological tests for monitoring *H. pylori* eradication treatment. *J Gastroenterol* 31(9): 33-36.
- Hirtz D, Thurman DJ, Gwinn-Hardy K, Mohamed M, Chaudhuri AR, Zalutsky R. 2007. How common are the “common” neurologic disorders? *Neurology* 68(5): 326-337.
- Hitzlet I, Kohler E, Engler DB, Yazgan AS, Müller A. 2012. The role of Th cell subsets in the control of *Helicobacter* infections and in T cell-driven gastric immunopathology. *Front Immunol* 3: 142.
- Hoffmeister A, Rothenbacher D, Bode G, Persson K, Marz W, Nauck MA, Brenner H, Hombach V, Koenig W. 2001. Current infection with *Helicobacter pylori*, but not seropositivity to *Chlamidia pneumoniae* or cytomegalovirus, is associated with an atherogenic, modified lipid profile. *Arterioscler Thromb vasc Biol*. 21(3): 427-432.
- Hong L, Zhao Y, Han Y, Guo W, Wang J, Li X, Han Y, Fan D. 2007. Reversal of migraine symptoms by *Helicobacter pylori* eradication therapy in patients with hepatitis-B-related liver cirrhosis. *Helicobacter* 12(4): 306-308.
- Honjo K, van Reekum, Verhoeff NP. 2009. Alzheimer’s disease and infection: do infectious agents contribute to progression of Alzheimer’s disease? *Alzheimers Dement* 5(4): 348-360.
- Horiki N, Omata F, Uemura M, Suzuki S, Ishii N, Iizuka Y, Fukuda K, Fujita Y, Katsurahara M, Ito T, Cesar GE, Imoto I, Takei Y. 2009. Annual change of primary resistance to clarithromycin among *Helicobacter pylori* isolates from 1996 through 2008 in Japan. *Helicobacter* 14(5): 86-90.
- Hoy B, Löwer M, Weydig C, Carra G, Tegtmeyer N, Geppert T, Schröder P, Sewald N, Backert S, Schneider G, Wessler S. 2010. *Helicobacter pylori* HtrA is a new secreted virulence factor that cleaves E-cadherin to disrupt intercellular adhesion. *EMBO Rep* 11(10): 798-804.
- Huang Y, Fan XG, Wang ZM, Zhou JH, Tian XF, Li N. 2004. Identification of *Helicobacter* species in human liver samples from patients with primary hepatocellular carcinoma. *J Clin Pathol* 57(12): 1273-1277.
- Huang B, Chen Y, Xie Q, Lin G, Wu Y, Feng Y, Li J, Zhuo Y, Zhang P. 2011. CagA-positive *Helicobacter pylori* strains enhanced coronary atherosclerosis by increasing serum OxLDL and HsCRP in patients with coronary heart disease. *Dig Dis Sci* 56(1): 109-114.
- Huber MR, Kumar S, Tefferi A. 2003. Treatment advances in adult immune thrombocytopenic purpura. *Ann Hematol* 82(12): 723-737.
- Hwang CY, Han HR, Youn HY. 2002. Prevalence and clinical characterization of gastric *Helicobacter* species infection in dogs and cats in Korea. *J Vet Sci* 3(2): 123-133.
- Ierardi E, Monno RA, Gentile A, Francavilla R, Burattini O, Marangi S, Pollice L, Francavilla A. 2001. *Helicobacter heilmannii* gastritis: a histological and immunohistochemical trait. *J Clin Pathol* 54(10): 774-777.
- Ilver D, Arnqvist A, Ögren J, Frick IM, Kersulyte D, Incecik ET, Berg DE, Covacci A, Engstrand L, Boren T. 1998. *Helicobacter pylori* adhesion binding fucosylated histoblood group antigens revealed by retagging. *Science* 279(5349): 373-377.
- Iwanczak B, Biernat M, Iwanczak F, Grabinska J, Matusiewicz K, Gosciniak G. 2012. The clinical aspects of *Helicobacter heilmannii* infection in children with dyspeptic symptoms. *J Physiol Pharmacol* 63(2): 133-136.
- Janssen M, Dijkmans BA, van der Sluys FA, van der Wielen JG, Havenga K, Vandenbroucke JP, Lamers CB, Zwinderman AH, Cats A. 1992. Upper gastrointestinal complaints and complications in chronic rheumatic patients in comparison with other chronic diseases. *Br J Rheumatol* 31(11): 747-752.
- Jalava K, On SL, Vandamme PA, Happonen I, Sukura A, Hänninen ML. 1998. Isolation and identification of *Helicobacter* spp. from canine and feline gastric mucosa. *Appl. Environ. Microbiol.* 64(10): 3998-4006.
- Jeon CY, Haan MN, Cheng C, Clayton ER, Mayeda ER, Miller JW, Aiello AE. 2012. *Helicobacter pylori* infection is associated with an increased rate of diabetes. *Diabetes Care* 35(3): 520-525.
- Joosten M, Flahou B, Meyns T, Smet A, Arts J, De Cooman L, Pasmans F, Ducatelle R, Haesebrouck F. 2013. Case report: *Helicobacter suis* infection in a pig veterinarian. *Helicobacter* 18(5): 392-396.
- Joosten M, Lindén S, Rossi M, Tay ACY, Skoog E, Padra M, Peters F, Perkins T, Vandamme P, Van Nieuwerburgh F, D’Herde K, Van den Broeck W, Flahou B, Deforce D, Ducatelle R, Marshall B, Haesebrouck F, Smet A. 2015. Divergence between the Highly Virulent Zoonotic

Pathogen *Helicobacter heilmannii* and Its Closest Relative, the Low-Virulence "*Helicobacter ailurogastricus*" sp. nov. Infect Immun 84 (1): 293-306.

Josenhans C, Labigne A, Suerbaum S. 1995. Compatative ultrastructural and functional studies of *Helicobacter pylori* and *Helicobacter mustelae* flagellin mutants: both flagellin subunits, FlaA and FlaB, are necessary for full motility of *Helicobacter* species. J Bacteriol 177(11): 3010-3020.

Kaklikkaya N, Ozgur O, Aydin F, Cobanoglu U. 2002. *Helicobacter heilmannii* as a causative agent of chronic active gastritis. Scand J Infect Dis 34(10): 768-770.

Kannarat GT, Boss JM, Tansey MG. 2013. The role of innate and adaptive immunity in Parkinson's disease. J. Parkinson's Dis. 3(4): 493-514.

Kang JM, Kim M, Lee BH, Park HK, Jo HJ, Shin CM, Lee SH, Song IS. 2011. Risk factors for peptic ulcer bleeding in terms of *Helicobacter pylori*, NSAID's, and antiplatelet agents. Scand J Gastroenterol 46(11): 1295-1301.

Karzewska E, Klesiewics K, Nowak P, Sito E, Skiba I, Zwolinska-Wcislo M, Mach T, Budak A. 2014. Empirical versus targeted treatment of *Helicobacter pylori* infections in Southern Poland according to the results of local antimicrobial resistance monitoring, 63:799-807, Chapter 12. In Roesler BM (ed), Trends in *Helicobacter pylori* infection, Published April 3, 2014.

Kato T, Yagi N, Kamada T, Shimbo T, Watanabe H, Ida K; Study Group for Establishing Endoscopic Diagnosis of Chronic Gastritis. 2013. Study group for establishing endoscopic diagnosis of chronic gastritis: Diagnosis of *Helicobacter pylori* infection in gastric mucoas by endoscopic features: a multicenter prospective study. Dig Endosc 25(5): 508-518.

Kersulyte D, Rossi M, Berg DI. 2013. Sequence divergence and conservation in genomes of *Helicobacter cetorum* strains from a dolphin and a whale. PlosOne 8(12): e83177.

Ki MR, Goo MJ, Park JK, Hong IH, Ji AR, Han SY, You SY, Lee EM, Kim AY, Park SJ, Lee HJ, Kim SY, Jeong KS. 2010. *Helicobacter pylori* accelerates hepatic fibrosis by sensitizing transforming growth factor- β 1-induced inflammatory signaling. Lab Invest 90(10): 1507-1516.

Kim SJ, Chang JH, Chung SI, Yum SJ. 1999. Molecular cloning and characterization of the *Helicobacter pylori* flhD gene, an essential factor in flagellar structure and motility. J Bacteriol 181(22): 6969-6976.

Kim KM, Lee SG, Park MG, Song JY, Kang HL, Lee WK, Cho MJ, Rhee KH, Youn HS, Baik SC. 2007. Gamma glutamyltranspeptidase of *Helicobacter pylori* induces mitochondria-mediated apoptosis in AGC cells. Biochem Biophys Res Comm 355(2): 562-567.

Kim KM, Lee SG, Kim JM, Kim DS, Song JY, Kang HL, Lee WK, Cho MJ, Rhee KH, Youn HS, Baik SC. 2010. *Helicobacter pylori* gamma-glutamyltranspeptidase induces cell cycle arrest at G1-S phase transition. J Microbiol 48(3): 372-377.

Kim SY, Buckwalter M, Soreq H, Vezzani A, Kaufer D. 2012. Blood-brain barrier dysfunction induced inflammatory signaling in brain pathology and epileptogenesis. Epilepsia 53(6): 37-44.

Kim DJ, Park JH, Franchi L, Backert S, Nunez G. 2013. The Cag pathogenicity island and interaction between TLR2/NOD2 and NLRP3 regulate IL-1 β production in *Helicobacter pylori* infected dendritic cells. Eur J Immunol 43: 2650-2658.

Kimura M, Goto S, Wada A, Yahiro K, Niidome T, Hatakeyama T, Aoyagi H, Hirayama T, Kondo T. 1999. Vacuolating cytotoxin purified from *Helicobacter pylori* causes mitochondrial damage in human gastric cells. Microb Pathog 26(1): 45-52.

Kivistö R, Linros J, Rossi M, Rautelin H, Hänninen ML. 2010. Characterization of multiple *Helicobacter bizzozeronii* isolates from a finnish patient with severe dyspeptic symptoms and chronic active gastritis. Helicobacter 15(1): 58-66.

Knörr C, Amrehn C, Seeberger H, Rosenwald A, Stilgenbauer S, Ott G, Müller-Hermelink HK, Greiner A. 1999. Expression of costimulatory molecules in low-grade mucosa-associated lymphoid tissue-type lymphomas *in vivo*. Am J Pathol 155(6): 2019-2027.

Kondadi PK, Pacini C, Revez J, Hänninen ML, Rossi M. 2013. Contingency nature of *Helicobacter bizzozeronii* oxygen-insensitive NAD(P)H-nitroreductase (HBZC1_00960) and its role in metronidazole resistance. Vet Res.16(44): 56.

Konturek PC, Brzozowski T, Konturek SJ. 2011. Stress and the gut: Pathophysiology, clinical consequences, diagnostic approach and treatment options. J Physiol Pharmacol 62: 591-599.

Kopta LA, Paquette JA, Bowersock TL, Choromanski LJ, Godbee TK, Galvin JE, Foss DL. 2010. Information of *Helicobacter suis* in pig-producing regions of North America. Conference of Research Workers in Animal Diseases 2010, Chicago, Illinois.

Kordower JH, Olanow CW, Dodiya HB, Chu Y, Beach TG, Adler CH, Halliday GM, Bartus RT. 2013. Disease duration and the integrity of the nigrostriatal system in Parkinson's disease. Brain 136(8): 2419-2431.

- Korkmaz H, Kesli R, Karabagli P, Terzi Y. 2013. Comparison of the diagnostic accuracy of the five different stool antigen tests for the diagnosis of *Helicobacter pylori* infection. *Helicobacter* 18(5): 384-391.
- Kountouras J, Mylopoulos N, Boura P, Bessas C, Chatzopoulos D, Venizelos J, Zavos C. 2001. Relationship between *Helicobacter pylori* infection and glaucoma. *Ophthalmology* 108(3): 599-604.
- Kountouras J, Mylopoulos N, Chatzopoulos D, Zavos C, Boura P, Konstas AG, Venizelos J. 2002. Eradication of *Helicobacter pylori* may be beneficial in the management of chronic open angle glaucoma. *Arch Intern Med* 162(11): 1237-1244.
- Kountouras J, Mylopoulos N, Konstas AG, Zavos C, Chatzopoulos D, Boukla A. 2003. Increased levels of *Helicobacter pylori* IgG antibodies in aqueous humor of patients with primary open-angle and exfoliation glaucoma. *Graefes Arch Clin Exp Ophthalmol*. 241(11): 884-890.
- Kountouras J, Deretzi G, Zavos C, Karatzoglou P, Touloumis L, Nicolaides T, Chatzopoulos D, Venizelos. 2005. Association between *Helicobacter pylori* infection and acute inflammatory demyelinating polyradiculoneuropathy. *Eur J Neurol* 12(2): 139-143.
- Kountouras J, Tsolaki M, Gavalas E, Boziki M, Zavos C, Karatzoglou P, Chatzopoulos D, Venizelos I. 2006. Relationship between *Helicobacter pylori* infection and Alzheimer disease. *Neurology* 66(6): 938-940.
- Kountouras J, Boziki M, Gavalas E, Zavos C, Deretzi G, Grigoriadis N, Tsolaki M, Chatzopoulos D, Katsinelos P, Tzilves D, Zabouri A, Michailidou I. 2009a. Increased cerebrospinal fluid *Helicobacter pylori* antibody in Alzheimer's disease. *Int J Neurosci* 119(6): 765-777.
- Kountouras J, Boziki M, Gavalas E, Zavos C, Deretzi G, Chatzigeorgiou S, Katsinelos P, Grigoriadis N, Giartza-Taxidou E, Venizelos I. 2010. Five-year survival after *Helicobacter pylori* eradication in Alzheimer's disease patients. *Cogn Behav Neurol* 23(3): 199-204.
- Kowalski M, Konturek PC, Pieniazek P, Karczewska E, Kluczka A, Grove R, Kranig W, Nasseri R, Thale J, Hahn EG, Konturek SJ. 2001. Prevalance of *Helicobacter pylori* infection in coronary artery disease and effect of its eradication on coronary lumen reduction after percutaneous coronary angioplasty. *Dig Liver Dis* 33(3): 222-229.
- Kowalski M, Rees W, Konturek PC, Grove R, Scheffold T, Meixner H, Brunec M, Franz N, Konturek JW, Pieniazek P, Hahn EG, Konturek SJ, Thale J, Warnecke H. 2002. Detection of *Helicobacter pylori* specific DNA in human atheromatous coronary arteries and its association to prior myocardial infarction and unstable angina. *Dig Liver Dis* 34(6): 398-402.
- Krakowka S, Eaton KA, Rings DM, Argenzio RA. 1998. Production of gastroesophageal erosions and ulcers (GEU) in gnotobiotic swine monoinfected with fermentative commensal bacteria and fed high-carbohydrate diet. *Veterinary Pathology* 35(4): 274-282.
- Kratzer I, Vasiljevic A, Rey C, Fevre-Montange M, Saunders N, Strazielle N, Ghersi-Egea JF. 2012. Complexity and developmental changes in the expression pattern of claudins at the blood-CSG barrier. *Histochem Cell Biol* 138(6): 861-879.
- Kucukazman M, Yavuz B, Sacikara M, Asilturk Z, Ata N, Ertugrul DT, Yalcin AA, Yenigun EC, Kizilca G, Okten H, Akin KO, Nazligul Y. 2009. The relationship between updated Sydney System score and LDL cholesterol levels in patients infected with *Helicobacter pylori*. *Dig Dis Sci* 54(3): 604-607.
- Kusters JG and Kuipers EJ. 2001. Antibiotic resistance of *Helicobacter pylori*. *Journal of Applied Microbial* 90(S6): 134S-144S.
- Kusters JG, Van Vliet AHM, Kuipers EJ. 2006. Pathogenesis of *Helicobacter pylori* infection. *Clin Microbiol Rev* 19(3): 449-490.
- Larussa T, Leone I, Suraci E, Imeneo M, Luzzza M. 2015. Review article: *Helicobacter pylori* and T Helper Cells: mechanisms of immune escape and tolerance. *J Immun Research* ID 981328.
- Lash JG and Genta RM. 2013. Adherence to the Sydney System guidelines increases the detection of *Helicobacter* gastritis and intestinal metaplasia in 400738 sets of gastric biopsies. *Aliment Pharmacol Ther* 38(4): 424-431.
- Laurila A, Bloigu A, Nayha S, Hassi J, Leinonen M, Saikku P. 1999. Association of *Helicobacter pylori* infection with elevated serum lipids. *Atherosclerosis* 142(1): 207-210.
- Lee E, Burnett CA, Lulich N, Cameron LL, Sestito JP. 2002. Proportionate mortality of crop and livestock farmers in the United States, 1984-1993. *Am J Ind Med* 42(5): 410-420.
- Lee H, Pienaar IS. 2014. Disruption of the blood-brain barrier in Parkinson's disease: curse or route to cure? *Front Biosc (Landmark Ed)* 19: 272-280.
- Lertsethtakarn P, Ottemann KM, Hendrixson DR. 2011. Motility and chemotaxis in *Campylobacter* and *Helicobacter*. *Annu Rev Microbiol* 65: 389-410.

- Li W, Minohara M, Su JJ, Matsuoka T, Osoegawa M, Ishizu T, Kira J. 2007. *Helicobacter pylori* infection is a potential protective factor against conventional multiple sclerosis in the Japanese population. *J Neuroimmunol* 184(1-2): 227-231.
- Li W, Minohara M, Piao H, Matsushita T, Masaki K, Matsuoka T, Isobe N, Su JJ, Ohyagi Y, Kira J. 2009. Association of anti-*Helicobacter pylori* neutrophil-activating protein antibody response with anti-aquaporin-4 autoimmunity in Japanese patients with multiple sclerosis and neuromyelitis optica. *Mult Scler* 15(12): 1411-1421.
- Liang J, Ducatelle R, Pasmans F, Smet A, Haesebrouck F, Flahou B. 2013. Multilocus sequence typing of the porcine and human gastric pathogen *Helicobacter suis*. *J Clin Microbiol* 51(3): 920-926.
- Lindberg JE, Ogle B. 2001. Digestive Physiology of Pigs: Proceedings of the 8th Symposium. Chapter 69: 249-252.
- Liu C, Smet A, Blaecher C, Flahou B, Ducatelle R, Linden S, Haesebrouck F. 2014. Gastric *de novo* expression and spasmolytic polypeptide-expressing metaplasia during *Helicobacter heilmannii* infection. *Infect and Immun* 82(8): 3227-3239.
- Liu C, Rossi M, Linden S, Padra M, Blaecher C, Bauwens E, Joosten M, Flahou B, Van den Broucke W, Ducatelle R, Haesebrouck F, Smet A. 2016. The *Helicobacter heilmannii* *HofE* and *HofF* Genes are Essential for Colonization of the Gastric Mucosa and Play a Role in IL-1 β -Induced Gastric MUC13 Expression. *Helicobacter*. doi: 10.1111/hel.12307.
- Logan RP. 1998. Urea breath tests in the management of *Helicobacter pylori* infection. *Gut* 43(1): 47-50.
- Long Y, Gao C, Qiu W, Hu X, Shu Y, Peng F, Lu Z. 2013. *Helicobacter pylori* infection in neuromyelitis optica and multiple sclerosis. *Neuroimmunomodulation* 20(2): 107-112.
- Longo-Mbenza B, Nkondi Nsenga J, Vangu Ngoma D. 2007. Prevention of the metabolic syndrome insulin resistance and the atherosclerotic diseases in Africans infected with *Helicobacter pylori* infection and treated by antibiotics. *Int J Cardiol* 121(3): 229-238.
- Löwer M, Geppert T, Schneider P, Hoy B, Wessler S, Schneider G. 2011. Inhibitors of *Helicobacter pylori* protease HtrA found by 'virtual ligand' screening combat bacterial invasion of epithelia. *PLoS One* 6: e17986.
- Lundgren A, Strömberg E, Sjöling A, Lindholm C, Enarsson K, Edebo A, Johnsson E, Suri-Payer E, Larsson P, Rudin A, Svennerholm AM, Lundin BS. 2005. Mucosal FOXP3-expressing CD4+CD25 high regulatory T cells in *Helicobacter pylori*-infected patients. *Infection and immunity* 73 (1): 523-531.
- Luther J, Dave M, Higgings PD, Kao JY. 2010. Association between *Helicobacter pylori* infection and inflammatory bowel disease: a meta-analysis and systematic review of the literature. *Inflamm Bowel Dis* 16(6): 1077-1084.
- Luzza F, Oderda G, Maletta M, Imeneo M, Mesuraca L, Chioboli E, Lerro P, Guandalini S, Pallone F. 1997. Salivary immunoglobulin G assay to diagnose *Helicobacter pylori* infection in children. *J Clin Microbiol* 35(12): 3358-3360.
- Luzza F, Parrello T, Monteleone G, Sebkova L, Romano M, Zarrilli R, Imeneo M, Pallone F. 2000. Up-regulation of IL-17 is associated with bio-active IL-8 expression in *Helicobacter pylori*-infected human gastric mucosa. *The Journal of Immunology* 165 (9): 5332-5337.
- Maeda S, Akanuma M, Mitsuno Y, Hirata Y, Ogura K, Yoshida H, Shiratori Y, Omata M. 2001. Distinct mechanisms of *Helicobacter pylori*-mediated NF-kappa B activation between gastric cancer cells and monocytic cells. *J Biol Chem* 276: 44856-44864.
- Magen E, Mishal J, Schlesinger M, Scharf S. 2007. Eradication of *Helicobacter pylori* infection equally improves chronic urticaria with positive and negative autologous serum skin test. *Helicobacter* 12(5): 567-571.
- Magen E, Delgado JS. 2014. *Helicobacter pylori* and skin autoimmune diseases. *World Journal of Gastroenterol* 20(6): 1510-1516.
- Mahdavi J, Sonden B, Hurtig M, Olfat FO, Forsberg L, Roche N, Angstrom J, Larsson T, Teneberg S, Karlsson KA, Altraja S, Wadström T, Kersuluyte D, Berg DE, Dubois A, Petersson C, Magnusson KE, Norberg T, Lindh F, Lundskog BB, Arnqvist A, Hammarström L, Boren T. 2002. *Helicobacter pylori* SabA adhesion in persistent infection and chronic inflammation. *Science* 297(5581): 573-578.
- Majka J, Rog T, Konturek PC, Konturek SJ, Bielanski W, Kowalsky M, Szczudlik A. 2002. Influence of chronic *Helicobacter pylori* infection on ischemic cerebral stroke risk factors. *Med Sci Monit* 8(10): CR675-CR684.
- Malaguarnera M, Bella R, Alagona G, Ferri R, Carnemolla A, Pennisi G. 2004. *Helicobacter pylori* and Alzheimer's disease: a possible link. *Eur J Int Med* 15(4): 381-386.
- Malaty HM, Logan ND, Graham DY, Ramchatesingh JE, Reddy SG. 2000. *Helicobacter pylori* infection in asymptomatic children: comparison of diagnostic tests. *Helicobacter* 5(3): 155-159.

- Malfertheiner P, Megraud F, O'Morain C, Bazzoli F, El-Omar E, Graham D, Hunt R, Rokkas T, Vakil N, Kuipers EJ. 2007. Current concepts in the management of *Helicobacter pylori* infection: the Maastricht III consensus report. *Gut* 56(6): 772-781.
- Malfertheiner P, Megraud F, O'Morain CA, Atherton J, Axon AT, Bazzoli F, Gensini GF, Gisbert JP, Graham DY, Rokkas T, El-Omar EM, Kuipers EJ, European Helicobacter Study Group. 2012. European *Helicobacter* study group: management of *Helicobacter pylori* infection: the Maastricht IV/Florence consensus report. *Gut* 61(5): 646-664.
- Mandell L, Moran AP, Cocchiarella A, Houghton J, Taylor N, Fox JG, Wang TC, Kurt-Jones EA. 2004. Intact gram-negative *Helicobacter pylori*, *Helicobacter felis*, and *Helicobacter hepaticus* bacteria activate innate immunity via toll-like receptor 2 but not toll-like receptor 4. *Infect Immun* 72: 6446-6454.
- Marshall BJ, Warren JR. 1984. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet*. 1(8390): 1311-1315.
- Martin Hübner A, Tenbaum SP. 2008. Complete remission of palmoplantar psoriasis through *Helicobacter pylori* eradication: a case report. *Clin Exp Dermatol* 33(3): 339-340.
- Martin ME, Bhatnagar S, George MD, Paster BJ, Canfield DR, Eisen JA, Solnick JV. 2013. The impact of *Helicobacter pylori* infection on the gastric microbiota of the rhesus macaque. *PLoS ONE* 8(10): e76375.
- Mashimo M, Nishikawa M, Higuchi K, Hirose M, Wei Q, Haque A, Sasaki E, Shiba M, Tominaga K, Watanabe T, Fujiwara Y, Arakawa T, Inoue M. 2006. Production of reactive oxygen species in peripheral blood is increased in individuals with *Helicobacter pylori* infection and decreased after its eradication. *Helicobacter* 11(4): 266-271.
- Matsui H, Takahashi T, Murayama SY, Uchiyama I, Yamaguchi K, Shigenobu S, Matsumoto T, Kawakubo M, Horiuchi K, Ota H, Osaki T, Kamiya S, Smet A, Flahou B, Ducatelle R, Haesebrouck F, Takahashi S, Nakamura S, Nakamura M. 2014. Development of new PCR primers by comparative genomics for the detection of *Helicobacter suis* in gastric biopsy specimens. *Helicobacter* 19(4): 260-271.
- Matsumoto T, Kawakubo M, Shiohara M, Kumagai T, Hidaka E, Yamauchi K, Oana K, Matsuzawa K, Ota H, Kawakami Y. 2009. Phylogeny of a novel "*Helicobacter heilmannii*" organism from a Japanese patient with chronic gastritis based on DNA sequence analysis of the 16S rRNA and urease genes. *J Microbiol*. 47(2): 201-207.
- Matsumoto T, Kawakubo M, Akamatsu T, Koide N, Ogiwara N, Kubota S, Sugano M, Kawakami Y, Katsuyama T, Ota H. 2014. *Helicobacter heilmannii* sensu stricto-related gastric ulcers: a case report. *World J Gastroenterol*. 20(12): 3376-3382.
- Mayer EA, Knight R, Mazmanian SK, Cryan JF, Tillisch K. 2014. Gut microbes and the brain: Paradigm shift in neuroscience. *Journal of neuroscience* 34(46): 15490-15496.
- McGee DJ, Langford ML, Watson EL, Carter JE, Chen YT, Ottemann KM. 2005. Colonization and inflammation deficiencies in Mongolian gerbils infected by *Helicobacter pylori* chemotaxis mutants. *Infect immune* 73(3): 1820-1827.
- McGuckin MA, Linden SK, Sutton P, Florin TH. 2011. Mucin dynamics and enteric pathogens. *Nat Rev Microbiol* 9(4): 265-278.
- McNaught KS, Olanow CW. 2003. Proteolytic stress: a unifying concept for etiopathogenesis of Parkinson's disease. *Ann Neurol* 53(3): S73-S84.
- Meeker RB, Bragg DC, Poulton W, Hudson L. 2012. Transmigration of macrophages across the choroid plexus epithelium in response to the feline immunodeficiency virus. *Cell and Tissue Research* 347(2): 443-455.
- Mégraud F. 2004. *H. pylori* antibiotic resistance: prevalence, importance, and advances in testing. *Gut* 53(9): 1374-1384.
- Mégraud F. 2012. The most important diagnostic modalities for *Helicobacter pylori*, now and in the future. *Eur J Gastroenterol Hepatol* 9(1): 13-15.
- Melnichouk SL, Friendship RM, Dewey CE, Bildfell RJ, Smart NL. 1999. *Helicobacter*-like organisms in the stomach of pigs with and without gastric ulceration. *Swine Health and Production* 7(5): 201-205.
- Mendall MA, Goggin PM, Molineaux N, Levy J, Toosy T, Strachan D, Camm AJ, Northfield TC. 1994. Relation of *Helicobacter pylori* infection and coronary heart disease. *Br Heart J* 71(5): 437-439.
- Mendes EN, Queiroz DM, Rocha GA, Moura SB, Leite VH, Fonseca ME. 1990. Ultrastructure of a spiral microorganism from pig gastric mucosa (*Gastrospirillum suis*). *J Med Microbiol* 33(1): 61-66.

- Mendes EN, Queiroz DM, Rocha GA, Nogueira AM, Carvalho AC, Lage AP, Barbosa AJ. 1991. Histopathological study of porcine gastric mucosa with and without a spiral bacterium ("*Gastrosprillum suis*"). J Med Microbiol 35(6): 345-348.
- Mendoza ML, Martín-Rabadán P, Carrión I, Morillas JD, López-Alonso G, Díaz-Rubio M. 1993. *Helicobacter pylori* infection. Rapid diagnosis with brush cytology. Acta Cytol 37(2): 181-185.
- Mention K, Michaud L, Guimber D, Martin De Lasalle E, Vincent P, Turck D, Gottrand F. 1999. Characteristics and prevalence of *Helicobacter heilmannii* infection in children undergoing upper gastrointestinal endoscopy. J Pediatr Gastroenterol Nutr 29(5): 533-539.
- Meining A, Kroher G, Stolte M. 1998. Animal reservoirs in the transmission of *Helicobacter heilmannii*. Results of a questionnaire-based study. Scand J Gastroenterol. 33 (8): 795-798.
- Melnichouk SI, Friendship RM, Dewey CE, Bildfell RJ, Smart NL. 1999. *Helicobacter*-like organisms in the stomach of pigs with and without gastric ulceration. Swine Health Prod. 7(5): 201-205.
- Meron MK, Amital H, Shepshelovich D, Barzilai O, Ram M, Anaya JM, Gerli R, Bizzaro N, Shoenfeld Y. 2010. Infectious aspects and the etiopathogenesis of rheumatoid arthritis. Clin Rev Allergy Immunol. 38(2-3): 287-291.
- Mhyre TR, Boyd JT, Hamill RW, Maguire-Zeiss KA. 2012. Parkinson's disease. Subcell Biochem 65: 389-455.
- Millet S, Kumar S, De Boever J, Meyns T, Aluwé M, De Brabander D, Ducatelle R. 2012. Effect of particle size distribution and dietary crude fiber content on growth performance and gastric mucosa integrity of growing-finishing pigs. Vet J 192(3): 316-321.
- Milman N, Rosenstock S, Andersen L, Jorgensen T, Bonnevie O. 1998. Serum ferritin, hemoglobin, and *Helicobacter pylori* infection: a seroepidemiologic survey comprising 2794 Danish adults. Gastroenterol 115(2): 268-274.
- Mimura T, Yoshida M, Nishiumi S, Tanaka H, Nobutani K, Takenaka M, Suleiman YB, Yamamoto K, Ota H, Takahashi S, Matsui H, Nakamura M, Miki I, Azuma T. 2011. IFN- γ plays an essential role in the pathogenesis of gastric lymphoid follicles formation caused by *Helicobacter suis* infection. FEMS Immunol Med Microbiol. 63(1): 25-34.
- Mimuro H, Suzuki T, Nagai S, Rieder G, Suzuki M, Nagai T, Fujita Y, Nagamatsu K, Ishijima N, Koyasu S, Haas R, Sasakawa C. 2007. *Helicobacter pylori* dampens gut epithelial self-renewal by inhibiting apoptosis, a bacterial strategy to enhance colonization of the stomach. Cell Host Microbe 2(4): 250-263.
- Misra SP, Dwivedi M, Misra V, Gupta SC. 1993. Imprint cytology--a cheap, rapid and effective method for diagnosing *Helicobacter pylori*. Postgrad Med J 69(810): 291-295.
- Miyaji H, Azuma T, Ito S, Abe Y, Ono H, Suto H, Ito Y, Yamazaki Y, Kohli Y, Kuriyama M. 1999. The effect of *Helicobacter pylori* eradication therapy on gastric antral myoelectrical activity and gastric emptying in patients with non-ulcer dyspepsia. Aliment Pharmacol Ther. 13(11): 1473-80.
- Mizuno T, Ando T, Nobata K, Tsuzuki T, Maeda O, Watanabe O, Minami M, Ina K, Kusugami K, Peek RM, Goto H. 2005. Interleukin-17 levels in *Helicobacter pylori*-infected gastric mucosa and pathologic sequelae of colonization. World J Gastroenterol 11(40): 6305-6311.
- Mobley HL, Garner RM, Bauerfeind P. 1995. *Helicobacter pylori* nickel transport gene nixA: synthesis of catalytically active urease in *Escherichia coli* independent of growth conditions. Mol Microbiol 16(1): 97-109.
- Moens S and Vanderleyden J. 1996. Functions of bacterial flagella. Crit Rev Microbiol 22(2): 67-100.
- Mohammadi M, Czinn S, Redline R, Nedrud J. 1996. *Helicobacter*-specific cell-mediated immune responses display a predominant Th1 phenotype and promote a delayed-type hypersensitivity response in the stomachs of mice. J Immunol 156(12): 4729-4738.
- Mohebi N, Mamarabadi M, Moghaddasi M. 2013. Relation of *Helicobacter pylori* infection and multiple sclerosis in Iranian patients. Neurol Int 5(2): 31-33.
- Monzon H, Forne M, Esteve M, Rosinach M, Loras C, Espinos JC, Viver JM, Salas A, Fernandez-Banares F. 2013. *Helicobacter pylori* infection as a cause of iron deficiency anaemia of unknown origin. World J Gastroenterol 19(20): 4166-4171.
- Montemurro P, Barbuti G, Dundon WG, Del Giudice G, Rappuoli R, Colucci M, De Rinaldis P, Montecucco C, Semeraro N, Papini E. 2001. *Helicobacter pylori* neutrophil-activating protein stimulates tissue factor and plasminogen activator inhibitor-2 production by human blood mononuclear cells. J Infect Dis 183(7): 1055-1062.
- Montemurro P, Nishioka H, Dundon WG, de Bernard M, Del Giudice G, Rappuoli R, Montecucco C. 2002. The neutrophil-activating protein (HP-NAP) of *Helicobacter pylori* is a potent stimulant of mast cells. Eur J Immunol 32(3): 671-676.

- Morgner A, Lehn N, Andersen LP, Thiede C, Bennedsen M, Trebesius K, Neubauer B, Neubauer A, Stolte M, Bayerdörffer E. 2000. *Helicobacter heilmannii*-associated primary gastric low grade MALT lymphoma: complete remission after curing the infection. *Gastroenterology* 118(5): 821-828.
- Mortazavi MM, Griessenauer CJ, Adeeb N, Deep A, Bavarsad Shahripour R, Loukas M, Tubbs RI, Tubbs RS. 2014. The choroid plexus: a comprehensive review of its history, anatomy, function, histology, embryology and surgical considerations. *Child's nervous system: ChNS: official journal of the International Society for Pediatric Neurosurgery* 30(2): 205-214.
- Musumba C, Jorgensen A, Sutton L, Van Eker D, Moorcroft J, Hopkins M, Pritchard DM, Pirmohamed M. 2012. The relative contribution of NSAID's and *Helicobacter pylori* to the aetiology of endoscopically-diagnosed peptic ulcer disease: observations from a tertiary referral hospital in the UK between 2005 and 2010. *Aliment Pharmacol Ther.* 36(1): 48-56.
- Nakamura M, Murayama SY, Serizawa H, Sekiya Y, Eguchi M, Takahashi S, Nishikawa K, Takahashi T, Matsumoto T, Yamada H, Hibi T, Tsuchimoto K, Matsui H. 2007. "*Candidatus Helicobacter heilmannii*" from a Cynomolgus Monkey Induces Gastric Mucosa-Associated Lymphoid Tissue Lymphomas in C57BL/6 Mice. *Infect immun.* 75(3): 1214-1222.
- Ndip RN, MacKay WG, Farthing MJ, Weaver LT. 2003. Culturing *Helicobacter pylori* from clinical specimens: review of microbiological methods. *J Pediatr Gastroenterol Nutr.* 36(5): 616-622.
- Nielsen HH, Qiu J, Friis S, Wermuth L, Ritz B. 2012. Treatment for *Helicobacter pylori* infection and risk of Parkinson's disease in Denmark. *Eur J Neurol* 19(6): 864-869.
- Nilsson HO, Mulchandani R, Tranberg KG, Stenram U, Wadström T. 2001. *Helicobacter* species identified in liver from patients with cholangiocarcinoma and hepatocellular carcinoma. *Gastroenterology* 120(1): 323-324.
- Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H, Hashimoto N, Furuse M, Tsukita S. 2003. Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell Biol* 161(3): 653-660.
- Odaka T, Yamaguchi T, Koyama H, Saisho H, Nomura F. 2002. Evaluation of the *Helicobacter pylori* stool antigen test for monitoring eradication therapy. *Am J Gastroenterol* 97(3): 594-599.
- Odenbreit S, Püls J, Sedlmaier B, Gerland E, Fischer W, Haas R. 2000. Translocation of *Helicobacter pylori* CagA into gastric epithelial cells by type IV secretion. *Science* 287(5457): 1497-1500.
- Odenbreit S. 2005. Adherence properties of *Helicobacter pylori*: impact on pathogenesis and adaption to the host. *Int J Med Microbiol* 295(5): 317-324.
- Odenbreit S, Linder S, Gebert-Vogl B, Rieder G, Moran AP, Haas R. 2006. Interleukin-6 induction by *Helicobacter pylori* in human macrophages is dependent on phagocytosis. *Helicobacter*.11(3): 196-207.
- Odenbreit S, Swoboda K, Barwig I, Ruhl S, Boren T, Koletzko S, Haas R. 2009. Outer membrane protein expression profile in *Helicobacter pylori* clinical isolates. *Infect Immun* 77(9): 3782-3790.
- Oguri A, Ohmiya N, Taguchi A, Itoh A, Hirooka Y, Niwa Y, Maeda O, Ando T, Goto H. 2007. Rugal hyperplastic gastritis increases the risk of gastric carcinoma, especially diffuse and p53-independent subtypes. *Eur J Gastroenterol Hepatol* 19(7): 561-566.
- Okiyama Y, Matsuzawa K, Hidaka E, Sano K, Akamatsu T, Ota H. 2005. *Helicobacter heilmannii* infection: Clinical, endoscopic and histopathological features in Japanese patients. *Pathology international* 55(7): 398-404.
- Olczak AA, Seyler RW Jr, Olson JW, Maier RJ. 2003. Association of *Helicobacter pylori* antioxidant activities with host colonization proficiency. *Infect Immun* 71(1): 580-583.
- Oldenburg B, Diepersloot RJ, Hoekstra JB. 1996. High seroprevalence of *Helicobacter pylori* in diabetes mellitus patients. *Dig Dis Sci* 41(3): 458-461.
- Oleastro M and Ménard A. 2013. The role of *Helicobacter pylori* outer membrane proteins in adherence and pathogenesis. *Biology (Basel)* 2(3): 1110-1134.
- Oliva MM, Lazenby AJ, Perman JA. 1993. Gastritis associated with *Gastrosprillum hominis* in children. Comparison with *Helicobacter pylori* and review of the literature. *Modern Pathol* 6(5): 513-515.
- Onsun N, Arda Ulusal H, Su O, Beycan I, Biyik Ozkaya D, Senocak M. 2012. Impact of *Helicobacter pylori* infection on severity of psoriasis and response to treatment. *Eur J Dermatol* 22(1): 117-120.

- O'Rourke JL and Lee A. 2003. Animal models of *Helicobacter pylori* infection and disease. *Microbes infect* 5(8): 741-748.
- O' Rourke JL, Solnick JV, Neilan BA, Seidel K, Hayter R, Hansen LM, Lee A. 2004. Description of '*Candidatus Helicobacter heilmannii*' based on DNA sequence analysis of the 16S rRNA and the urease genes. *Int J Syst Evol Microbiol*.54(6): 2203-2211.
- O'Toole PW, Kostrzynska M, Trust TJ. 1994. Non-motile mutants of *Helicobacter pylori* and *Helicobacter mustelae* defective in flagellar hook production. *Mol Microbiol* 14(4): 691-703.
- Otteman KM, Miller F. 1997. Roles of motility in bacterial host interactions. *Mol Microbiol* 24(6): 1109-1117.
- Ozaslan E, Koseoglu T, Purnak T, Yildiz A. 2010. A forgotten cause of false negative rapid urease test: formalin contamination of the sample. *Hepatogastroenterology* 57: 99-100.
- Oshima T, Ozono R, Yano Y, Oishi Y, Teragawa H, Higashi Y, Yoshizumi M, Kambe M. 2005. Association of *Helicobacter pylori* infection with systemic inflammation and endothelial dysfunction in healthy male subjects. *J Am Coll Cardiol* 45(8): 1219-1222.
- Papamichael KX, Papaioannou G, Kerge H, Roussos A, Mantzaris GJ. 2009. *Helicobacter pylori* infection and endocrine disorders: Is there a link? *World J Gastroenterol* 15(22): 2701-2707.
- Papamichael K, Konstantopoulos P, Mantzaris. 2014. *Helicobacter pylori* infection and inflammatory bowel disease: Is there a link? *World Journal of Gastroenterology*. 20(21): 6374-6385.
- Park JH, Lee BJ, Lee YS, Park JH. 2000. Association of tightly spiraled bacterial infection and gastritis in pigs. *J Vet Med Sci* 62(7): 725-729.
- Park JH, Seok SH, Baek MW, Lee HY, Kim DJ, Park JH. 2008. Gastric lesions and immune responses caused by long-term infections with *Helicobacter heilmannii* in C57BL/6 mice. *J Comp Pathol*. 139(4): 208-217.
- Park MJ, Choi SH, Kim D, Kang SJ, Chung SJ, Choi SY, Yoon DH, Lim SH, Kim YS, Yim JK, Kim JS, Jung HC. 2011. Association between *Helicobacter pylori* seropositivity and the coronary artery calcium score in a screening population. *Gut Liver* 5(3): 321-327.
- Parkinson J. 1817. An assay on the shaking Palsy. *J Neuropsychiatry Clin Neurosci*. 14(2): 223-36; discussion 222.
- Parsonnet J and Isaacson PG. 2004. Bacterial infections and Malt-lymphoma's. *N Engl J Med* 350: 213-215.
- Paster BJ, Lee A, Fox, JG, Dewhirst FE, Tordoff LA, Frasser GJ, O'Rourke JL, Taylor NS, Ferrero R. 1991. Phylogeny of *Helicobacter felis* sp. nov., *Helicobacter mustelae*, and related bacteria. *Int J Syst Bacteriol* 41(1): 31-38.
- Pay S, Simsek I, Erdem H, Dinç A. 2007. Immunopathogenesis of Behçet's disease with special emphasize on the possible role of antigen presenting cells. *Rheumatol Int* 27(5): 417-424.
- Peach HG, Bath NE, Farish SJ. 1998. *Helicobacter pylori* infection: an added stressor on iron status of women in the community. *Med J Aust* 169(4): 188-190.
- Peek RM, Thompson SA, Donahue JP, Tham KT, Atherton JC, Blaser MJ, Miller GG. 1998. Adherence to gastric epithelial cells induces expression of a *Helicobacter pylori* gene, *IceA*, that is associated with clinical outcome. *Proc Assoc Am Physicians* 110(6): 531-544.
- Peek RM, van Doorn LJ, Donahue JP, Tham KT, Figueiredo C, Blaser MJ, Miller GG. 2000. Quantitative detection of *Helicobacter pylori* gene expression *in vivo* and relationship to gastric pathology. *Infect Immun* 68(10): 5488-5495.
- Peek RM JR, Fiske C, Wilson KT. 2010. Role of innate immunity in *Helicobacter pylori*-induced gastric malignancy. *Physiol Rev* 90(3): 831-858.
- Pellicano R, Mazzaferro V, Grigioni WF, Cutufia MA, Fagoonee S, Silengo L, Rizzetto M, Ponzetto A. 2004. *Helicobacter* species sequences in liver samples from patients with and without hepatocellular carcinoma. *World J Gastroenterol* 10(4): 598-601.
- Pfeiffer RF. 2003. Gastrointestinal dysfunction in Parkinson's disease. *Lancet Neurol* 2(2): 107-116.
- Pfeiffer RF. 2011. Gastrointestinal dysfunction in Parkinson's disease. *Parkinsonism Relat Disord*. 17(1): 10-15.
- Pierantozzi M, Pietroiusti A, Sancesario G, Lunardi G, Fedele E, Giacomini P, Frasca S, Galante A, Marciani MG, Stanzione P. 2001. Reduced L-dopa absorption and increased clinical fluctuations in *Helicobacter pylori*-infected Parkinson's disease patients. *Neurol Sci* 22(1): 89-91.
- Pirouz T, Zounubi L, Keivani H, Rakhshani N, Hormazdi M. 2009. Detection of *Helicobacter pylori* in paraffin-embedded specimens from patients with chronic liver diseases, using the amplification method. *Dig Dis Sci* 54(7): 1456-1459.

- Pisani V, Stefani A, Pierantozzi M, Natoli S, Stanzione P, Franciotta D, Pisani A. 2012. Increased blood-cerebrospinal fluid transfer of albumin in advanced Parkinson's disease. *Journal of Neuroinflamm* 9: 188.
- Polenghi A, Bossi F, Fischetti F, Durigutto P, Cabrelle A, Tamassia N, Cassatella MA, Montecucco C, Tedesco F, de Bernard M. 2007. The neutrophil-activating protein of *Helicobacter pylori* crosses endothelia to promote neutrophil adhesion *in vivo*. *J Immunol* 178(3): 1312-1320.
- Ponzetto A, Pellicano R, Leone N, Berrutti M, Turrini F, Rizzetto M. 2000. *Helicobacter pylori* seroprevalence in cirrhotic patients with hepatitis B virus infection. *Neth J Med* 56(6): 206-210.
- Ponzetto A, Pellicano R, Leone N, Cutafia MA, Turrini F, Grigioni WF, D'Errico A, Mortimer P, Rizzetto M, Silengo L. 2000. *Helicobacter* infection and cirrhosis in hepatitis C virus carriage: is it an innocent bystander or a troublemaker. *Med Hypotheses* 54(2): 275-277.
- Pot RG, Stoof J, Nuijten PJ, de Haan LA, Loeffen P, Kuipers EJ, van Vliet AH, Kusters JG. 2007. UreA2B2: a second urease system in the gastric pathogen *Helicobacter felis*. *FEMS Immunol Med Microbiol*. 50(2): 273-279.
- Pounder RE and Ng D. 1995. The prevalence of *Helicobacter pylori* infection in different countries. *Aliment Pharmacol Ther* 9(2): 33-39.
- Powell FC. 2005. Clinical practice. Rosacea. *N Engl J Med* 352(8): 793-803.
- Pringsheim T, Jette N, Frolkis A, Steeves D. 2014. The prevalence of Parkinson's disease: a systematic review and meta-analysis. *Movement disorders* 29(13): 1583-1590.
- Provencio JJ, Kivisäkk P, Tucky BH, Luciano MG, Ransohoff RM. 2005. Comparison of ventricular and lumbar cerebrospinal fluid cells in non-inflammatory neurological disorder (NIND) patients. *Journal of Neuroimmunology* 163(1-2): 179-184.
- Qayoom S, Ahmad QM. 2003. Psoriasis and *Helicobacter pylori*. *Indian J Dermatol Venereol* 69(2): 133-134.
- Qin H, Tang G, Yi P, Pan X, Huang H, Chang R, Shi Z, Ashraf MA. 2016. Diagnosis of the Genus *Helicobacter* through a hemi-nested PCR assay of 16S rRNA. *Saudi Pharmaceutical Journal* doi:10.1016/j.jsps.2016.04.015.
- Quadri R, Rossi C, Catalfamo E, Masoero G, Lombardo L, Della Monica P, Rovera L, Pera A, Cavello Perib P. 2000. *Helicobacter pylori* infection in type 2 diabetic patients. *Nutr Metab Cardiovasc Dis* 10(5): 263-266.
- Queiroz DM, Rocha GA, Mendes EN, Lage AP, Carvalho AC, Barbosa AJ. 1990. A spiral microorganism in the stomach of pigs. *Vet Microbiol*. 24(2): 199-204.
- Queiroz DM, Rocha GA, Mendes EN, De Moura SB, De Oliveira AM, Miranda D. 1996. Association between *Helicobacter* and gastric ulcer disease of pars oesophagea in swine. *Gastroenterology* 111(1): 19-27.
- Queiroz DM, Saito M, Rocha GA, Rocha AM, Melo FF, Checkley W, Braga LL, Silva IS, Gilman RH, Crabtree JE. 2013. *Helicobacter pylori* infection in infants and toddlers in South America: concordance between [13C]urea breath test and monoclonal *H. pylori* stool antigen test. *J Clin. Microbiol*. 51(11): 3735-3740.
- Rabelo-Gonçalves EMA, Roesler BM, Zeitune JMR. 2015. Extragastric manifestations of *Helicobacter pylori* infection: Possible role of bacterium in liver and pancreas diseases. *World Journal of Hepatology* 7(30): 2968-2979.
- Rad R, Ballhorn W, Volland P, Eisenächer K, Mages J, Rad L, Ferstl R, Lang R, Wagner H, Schmid RM, Bauer S, Prinz C, Kirschning CJ, Krug A. 2009. Extracellular and intracellular pattern recognition receptors cooperate in the recognition of *Helicobacter pylori*. *Gastroenterol* 136: 2247-2257.
- Rad R, Brenner L, Krug A, Volland P, Mages J, Lang R, Schwendy S, Reindl W, Dossumbekova A, Ballhorn W, Wagner H, Schmid RM, Bauer S, Prinz C. 2007. Toll-like receptor dependent activation of antigen-presenting cells affects adaptive immunity to *Helicobacter pylori*. *Gastroenterology* 133: 150-163.
- Radic M. 2014. Role of *Helicobacter pylori* infection in autoimmune systemic rheumatic diseases. *World J Gastroenterol*. 20(36): 12839-12846.
- Raghavan S and Quiding-Järbrink M. 2012. Immune modulation by regulatory T cells in *Helicobacter pylori*-associated diseases. *Endocr Metab Immune Disord Drug Targets* 12(1): 71-85.
- Rahbani-Nobar MB, Javadzadeh A, Ghojzadeh L, Rafeey M, Ghorbanihaghjo A. 2011. The effect of *Helicobacter pylori* treatment on remission of idiopathic central serous chorioretinopathy. *Mol Vis* 17: 99-103.

- Ramarao N, Gray-Owen SD, Meyer TF. 2000. *Helicobacter pylori* induces but survives the extracellular release of oxygen radicals from professional phagocytes using its catalase activity. *Mol Microbiol* 38(1): 103-113.
- Ramsauer M, Krause D, Dermietzel R. 2002. Angiogenesis of the blood-brain barrier *in vitro* and the function of cerebral pericytes. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology* 16(10): 1274-1276.
- Raymond J, Kalach N, Bergeret M, Barbet JP, Benhamou PH, Gendrel D, Dupont C. 1996. Evaluation of a serological test for the diagnosis of *Helicobacter pylori* in children. *Eur J Clin Microbiol Infect Dis* 15(5): 415-417.
- Reale M, Iarlori C, Thomas A, Gambi D, Perfetti B, Nicola MDi, Onofri M. 2009. Peripheral cytokines profile in Parkinson's disease. *Brain Behaviour Immunity* 23: 55-63.
- Rebora A, Drago F, Picciotto A. 1994. *Helicobacter pylori* in patients with rosacea. *Am J Gastroenterol* 89(9): 1603-1604.
- Rees K, Stowe R, Patel S, Ives N, Breen K, Clarke CE, Ben-Shlomo Y. 2011. *Helicobacter pylori* eradication for Parkinson's disease. *Cochrane Database Syst Rev* doi10.1002/14651858.CD008453.pub2.
- Ricci C, Holton J, Vaira D. 2007. Diagnosis of *Helicobacter pylori*: invasive and non-invasive tests. *Best Pract Res Clin Gastroenterol*. 21(2): 299-3134.
- Ritch R. 2007. Natural compounds: evidence for a protective role in eye disease. *Can J Ophthalmol* 42(3): 425-438.
- Ritchie B, Brewster D, Tran CD, McNeil Y, Zacharakis B, Davidson GP, Butler RN. 2009. Lack of diagnostic accuracy of the monoclonal stool antigen test for detection of *Helicobacter pylori* infection in young Australian aboriginal children. *Pediatr Infect Dis J*. 28: 287-289.
- Robertson ID, Accioly JM, Moore KM, Driesen SJ, Pethick DW, Hampson DJ. 2002. Risk factors for gastric ulcers in Australian pigs at slaughter. *Prev Vet Med* 53(4): 293-303.
- Robinson K, Kenefeck R, Pidgeon EL, Shakib S, Patel S, Polson RJ, Zaitoun AM, Atherton JC. 2008. *Helicobacter pylori*-induced peptic ulcer disease is associated with inadequate regulatory T cell responses. *Gut* 57(10): 1375-1385.
- Robinson K and Atherton JC. 2010. *Helicobacter pylori*-induced acquired immunity and immunoregulation. In: Sutton P, Mitchell HM (eds), *Helicobacter pylori* in the 21st century, CAB International, Oxfordshire, UK, p94-115.
- Rogers AB and Fox JG. 2004. Inflammation and cancer. I. Rodent models of infectious gastrointestinal and liver cancer. *Am J Physiol gastrointest liver Physiol*. 286 (3): G361-366.
- Rokkas T, Gisbert JP, Niv Y, O'Morain C. 2015. The association between *Helicobacter pylori* infection and inflammatory bowel disease based on meta-analysis. *United European Gastroenterology Journal* 3(6): 539-550.
- Romagnani S. 2000. T-cell subsets (Th1 versus Th2). *Annals of Allergy, Asthma & Immunol* 85 (1): 9-18,21
- Roosendaal R, Vos JH, Roumen T, van Vugt R, Cattoli G, Bart A, Klaasen HL, Kuipers EJ, Vandenbroucke-Grauls CM, Kusters JG. 2000. Slaughter pigs are commonly infected with closely related but distinct gastric ulcerative lesion-inducing gastrospirilla. *J Clin Microbiol*. 38(7): 2661-2664.
- Rosenthal R, Milatz S, Krug SM, Oelrich B, Schulzke JD, Amasheh S, Günzel D, Fromm M. 2010. Claudin-2, a component of the tight junction, forms a paracellular water channel. *J Cell Sci* 123(11): 1913-1921.
- Roubaud-Baudron C, Krolak-Salmon P, Quadrio I, Mégraud F, Salles N. 2012. Impact of chronic *Helicobacter pylori* infection on Alzheimer's disease: preliminary results. *Neurobiol Aging* 33(5): 1009.e11-1009.e19.
- Rubicz R, Leach CT, Kraig E, Dhurandhar NV, Grubbs B, Blangero J, Yolken R, Göring HH. 2011. Seroprevalence of 13 common pathogens in a rapidly growing U.S. minority population: Mexican Americans from San Antonio, TX. *BMC Res Notes* 4: 433.
- Ruuska T, Vaajalahti P, Arajärvi P, Mäki M. 1994. Prospective evaluation of upper gastrointestinal mucosal lesions in children with ulcerative colitis and Crohn's disease. *J Pediatr gastroenterol Nutr* 19(2): 181-186.
- Sacca SC, Vagge A, Pulliero A, Izzotti A. 2014. *Helicobacter pylori* infection and eye diseases: a systematic review. *Medicine* 93(28): e216.
- Saez J, Belda S, Santibáñez M, Rodríguez JC, Sola-Vera J, Galiana A, Ruiz-García M, Brotons A, López-Girona E, Girona E, Sillero C, Royo G. 2012. Real-time PCR for diagnosing *Helicobacter pylori* infection in patients with upper gastrointestinal bleeding: comparison with other classical diagnostic methods. *J. Clin. Microbiol* 50(10): 3233-3237.

- Sakakibara R, Uchiyama T, Yamanishi T, Shirai K, Hattori T. 2008. Bladder and bowel dysfunction in Parkinson's disease. *J Neural Transm.* 115(3): 443-460.
- Salama NR, Hartung ML, Muller A. 2013. Life in the human stomach: persistence strategies of the bacterial pathogen *Helicobacter pylori*. *Nat Rev Microbiol* 11(6): 385-399.
- Saldinger PF, Porta N, Launois P, Louis JA, Waanders GA, Bouzourène H, Michetti P, Blum AL, Corthésy-Theulaz IE. 1998. Immunization of BALB/c mice with *Helicobacter urease* B induces a T helper 2 response absent in *Helicobacter* infection. *Gastroenterol* 115(4): 891-897.
- Salomon H. 1898. Über das Spirillum des Säugetiermagens und sein Verhalten zu den Belegzellen. *Zbl Bakt Parasitenkd Infektionskr Hyg Abt* 19: 433-442.
- Sapierzyński R, Fabisiak M, Kizerwetter-Swida M, Cywińska A. 2007. Effect of *Helicobacter* sp. infection on the number of antral gastric endocrine cells in swine. *Pol J Vet Sci.* 10(2): 65-70.
- Satin B, Del Giudice G, Della Bianca V, Dusi S, Laudanna C, Tonello F, Kelleher D, Rappuoli R, Montecucco C, Rossi F. 2000. The neutrophil-activating protein (HP-NAP) of *Helicobacter pylori* is a protective antigen and a major virulence factor. *J Exp Med* 191(9): 1467-1676.
- Sato M, Shimoyama T, Takahashi R, Kajiyama H, Sano Y, Sakaedani N, Kato A, Hirata H, Fukuda Y. 2012. Characterization and usefulness of stool antigen tests using a monoclonal antibody to *Helicobacter pylori* catalase. *J. Gastroenterol Hepatol.* 27(3): 23-28.
- Savarino V, Vigneri S, Celle GC. 1999. The ¹³C urea breath test in the diagnosis of *Helicobacter pylori* infection. *Gut* 45: 118-122.
- Sawalha AH, Schmid WR, Binder SR, Bacino DK, Harley JB. 2004. Association between systemic lupus erythematosus and *Helicobacter pylori* seronegativity. *J Rheumatol* 31(8): 1546-1550.
- Schabereiter-Gurtner C, Hirschl AM, Dragosics B, Hufnagl P, Puz S, Kovách Z, Rotter M, Makristathis A. 2004. Novel real-time PCR assay for detection of *Helicobacter pylori* infection and simultaneous clarithromycin susceptibility testing of stool and biopsy specimens. *J Clin Microbiol* 42(10): 4512-4518.
- Schildt J, Grimm T, Radke M, Lobeck H. 2000. Akute oberbauchschmerzen und hematemesis. *Monatsschr Kinderheilkd* 148: 900-901.
- Schmausser B, Andrulis M, Endrich S, Lee SK, Josenhans C, Müller-Hermelink HK, Eck M. 2004. Expression and subcellular distribution of toll-like receptors TLR4, TLR5 and TLR9 on the gastric epithelium in *Helicobacter pylori* infection. *Clin Exp Immunol* 136(3): 521-526.
- Schmees C, Prinz C, Treptau T, Rad R, Hengst L, Volland P, Bauer S, Brenner L, Schmid RM, Gerhard M. 2007. Inhibition of T cell proliferation by *Helicobacter pylori* gamma glutamyl transpeptidase. *Gastroenterology* 132(5): 1820-1833.
- Schott T, Kondadi PK, Hänninen ML, Rossi M. 2011. Comparative genomics of *Helicobacter pylori* and the human-derived *Helicobacter bizzozeronii* CIII-1 strain reveal the molecular basis of the zoonotic nature of non-pylori gastric *Helicobacter* infections in humans. *BMC Genomics*.12: 534.
- Schulz C, Schütte K, Malfertheiner P. 2014. Does *H. pylori* eradication therapy benefit patients with hepatic encephalopathy?: systematic review. *J Clin Gastroenterol* 48(6): 491-499.
- Schwab RS. 1961. Symptomatology and medical treatment of Parkinson's disease. *Int J Neurol* 2: 61-75.
- Scott DR, Marcus EA, Weeks DL, Lee A, Melchers K, Sachs G. 2000. Expression of the *Helicobacter pylori ureI* gene is required for acidic pH activation of cytoplasmic urease. *Infect Immun* 68(2): 470-477.
- Scott DR, Marcus EA, Weeks DL, Sachs G. 2002. Mechanisms of acid resistance due to the urease system of *Helicobacter pylori*. *Gastroenterology* 123(1): 187-195.
- Scotti C, Sommi P, Paschetto MV, Cappelletti D, Stivala S, Mignosi P, Savio M, Chiarelli LR, Valentini G, Bolanos-Garcia VM, Merrell DS, Franchini S, Verona ML, Bolis C, Solcia E, Manca R, Franciotta D, Casasco A, Filipazzi P, Zardini E, Vannini V. 2010. Cell-cycle inhibition by *Helicobacter pylori* L-asparaginase. *PLoS One*.5(11): e13892.
- Segal ED, Cha J, Lo J, Falkow S, Tompkins LS. 1999. Altered states: involvement of phosphotylated CagA in the induction of the host cellular growth changes of by *Helicobacter pylori*. *Proc Natl Acad, Sci USA* 96(25): 14559-14564.
- Selkoe DJ. 2002. Deciphering the genesis and fate of amyloid beta-protein yields novel therapies for Alzheimer disease. *J Clin Invest* 110(10): 1375-1381.

- Seo WJ, Park CS, Cho YJ, Cha KW, Lee SW, Lim ST, Sung YH, Baek AR. 2003. A case of gastric ulcer induced by *Helicobacter heilmannii*-like organisms. Korean J Gastroenterol 42(1): 63-66.
- Serrano C, Diaz MI, Valdivia A, Godoy A, Pena A, Rollan A, Kirberg A, Hebel E, Fierro J, Klapp G, Venegas A, Harris PR. 2007. Relationship between *Helicobacter pylori* virulence factors and regulatory cytokines as predictors of clinical outcome. Microbes Infect 9(4): 428-434.
- Seyler RW Jr, Olson JW, Maier RJ. 2001. Superoxide dismutase-deficient mutants of *Helicobacter pylori* are hypersensitive to oxidative stress and defective in host colonization. Infect Immun 69(6): 4034-4040.
- Sheng YH, Triyana S, Wang R, Das I, Gerloff K, Florin TH, Sutton P, McGuckin MA. 2013. MUC1 and MUC13 differentially regulate epithelial inflammation in response to inflammatory and infectious stimuli. Mucosal Immunol. 6(3): 557-68.
- Shi Y, Liu XF, Zhuang Y, Zhang JY, Liu T, Yin Z, Wu C, Mao XH, Jia KR, Wang FJ, Guo H, Flavell RA, Zhao Z, Liu KY, Xiao B, Guo Y, Zhang WJ, Zhou WY, Guo G, Zou QM. 2010. *Helicobacter pylori*-induced Th17 responses modulate Th1 cell responses, benefit bacterial growth, and contribute to pathology in mice. J Immunol. 184: 5121-1529.
- Shi WJ, Liu W, Zhou XY, Ye F, Zhang GX. 2013. Associations of *Helicobacter pylori* infection and cytotoxin-associated gen: A status with autoimmune thyroid diseases: a metanalysis. Thyroid 23(10): 1294-1300.
- Shibayama K, Kamachi K, Nagata N, Yagi T, Nada T, Doi Y, Shibata N, Yokoyama K, Yamane K, Kato H, Iinuma Y, Arakawa Y. 2003. A novel apoptosis-inducing protein from *Helicobacter pylori*. Mol Microbiol 47(2): 443-451.
- Shibayama K, Wachino J, Arakawa Y, Saidijam M, Rutherford NG, Henderson PJ. 2007. Metabolism of glutamine and glutathione via gamma-glutamyltranspeptidase and glutamate transport in *Helicobacter pylori*: possible significance in pathophysiology of the organism. Mol Microbiol 64(2): 396-406.
- Shibayama T, Sawaya M, Ishiguro A, Hanabata N, Yoshimura T, Fukuda S. 2011. Applicability of a rapid stool antigen test, using monoclonal antibody to catalase, for the management of *Helicobacter pylori* infection. J. Gastroenterol 46(4): 487-491.
- Shinozaki JK, Sellon RK, Cantor GH, Besser TE, Mealey KL, Vaden SL. 2002: Fecal polymerase chain reaction with 16S ribosomal RNA primers can detect the presence of gastrointestinal *Helicobacter* in dogs. J. Vet. Intern Med. 16(4): 426-432.
- Shiomi S, Toriiez A, Imamura S, Konishi H, Mitsufuji S, Iwakura Y, Yamaoka Y, Ota H, Yamamoto T, Imanishi J, Kita M. 2008. IL-17 is involved in *Helicobacter pylori*-induced gastric inflammatory responses in a mouse model. Helicobacter 13(6): 518-524.
- Showji Y, Nozowa R, Sato K, Suzuki H. 1996. Seroprevalence of *Helicobacter pylori* infection in patients with connective tissue diseases. Microbiol Immunol 40(7): 499-503.
- Shojaee Tabrizi A, Jamshidi Sh, Oghalaei A, Zahraei Salehi T, Bayati Eshkaftaki A, Mohammadi M. 2010. Identification of *Helicobacter* spp. in oral secretions vs. gastric mucosa of stray cats. Vet Microbiol. 140(1-2): 142-146.
- Simon L, Tornoczky J, Toth M, Jambor M, Sudar Z. 1989. The significance of *Campylobacter pylori* infection in gastroenterologic and diabetic practice. Orv Hetil 130(25): 1325-1329.
- Smet A, Van Nieuwerburgh F, Flahou B, Pasmans F, Deforce D, Ducatelle R, Haesebrouck F. 2011. Proceedings EHSg Dublin: Whole genome sequence of *Helicobacter heilmannii*. Helicobacter 16(S1): 89.
- Smet A, Van Nieuwerburgh F, Ledesma J, Flahou B, Deforce D, Ducatelle R, Haesebrouck F. 2013. Genome sequence of *Helicobacter heilmannii* sensu stricto ASB1 isolated from the gastric mucosa of a kitten with severe gastritis. Genome Announc 1(1): e00033-12.
- Smet A, Tay A, Thirriot F, Perkins T, Backert S, Flahou B, Ducatelle R, Skoog E, Padra M, Linden S, Rossi M, Haesebrouck F. 2014. Comparative genomics of gastric *Helicobacter* species: analysis of outer membrane proteins. Helicobacter 19(S1), 97.
- Smith MF, Mitchell A, Li G, Ding S, Fitzmaurice AM, Ryan K, Crowe S, Goldberg JB. 2003. Toll-like receptor (TLR)2 and TLR5, but not TLR4, are required for *Helicobacter pylori*-induced NF-kappa B activation and chemokine expression by epithelial cells. J Biol Chem. 278(35): 32552-32560.
- Smoot DT, Mobley HL, Chippendale GR, Lewison JF, Resau JH. 1990. *Helicobacter pylori* urease activity is toxic to human gastric epithelial cells. Inf Immun 58(6): 1992-1994.
- Smyk DS, Koutsoumpas AL, Mytilinaiou MG, Rigopoulou EI, Sakkas LI, Bogdanos DP. 2014. *Helicobacter pylori* and autoimmune disease: Cause or bystander. World J Gastroenterol. 20(3): 613-629.

- Smythies LE, Waites KB, Lindsey JR, Harris PR, Ghiara P, Smith PD. 2000. *Helicobacter pylori*-induced mucosal inflammation is Th1 mediated and exacerbated in IL-4, but not IFN-gamma, gene-deficient mice. *J Immunol* 165(2): 1022-1029.
- Solnick JV. 2003. Clinical significance of *Helicobacter* species other than *Helicobacter pylori*. *Clin Infect Dis* 36(3): 349-354.
- Sommer F, Faller G, Röllinghoff M, Kirchner T, Mak TW, Lohoff M. 2001. Lack of gastritis and of an adaptive immune response in interferon regulatory factor-1-deficient mice infected with *Helicobacter pylori*. *Eur J Immunol* 31(2): 396-402.
- Souto R and Colombo AP. 2008. Detection of *Helicobacter pylori* by polymerase chain reaction in the subgingival biofilm and saliva of non-dyspeptic periodontal patients. *J. Periodontol* 79(1): 97-103.
- Stalke P, Zóttowska A, Orłowski M, Ellert-Zygadłowska J, Witczak-Malinowska K, Michalska Z, Lakomy EA, Trocha H, Stepiński J. 2001. Correlation between liver damage and degree of gastric mucosa colonization by *Helicobacter pylori* in subjects with parenchymatous liver damage. *Med Sci Monit* 7(1): 271-276.
- Stasi T, Rossi Z, Stipa E, Amadori S, Newland AC, Provan D. 2005. *Helicobacter pylori* eradication in the management of patients with idiopathic thrombocytopenic purpura. *Am J Med* 118(4): 414-419.
- Sterzenbach T, Lee SK, Brenneke B, von Goetz F, Schauer DB, Fox JG, Suerbaum S, Josenhans C. 2007. Inhibitory effect of enterohepatic *Helicobacter hepaticus* on innate immune responses of mouse intestinal epithelial cells. *Infect Immun* 75(6): 2717-2728.
- Stingl K, Altendorf K, Bakker EP. 2002. Acid survival of *Helicobacter pylori*: how does urease activity trigger cytoplasmic pH homeostasis? *Trends Microbiol* 10(2): 70-74.
- Stoffel MH, Friess AE, Burnens A, Schmassmann A, Neiger R. 2000. Distinction of gastric *Helicobacter* spp. in humans and domestic pets by scanning electron microscopy. *Helicobacter*. 5(4): 232-239.
- Stolp HB, Dziegielewska KM. 2009. Review: Role of developmental inflammation and blood-brain barrier dysfunction in neurodevelopmental and neurodegenerative disorders. *Neuropathology and applied Neurobiology* 35(2): 132-146.
- Stolte M, Eidt S. 1993. Healing gastric MALT lymphomas by eradicating *H. pylori*. *Lancet* 342(8871): 568.
- Stolte M, Wellens E, Bethke B, Ritter M, Eidt H. 1994. *Helicobacter heilmannii* (formerly *Gastrospirillum hominis*) gastritis: an infection transmitted by animals? *Scand. J. Gastroenterol.* 29(12): 1061-1064.
- Stolte M, Bayerdörffer E, Morgner A, Alpen B, Wündisch T, Thiede C, Neubauer A. 2002. *Helicobacter* and gastric MALT lymphoma. *Gut* 50(3): iii19-iii24.
- Strachan DP. 1989. Hay fever, hygiene, and household size. *BMJ* 299 (6710): 1259-1260.
- Strachan DP. 2000. Family size, infection and atopy: the first decade of the "hygiene hypothesis". *Thorax* 55 (1): S2-S10.
- Strang RR. 1965. The association of gastroduodenal ulceration with Parkinson's disease. *Med J Austr* 52: 842-843.
- Suerbaum S, Josenhans C. 2007. *Helicobacter pylori* evolution and phenotypic diversification in a changing host. *Nat Rev Microbiol.* 5(6): 441-452.
- Sunnerstam B, Kjerstad T, Jansson L, Giesecke J, Bergström M, Ejderhamn J. 1999. Detection of *Helicobacter pylori* in a pediatric population: comparison of three commercially available serological tests and one in-house enzyme immunoassay. *J Clin Microbiol.* 37(10): 3328-3331.
- Suzuki N, Wakasugi M, Nakaya S, Okada K, Mochida R, Sato M, Kajiya H, Takahashi R, Hirata H, Ezure Y, Koga Y, Fukuda Y, Shimoyama T. 2002. Production and application of new monoclonal antibodies specific for a fecal *Helicobacter pylori* antigen. *Clin. Diagn. Lab Immunol* 9(1): 75-78.
- Suzuki T, Matsushima M, Masui A, Watanabe K, Takagi A, Ogawa Y, Shirai T, Mine T. 2005. Effect of *Helicobacter pylori* eradication in patients with chronic idiopathic thrombocytopenic purpura-a randomized controlled trial. *Am J Gastroenterol* 100(6): 1265-1270.
- Suzuki H, Marshall BJ, Hibi T. 2006. Overview: *Helicobacter pylori* and extragastric disease. *Int J of Hematology* 84(4): 291-300.
- Svec A, Kordas P, Pavlis Z, Novotný J. 2000. High prevalence of *Helicobacter heilmannii*-associated gastritis in a small, predominantly rural area: further evidence in support of a zoonosis? *Scand. J. Gastroenterol* 35(9): 925-928.

- Sykora J, Hejda V, Varvarovská J, Stozicky F, Gottrand F, Siala K. 2003. *Helicobacter heilmannii* related gastric ulcer in childhood. J Pediatr Gastroenterol Nutr 36(3): 410-413.
- Szabo S. 1979. Dopamine disorder in duodenal ulceration. Lancet ii: 880-882.
- Szabo I, Brutsche S, Tombola F, Moschioni M, Satin B, Telford JL, Rappuoli R, Montecucco C, Papini E, Zoratti M. 1999. Formation of anion-selective channels in the cell plasma membrane by the toxin VacA of *Helicobacter pylori* is required for his biological activity. EMBO J 18(20): 5517-5527.
- Szeredi L, Palkovics G, Solymosi N, Tekes L, Méhesfalvi J. 2005. Study on the role of gastric *Helicobacter* infection in gross pathological and histological lesions of the stomach of finishing pigs. Acta Vet. Hung. 53(3): 371-383.
- Takahashi T, Yujiri T, Shinohara K, Inoue Y, Sato Y, Fujii Y, Okubo M, Zaitzu Y, Ariyoshi K, Nakamura Y, Nawata Y, Oka Y, Shirai M, Tanizawa Y. 2004. Molecular mimicry by *Helicobacter pylori* CagA protein may be involved in the pathogenesis of *H. pylori*-associated chronic idiopathic thrombocytopenic purpura. Br J Hematol 124(1): 91-96.
- Takashima T, Adachi K, Kawamura A, Yuki M, Fujishiro H, Rumi MA, Ishihara S, Watanebe M, Kinoshita Y. 2002. Cardiovascular risk factors in subjects with *Helicobacter pylori* infection. Helicobacter 7(2): 86-90.
- Takeda K, Ankira S. 2004. TLR signaling pathways. Semin Immunol 16(1): 3-9.
- Tamer GS, Tengiz I, Ercan E, Duman C, Alioglu E, Turk UO. 2009. *Helicobacter pylori* seropositivity in patients with acute coronary syndromes. Dig Dis Sci 54(6): 1253-1256.
- Tan AH, Mahadeva S, Marras C, Thalha AM, Kiew CK, Yeat CM, Ng SW, Ang SP, Chow SK, Loke MF, Vadivelu JS, Ibrahim N, Yong HS, Tan CT, Fox SH, Lang AE, Lim SY. 2015. *Helicobacter pylori* infection is associated with worse severity of Parkinson's disease. Parkinsonism and related disorders 21(3): 221-225.
- Tanaka E, Singh G, Saito A, Syouji A, Yamada T, Urano W, Nakajima A, Taniguchi A, Tomatsu T, Hara M, Saito T, Kamatani N, Yamanaka H. 2005. Prevalence of *Helicobacter pylori* infection and risk of upper gastrointestinal ulcer in patients with rheumatoid arthritis in Japan. Mod Rheumatol 15(5): 340-345.
- Tansey MG, McCoy MK, Frank-Cannon TC. 2007. Neuroinflammatory mechanisms in Parkinson's disease: Potential environmental triggers, pathways, and targets for early therapeutic intervention. Exp Neurol 208 (1): 1-25.
- Teoman I, Ozmeric N, Ozcan G, Alaaddinoglu E, Dumlu S, Akyon Y, Balos K. 2007. Comparison of different methods to detect *Helicobacter pylori* in the dental plaque of dyspeptic patients. Clin Oral. Invest. 11(3): 201-205.
- Terry K, Williams SM, Connolly L, Ottemann KM. 2005. Chemotaxis plays multiple roles during *Helicobacter pylori* animal infection. Infect Immun 73(2): 803-811.
- Testerman TL and Morris J. 2014. Beyond the stomach: An updated view of *Helicobacter pylori* pathogenesis, diagnosis and treatment. World J of Gastroenterol 20(36): 12781-12808.
- Thia KT, Loftus EV, Sandborn WJ, Yang SK. 2008. An update on the epidemiology of inflammatory bowel disease in Asia. Am J Gastroenterol. 103(12): 3167-3182.
- Tian XY, Zhu H, Zhao J, She Q, Zhang GX. 2012. Diagnostic performance of urea breath test, rapid urease test and histology for *Helicobacter pylori* infection in patients with partial gastrectomy: a meta-analyses. J Clin Gastroenterol 46(4): 285-292.
- Tomb JF, White O, Kerlavage AR, Clayton RA, Sutton GG, Fleischmann RD, Ketchum KA, Klenk HP, Gill S, Dougherty BA, Nelson K, Quackenbush J, Zhou L, Kirkness EF, Peterson S, Loftus B, Richardson D, Dodson R, Khalak HG, Glodek A, McKenney K, Fitzgerald LM, Lee N, Adams MD, Hickey EK, Berg DE, Gocayne JD, Utterback TR, Peterson JD, Kelley JM, Cotton MD, Weidman JM, Fujii C, Bowman C, Watthey L, Wallin E, Hayes WS, Borodovsky M, Karp PD, Smith HO, Fraser CM, Venter JC. 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. Nature 388(6642): 539-547.
- Trebesius K, Adler K, Vieth M, Stolte M, Haas R. 2001. Specific detection and prevalence of *Helicobacter heilmannii*-like organisms in the human gastric mucosa by fluorescent in situ hybridization and partial 16S ribosomal DNA sequencing. J. Clin. Microbiol 39(4): 1510-1516.
- Tseng CA, Wang WM, Wu DC. 2005. Comparison of the clinical feasibility of three rapid urease tests in the diagnosis of *Helicobacter pylori* infection. Dig Dis Sci 50(3): 449-452.
- Tsolaki F, Kountouras J, Topouzis F, Tsolaki M. 2015. *Helicobacter pylori* infection, dementia and primary open-angle glaucoma: are they connected? BMC Ophthalmology 15: 24.

- Tunca A, Türkay C, Tekin O, Kargili A, Erbayrak M. 2004. Is *Helicobacter pylori* infection a risk factor for migraine? A case control study. *Acta Neurol Belg* 104(4): 161-164.
- Tursi A. 2006. Onset of Crohn's disease after *Helicobacter pylori* eradication. *Inflamm Bowel Dis* 12(10): 1008-1009.
- Twelves D, Perkins KS, Counsell C. 2003. Systematic review of incidence studies of Parkinson's disease. *Mov Disord* 18(1): 19-31.
- Vafaeimanesh J, Hejazi SF, Damanpak V, Vahedian M, Sattari M, Seyyedmajidi M. 2014. Association of *Helicobacter pylori* infection with coronary artery disease: Is *Helicobacter pylori* a risk factor? *Scientific World Journal* 2014: 516354.
- Vale FF, Encarnação P, Vítor JM. 2008. A new algorithm for cluster analysis of genomic methylation: the *Helicobacter pylori* case. *Bioinformatics* 24(3): 383-388.
- Vale FF, Mégraud F, Vítor JM. 2009. Geographic distribution of methyltransferases of *Helicobacter pylori*: evidence of human host population, isolation and migration. *BMC Microbiol* 9: 193.
- Vandamme P, Peeters C. 2014. Time to revisit polyphasic taxonomy. *Antonie Van Leeuwenhoek* 106(1): 57-65.
- Vandenbroucke RE, Dejonckheere E, Van Lint P, Demeestere D, Van Wonterghem E, Vanlaere I, Puimège L, Van Hauwermeiren F, De Rycke R, McGuire C, Campestre C, López-Otin C, Matthys P, Leclercq G, Libert C. 2012. Matrix metalloprotease 8-dependent extracellular matrix cleavage at the blood-CSF barrier contributes to lethality during systemic inflammatory diseases. *Journal of Neuroscience* 32(29): 9805-9816.
- Van den Bulck K, Decostere A, Baele M, Driessen A, Debongnie JC, Burette A, Stolte M, Ducatelle R, Haesebrouck F. 2005. Identification of non-*Helicobacter pylori* spiral organisms in gastric samples from humans, dogs and cats. *J. Clin. Microbiol.* 42(5): 2256-2260.
- Van den Bulck K, Decostere A, Gruntar I, Baele M, Krt B, Ducatelle R, Haesebrouck F. 2005. *In vitro* antimicrobial susceptibility testing of *Helicobacter felis*, *Helicobacter bizzozeronii* and *Helicobacter salomonis*. *Antimicrob. Agents. Chemother.* 49(7): 2297-3000.
- Van den Bulck K, Decostere A, Baele M, Vandamme P, Mast J, Ducatelle R, Haesebrouck F. 2006. *Helicobacter cynogastricus* sp. nov., a *Helicobacter* species isolated from the canine gastric mucosa. *Int j Syst Evol Microbiol.* 56(7): 1559-1564.
- Van Loon S, Bart A, den Hertog EJ, Nikkels PG, Houwen RH, De Schryver JE, Oudshoorn JH. 2003. *Helicobacter heilmannii* gastritis caused by cat to child transmission. *J. Pediatr. Gastroenterol. Nutr.* 36(3): 407-409.
- Vawter MP, Dillon-Carter O, Tourtellotte WW, Carvey P, Freed WJ. 1996. TGFbeta1 and TGFbeta2 concentrations are elevated in Parkinson's disease in ventricular cerebrospinal fluid. *Exp Neurol* 142(2): 313-322.
- Veijola L, Myllyluoma E, Korpela R, Rautelin H. 2005. Stool antigen tests in the diagnosis of *Helicobacter pylori* infection before and after eradication therapy. *World J Gastroenterol* 11(46): 7340-7344.
- Velayos B, Fernández-Salazar L, Pons-Renedo F, Muñoz MF, Almaraz A, Aller R, Ruíz L, Del Olmo L, Gisbert JP, González-Hernández JM. 2012. Accuracy of the urea breath test performed immediately after emergency endoscopy in peptic ulcer bleeding. *Dig. Dis. Sci* 57(7): 1880-1886.
- Vercellino M, Votta B, Condello C, Piacentino C, Romagnolo A, Merola A, Capello E, Mancardi GL, Mutani R, Giordana MT, Cavalla P. 2008. Involvement of the choroid plexus in multiple sclerosis autoimmune inflammation: a neuropathological study. *Journal of neuroimmunology* 199(1-2): 133-141.
- Verhoef C, Pot RG, de Man RA, Zondervan PE, Kuipers EJ, IJzermans JN, Kusters JG. 2003. Detection of identical *Helicobacter* DNA in the stomach and in the non-cirrhotic liver of patients with hepatocellular carcinoma. *Eur J Gastroenterol Hepatol* 15(11): 1171-1174.
- Vermoote M, Vandekerckhove TT, Flahou B, Pasmans F, Smet A, De Groote D, Van Crieke W, Ducatelle R, Haesebrouck F. 2011. Genome sequence of *Helicobacter suis* supports its role in gastric pathology. *Vet Res* 42: 51.
- Vermoote M, Pasmans F, Flahou B, Van Deun K, Ducatelle R, Haesebrouck F. 2011: Antimicrobial susceptibility pattern of *Helicobacter suis* strains. *Vet. Microbiol* 153(3-4): 339-342.
- Vermoote M, Van Steendam K, Flahou B, Smet A, Pasmans F, Glibert P, Ducatelle R, Deforce D, Haesebrouck F. 2012. Immunization with the immunodominant *Helicobacter suis* urease subunit B induces partial protection against *H. suis* infection in a mouse model. *Vet Res* 43: 72.
- Vermoote M, Flahou B, Pasmans F, Ducatelle R, Haesebrouck F. 2013. Protective efficacy of vaccines based on the *Helicobacter suis* urease subunit B and γ -glutamyl transpeptidase. *Vaccine* 31(32): 3250-3256.

- Viala J, Chaput C, Boneca IG, Cardona A, Girardin SE, Moran AP, Athman R, Mémet S, Huerre MR, Coyle AJ, DiStefano PS, Sansonetti PJ, Labigne A, Bertin J, Philpott DJ, Ferrero RL. 2004. Nod1 responds to peptidoglycan delivered by *Helicobacter pylori* *cag* pathogenicity island. *Nat Immunol* 5(11): 1166-1174.
- Vitoriano I, Rocha-Gonçalves A, Carvalho T, Oleastro M, Calado CR, Roxo-Rosa M. 2011. Antigenic diversity among Portuguese clinical isolates of *Helicobacter pylori*. *Helicobacter* 16(2): 153-168.
- Vleminckx K, Kemler R. 1999. Cadherins and tissue formation: integrating adhesion and signaling. *BioEssays* 21(3): 211-220.
- Waluga M, Kukla M, Zorniak M, Bacik A, Kotulski R. 2015. From the stomach to other organs: *Helicobacter pylori* and the liver. *World Journal of hepatology* 7(18): 2136-2146.
- Wang M, Munch IC, Hasler PW, Prünke C, Larsen M. 2008. Central serous chorioretinopathy. *Acta Ophthalmol* 86: 126-145.
- Wang XI, Zhang S, Abreo F, Thomas J. 2010. The role of routine immunohistochemistry for *Helicobacter pylori* in gastric biopsy. *Am Diagn Pathol* 14(4): 256-259.
- Wang ZW, Li Y, Huang LY, Guan QK, Xu da W, Zhou WK, Zhang XZ. 2012. *Helicobacter pylori* infection contributes to high risk of ischemic stroke: evidence from a meta-analysis. *J Neurol* 259(12): 2527-2537.
- Wang Q, Yu C, Sun Y. 2013. The association between asthma and *Helicobacter pylori*: a meta-analysis. *Helicobacter* 18(3): 41-53.
- Wardi J, Shalev T, Shevah O, Boaz M, Avni Y, Shirin H. 2012. A rapid continuous-real-time-13C-urea breath test for the detection of *Helicobacter pylori* in patients after partial gastrectomy. *J. Clin. Gastroenterol* 46(4): 293-296.
- Warren JR and Marshall BJ. 1983. Unidentified curved bacilli on gastric epithelium in active chronic gastritis. *Lancet* 321(8336): 1273-1275.
- Watanabe T, Asano N, Kitani A, Fuss IJ, Chiba T, Strober W. 2011. Activation of type 1 IFN signaling by NOD1 mediates mucosal host defense against *Helicobacter pylori* infection. *Gut Microbiol.* 2(1): 61-65.
- Watanabe M, Kato J, Inoue I, Yoshimura N, Yoshida T, Mukoubayashi C, Deguchi H, Enomoto S, Ueda K, Maekita T, Iguchi M, Tamai H, Utsunomiya H, Yamamichi N, Fujishiro M, Iwane M, Tekeshita T, Mohara O, Ushijima T, Ichinose M. 2012. Development of gastric cancer in nonatrophic stomach with highly active inflammation identified by serum levels of pepsinogen and *Helicobacter pylori* antibody together with endoscopic rugal hyperplastic gastritis. *Int J Cancer* 131(11): 2632-2642.
- Weeks DL, Eskandari S, Scott DR, Sachs G. 2000. A H⁺-gated urea channel: the link between *Helicobacter pylori* urease and gastric colonization. *Science* 287(5452): 482-485.
- Wegmann W, Aschwanden M, Schaub N, Aenishänslin W, Gyr K. 1991. Gastritis associated with *Gastrospirillum hominis*--a zoonosis? *Schweiz Med Wochenschr.* 121(8): 245-254.
- Weller C, Charlett A, Oxlade NL, Dobbs SM, Dobbs RJ, Peterson DW, Bjarnason IT. 2005. Role of chronic infection and inflammation in the gastrointestinal tract in the etiology and pathogenesis of idiopathic parkinsonism. Part 3: predicted probability and gradients of severity of idiopathic parkinsonism based on *H. pylori* antibody profile. *Helicobacter* 10(4): 288-297.
- Wender M. 2003. Prevalence of *Helicobacter pylori* infection among patients with multiple sclerosis. *Neurol Neurochir Pol* 37(1): 45-48.
- Willhite DC, Blanke SR. 2004. *Helicobacter pylori* vacuolating cytotoxin enters cells, localizes in the mitochondria and induces mitochondrial membrane permeability changes correlated to toxin channel activity. *Cell Microbiol* 6(2): 143-154.
- Wilson KT, Crabtree JE. 2007. Immunology of *Helicobacter pylori*: insights into the failure of the immune response and perspectives on vaccine studies. *Gastroenterology.* 133(1): 288-308.
- Winge K, Skau AM, Stimpel H, Nielsen KK, Werdelin L. 2006. Prevalence of bladder dysfunction in Parkinson's disease. *Neurol Urodyn* 25(2): 116-122.
- Wirdefeldt K, Adami HO, Cole P, Trichopoulos D, Mandel J. 2011. Epidemiology and etiology of Parkinson's disease: a review of the evidence. *Eur J Epidemiol* 26(1): S1-S58.
- Wolburg H, Neuhaus J, Kniesel U, Krauss B, Schmid EM, Ocalan M, Farrell C, Risau W. 1994. Modulation of tight junction structure in blood-brain barrier endothelial cells. Effects of tissue culture, second messengers and cocultured astrocytes. *Journal of Cell Science* 107(5): 1347-1357.

- Wong F, Rayner-Hartley E, Byrne MF. 2014. Extraintestinal manifestations of *Helicobacter pylori*: a concise review. *World J of Gastroenterol* 20(34): 11950-11961.
- Wotherspoon AC, Ortiz-Hidalgo C, Falzon MR, Isaacson PG. 1991. *Helicobacter pylori*-associated gastritis and primary B-cell gastric lymphoma. *Lancet* 338(8776): 1175-1176.
- Wüppenhorst N, von Loewenich F, Hobmaier B, Vetter-Knoll M, Mohadjer S, Kist M. 2013. Culture of a gastric non-*Helicobacter pylori* *Helicobacter* from the stomach of a 14 year-old girl. *Helicobacter* 18(1): 1-5.
- Yiannopoulou KG, Efthymiou A, Karydakis K, Arhimandritis A, Bovaretos N, Tzivras M. 2007. *Helicobacter pylori* infection as an environmental risk factor for migraine without aura. *J Headache pain* 8(6): 329-333.
- Yakoob J, Abbas Z, Khan R, Naz S, Ahmad Z, Islam M, Awan S, Jafri F, Jafri W. 2012. Prevalence of non-*Helicobacter pylori* species in patients presenting with dyspepsia. *BMC Gastroenterol* 12: 3.
- Yamamoto K, Tanaka H, Nishitani Y, Nishiumi S, Miki I, Takenaka M, Nobutani K, Mimura T, Ben Suleiman Y, Mizuno S, Kawai M, Uchiyama I, Yoshida M, Azuma T. 2011. *Helicobacter suis* KB1 derived from pig gastric lymphoid follicles induces the formation of gastric lymphoid follicles in mice through the activation of B cells and CD4 positive cells. *Microbes and infection* 13(7): 697-708.
- Yamanishi S, Iizumi T, Watanabe E, Shimizu M, Kamiya S, Nagata K, Kumagai Y, Fukunaga Y, Takahashi H. 2006. Implications for induction of autoimmunity via activation of B-1 cells by *Helicobacter pylori* urease. *Infect Immun* 74(1): 248-256.
- Yamazaki S, Yamakawa A, Ito Y, Ohtani M, Higashi H, Hatekeyama M, Azuma T. 2003. The CagA protein of *Helicobacter pylori* is translocated into epithelial cells and binds to SHP-2 in human gastric mucosa. *J Infect Dis* 187(2): 334-337.
- Yan J, Yamaguchi T, Odaka T, Suzuki T, Ohyama N, Hara T, Sudo K, Nakamura K, Denda T, Takiguchi N, Yokosuka O, Nomura F. 2010. Stool antigen test is a reliable method to detect *Helicobacter pylori* in the gastric remnant after distal gastrectomy for gastric cancer. *J. Clin Gastroenterol* 44(1): 73-74.
- Yang H, Li X, Xu Z, Zhou D. 1995. "*Helicobacter heilmannii*" infection in a patient with gastric cancer. *Dig Dis Sci* 40(5): 1013-1014.
- Yang H, Goliger JA, Song M, Zhou D. 1998. High prevalence of *Helicobacter heilmannii* infection in China. *Dig Dis Sci* 43(7): 1493.
- Yang X, Gao Y, Zhao X, Tang Y, Su Y. 2011. Chronic *Helicobacter pylori* infection and ischemic stroke subtypes. *Neurol Res* 33(5): 467-472.
- Yip R, Limburg PJ, Ahlquist DA, Carpenter HA, O'Neill A, Kruse D, Stitham S, Gold BD, Gunter EW, Looker AC, Parkinson AJ, Nobmann ED, Petersen KM, Ellefson M, Schwartz S. 1997. Pervasive occult gastrointestinal bleeding in an Alaska native population with prevalent iron deficiency. Role of *Helicobacter pylori* gastritis. *JAMA* 277(14): 1135-1139.
- Yoshimura M, Isomoto H, Shikuwa S, Osabe M, Matsunaga K, Omagari K, Mizuta Y, Murase K, Murata I, Kohno S. 2002. A case of acute gastric mucosal lesions associated with *Helicobacter heilmannii* infection. *Helicobacter* 7(5): 322-326.
- Zavos C, Kountouras J, Sakkias G, Venizelos I, Deretzi G, Arapoglou S. 2012. Histological presence of *Helicobacter pylori* bacteria in the trabeculum and iris of patients with primary open-angle glaucoma. *Ophthalmic Res* 47(3): 150-156.
- Zhang S, Guo Y, Ma Y, Teng Y. 2008. Relationship between the cytotoxin-associated gen-A status of *H. pylori* strains and cerebral infarction in European Causasians and Chinese Han: a meta-analysis. *World J Gastroenterol* 14(8): 1286-1292.
- Zhang G, Ducatelle R, Pasmans F, D'Herde K, Huang L, Smet A, Haesebrouck F, Flahou B. 2013. Effects of *Helicobacter suis* γ -Glutamyl Transpeptidase on Lymphocytes: Modulation by Glutamine and Glutathione Supplementation and Outer Membrane Vesicles as a Putative Delivery Route of the Enzyme. *PLoS One*. 8(10): e77966.
- Zhang G, Ducatelle R, De Bruyne E, Joosten M, Bosschem I, Smet A, Haesebrouck F, Flahou B. 2015. Role of γ -glutamylpeptidase in the pathogenesis of *Helicobacter suis* and *Helicobacter pylori* infections. *Vet Research* 46: 31.
- Zhang YZ, Ran LY, Li CY, Chen XL. 2015. Diversity, Structures and Collagen-Degrading Mechanisms of 1 Bacterial Collagenolytic Proteases: an Overview. *Appl. Environ. Microbiol.* 81(18): 6098-6107.
- Zhou X, Zhang C, Wu J, Zhang G. 2013. Association between *Helicobacter pylori* infection and diabetes mellitus: a mate-analysis of observational studies. *Diabetes Res Clin Pract* 99(2): 200-208.
- Zhou X, Wu J, Zhang G. 2013. Association between *Helicobacter pylori* and asthma: a meta-analysis. *Eur J Gastroenterol Hepatol* 25(4): 460-468.

Zuberbier T, Aberer W, Asero R, Bindslev-Jensen C, Brzoza Z, Canonica GW, Church MK, Ensina LF, Giménez-Arnau A, Godse K, Gonçalo M, Grattan C, Hebert J, Hide M, Kaplan A, Kapp A, Abdul Latiff AH, Mathelier-Fusade P, Metz M, Nast A, Saini SS, Sánchez-Borges M, Schmid-Grendelmeier P, Simons FE, Staubach P, Sussman G, Toubi E, Vena GA, Wedi B, Zhu XJ, Maurer M. 2009. EAACI/GA(2)LEN/EDF/WAO guideline: definition, classification and diagnosis of urticaria. *Allergy* 64(10): 1417-1426.

SCIENTIFIC AIMS

Helicobacter pylori (*H. pylori*) is considered to be the most important pathogen involved in gastric disease in humans. Besides causing inflammation and lesions in the stomach, infection with this agent has also been linked to extragastric manifestations, including brain disorders (Holtmann and Talley, 2014). Gastric pathologies in humans have, however, also been associated with spiral-shaped, gastric *Helicobacter* species different from *H. pylori* (non-*H. pylori* helicobacters (NHPH)). *H. suis*, an organism with a corkscrew-like morphology that naturally colonizes the stomach of pigs and non-human primates, is considered to be the most prevalent NHPH in humans. The role of gastric bacteria, other than *H. pylori*, in brain functionality is largely unknown. There is, however, a case report suggesting that infections with *H. suis* might also be associated with Parkinson's disease (PD) (Dobbs et al., 2005).

Therefore, the **general aim** of this thesis was to evaluate if and how a gastric infection with *H. suis* might be linked to PD in human patients.

Since it became clear during these studies that the current diagnostic tests used for detection of *H. suis* infections suffered from severe shortcomings, an **additional aim** of the thesis was to improve diagnostic assays.

It has been reported that an *H. pylori* infection might be associated with PD. Overall in idiopathic parkinsonism (IP), successful *H. pylori* eradication was shown to improve brady/hypokinesia. A poorer response was seen in the anti-nuclear antibody seropositive, who may have had residual *Helicobacter*, undetected by culture and PCR. Overt failure was accompanied by marked deterioration. Apart from *H. pylori* infections, IP has also been associated with rural-living and farm-experience, especially live-stock farm-experience. To determine whether there might be an association between PD and *H. suis* infections in humans, the **first specific aim** of this thesis was to compare the prevalence of *H. suis* in gastric biopsies between patients suffering from IP and controls.

In our first study we showed that the frequency of *H. suis* was significantly higher in IP patients compared to control patients. However, it remained unclear if an infection with this gastric microorganism might play a role in aetiopathogenesis. The brain is protected from alterations in the systemic circulation by several specialized neuro-epithelial barriers, such as the blood-brain barrier and the blood-cerebrospinal fluid (CSF) barrier, that determine what will (and what will not) enter the brain. Dysfunction of these brain barriers has been implicated in the pathogenesis of PD. It has been suggested that systemic inflammation could play an important role in the onset of neurodegenerative disorders: it might be accompanied by barrier disruption associated with immune/inflammatory mediators. Since gastric *Helicobacter* infections have been proposed to be associated with systemic inflammation, the **second specific aim** of this thesis was to evaluate whether and how a *H. suis* infection could have an impact on the brain homeostasis.

In order to determine the importance of an *H. suis* infection in gastric and extragastric disorders in human patients, accurate diagnostic methods for the specific detection of *H. suis* in gastric biopsies are required. Currently, the diagnosis of infections with this spiral-shaped bacterium is hampered by its fastidious nature, the low colonization density and the patchy distribution of *H. suis* in the human stomach. Additionally, when using

the conventional *ureA*-based SYBR green RT-PCR for the detection of *H. suis* infections, non-specific interactions with host DNA and DNA from other *Helicobacter* spp. may occur, as was noticed during our first study. Thus, **the third specific aim** of this thesis was to optimize the currently used methods for demonstration of *H. suis* in the stomach. We first developed and validated a new, probe-based RT-PCR that allowed detection and quantification of *H. suis* in gastric tissue samples. Secondly, a mouse-passage-based protocol was optimized for the isolation of low numbers of *H. suis* bacteria.

EXPERIMENTAL STUDIES

Chapter 1

Significantly higher frequency of *Helicobacter suis* in patients with idiopathic parkinsonism than in control patients

C. Blaecher^{*,†}, A. Smet^{*}, B. Flahou^{*}, F. Pasmans^{*}, R. Ducatelle^{*}, D. Taylor^{†,‡}, C. Weller[†], I. Bjarnason[§], A. Charlett[¶], A. J. Lawson^{**}, R. J. Dobbs^{†,‡,§}, S. M. Dobbs^{†,‡,§} & F. Haesebrouck^{*}

^{*}Faculty of Veterinary Medicine, Ghent University, Merelbeke, Belgium.

[†]Institute of Pharmaceutical Science, King's College London, London, UK.

[‡]The Maudsley Hospital, London, UK.

[§]Department of Gastroenterology, King's College Hospital, London, UK.

[¶]Statistics Modelling and Economics Department of Centre for Infectious Disease Surveillance and Control, Public Health England, London, UK.

^{**}Gastrointestinal Bacteriology Reference Unit, Public Health England, London, UK.

Adapted from: Alimentary Pharmacology & Therapeutics 2013, 38 (11): 1347-1353

Abstract

Background: There is increased proportional mortality from Parkinson's disease (PD) amongst livestock farmers. The hypokinesia of Parkinson's disease has been linked to *Helicobacter pylori* (*H. pylori*). *Helicobacter suis* (*H. suis*) is the most common zoonotic *Helicobacter* in man.

Aims: To compare the frequency of *H. suis*, relative to *H. pylori*, in gastric biopsies of patients with idiopathic parkinsonism (IP) and controls from gastroenterology services.

Methods: Coded DNA extracts, archived at a *Helicobacter* Reference Laboratory, from IP patient and gastroenterology service biopsies were examined for *H. suis*, using species-specific RT-PCR.

Results: Relative risk of having *H. suis* in 60 IP patients compared with 256 controls was 10 times greater than that of having *H. pylori*. In patients with IP and controls, respectively, frequencies of *H. suis* were 27 (exact binomial 95% C.I. 15, 38) and 2 (0, 3)%, and of *H. pylori*, 28 (17, 40) and 16 (12, 21)%. Excess of *H. suis* in IP held when only the antral or corporal biopsy was considered. Of 16 IP patients with *H. suis*, 11 were from 19 with proven *H. pylori* eradication, 3 from 17 pre-*H. pylori* eradication, 2 from 24 *H. pylori* culture/PCR negative. Frequency was different between groups ($P = 0.001$), greatest where *H. pylori* had been eradicated. Even without known exposure to anti-*H. pylori* therapy, *H. suis* was more frequent in IP patients (5/41) than in controls (1/ 155) ($P = 0.002$). Partial multilocus sequence typing confirmed that strains from IP patients (6) and control (1) differed from RT-PCR standard strain.

Conclusions: Greater frequency of *H. suis* in idiopathic parkinsonism appears exaggerated following *H. pylori* eradication. Multilocus sequence testing comparison with porcine strains may clarify whether transmission is from pigs/ porcine products or of human-adapted, *H. suis*-like, bacteria.

INTRODUCTION

The broad brush of epidemiology has provided enigmatic aetiopathogenic clues in Parkinson's disease (PD). It has been linked, albeit inconsistently, with rural living and farm experience (Wirdefeldt et al., 2011). Such associations have long been explained away by putative exposure to agrochemicals. However, a mortality study in 26 US states found increased proportional mortality from PD among livestock, but not arable, farmers (Lee et al., 2002). From a total of 6 million death certificates, 267479 decedents were classified as either crop or livestock farmers. Amongst white males, the proportional mortality from PD was significantly higher in livestock farmers than in non-farmers, but lower in crop farmers. Zoonotic infections were considered as an explanation, but it was noted that the pesticides used in animal infestations are different from those used on crops. These effects were not replicated in the much smaller numbers of White female or of African American decedents.

The 1965 finding of prodromal peptic ulcers in PD (Strang, 1965) paved the way to exploring any link between (the then undiscovered) *Helicobacter pylori* (*H. pylori*) and PD (Dobbs et al., 2008; 2012; 2010; 2013). The 2012 Maastricht IV Consensus Report (Malfertheiner et al., 2012) acknowledges interesting associations. In a randomised placebo-controlled trial, we found that biopsy-proven *H. pylori* eradication had differential effects on objective measures of PD facets: improvement in hypokinesia and worsening rigidity over the year post-eradication, both plateauing over the next year (Dobbs et al., 2010). Overall, there was clinically relevant improvement. This was independent of any (stable, long t½) anti-parkinsonian medication. Receipt of levodopa was an exclusion. Moreover, the effect on hypokinesia was indication-specific (Dobbs et al., 2013). A longitudinal observational study confirmed improvement in hypokinesia following *H. pylori* eradication. Anti-microbials for other indications had no such effect, but successive courses were associated with cumulative increase in rigidity.

Could, then, zoonotic-transmission of gastric non-*H. pylori* *Helicobacter* species (NHPH) contribute to PD? The term NHPH represents a group of closely related, but distinct, bacterial species found in different animal species, such as *H. suis* in pigs, and *H. felis*, *H. salomonis*, *H. bizzozeronii*, *H. heilmannii* sensu stricto, *H. cynogastricus* and *H. baculiformis* in cats and dogs (Haesebrouck et al., 2009). Although infection with these bacteria has been associated with human gastric disease in a substantial number of patients (Debongnie et al., 1995; Morgner et al., 1995; Stolte et al., 1997; Trebesius et al., 2001; O'Rourke et al., 2004; Van den Bulck et al., 2005; Joosten et al., 2013), human NHPH strains have been isolated on only three occasions (Andersen et al., 1996; Kivisto et al., 2010; Wüppenhorst et al., 2013). Subsequent identification to species level has revealed these to be *H. bizzozeronii* in two cases (Kivisto et al., 2010; Javala et al., 2001) and *H. felis* in the other (Wüppenhorst et al., 2013). Based on histopathology of gastric biopsy, the prevalence of NHPH in patients with gastric disease has been estimated at between 0.2% and 6%, depending on geographical distribution (Stolte et al., 1997). Several studies have demonstrated that *H. suis* is the most frequent NHPH species in humans (Haesebrouck et al., 2009; Trebesius et al., 2001; Van den Bulck et al., 2005). However, its *in vitro* culture, starting from human gastric mucosa, has been unsuccessful so far, underlining the extreme fastidiousness of gastric NHPH in general and *H. suis* in particular.

We report the frequency of *H. suis* DNA in archived DNA extracts from gastric biopsies in a group of PD patients and a group of gastroenterology patients, as part of a service evaluation. The relative prevalence of *H. pylori*

immunoblot seropositivity in people with and without PD (37% and 31% respectively) has been reported (Charlett et al., 2009). Here, we standardise the frequency of *H. suis* positivity against that recorded for *H. pylori* (culture, or if negative, PCR), in order to compare the two sets of archived DNA extracts.

METHODS

Service evaluation

The term '*H. heilmannii*' was commonly used to describe spiral gastric helicobacters seen on histopathology. The diagnostic service of the Gastrointestinal Reference Unit, Public Health England, for PCR detection of '*H. heilmannii*-like organisms' in gastric biopsy material (Chisholm et al., 2003) was re-evaluated, using coded archived DNA extracts. As *H. suis* is the commonest reported NHPH in humans and a species-specific assay is now available (O'Rourke et al., 2004), it is a good starting point for evaluating whether NHPH cases are being missed. Interestingly, the original '*H. heilmannii*' assay was set up in response to a case of a spiral *Helicobacter* in PD (Dobbs et al., 2005).

Sourcing coded DNA extracts

The *Helicobacter* Reference Laboratory in the Gastrointestinal Bacteriology Reference Unit, Public Health England, held DNA from gastric biopsies, extracted (at time of receipt) over the last decade, stored at -80 °C. Biopsies had been received with request for *Helicobacter* culture (isolation/identification/anti-microbial susceptibility) and, if biopsy culture-negative, detection of *H. pylori*-specific DNA. Culture-negative biopsies had been tested using a PCR targeting *16S rRNA* (primer pair HP1/HP2) (Ho et al., 1991) and *vacA* (Vac3624F/Vac3853R) (Chisholm et al., 2001) genes. To detect '*Helicobacter heilmannii*-like organisms' a *16S rRNA* assay had been used (Chisholm et al., 2003). For the service evaluation, archived DNA extracts, identified only by their accession number, were couriered to the Laboratory of Veterinary Bacteriology and Mycology, Ghent University. All extracts archived over this decade (paired antral and corporal biopsies) from 60 patients with 'clinically definite' idiopathic parkinsonism (IP) (Calne et al., 1992), diagnosed at a National clinic, were retrieved. Clinically definite IP refers to any combination of three of the four cardinal features: resting tremor, rigidity, brady/hypokinesia, impaired postural reflexes. Alternatively sufficient is having two of the features, with one of first three asymmetrical (Responsiveness to a dopaminergic drug challenge was not a requirement). Details of anti-microbials used in any previous anti-*H. pylori* treatment (19/60) had been recorded on the request form.

DNA extracts from biopsies originating from English gastroenterological services, selected for attention to documenting any exposure to anti-*Helicobacter* therapy on the request form, were used as 'controls'. This criterion yielded extracts from 256 patients, archived over a similar period (single biopsy on given occasion except in four, where two sites sampled). There had been previous exposure to anti-*Helicobacter* therapy in 101, no recorded exposure in the rest. No request form mentioned Parkinson's disease, or any other issues outside the context of diagnostic endoscopy. Crude prevalence rate estimates for PD in 34 European studies range widely, from 65.6 to 12500 per 100000 (von Campenhausen et al., 2005): that is, 0–32 diagnosed cases would be expected in the gastroenterological services' patient group, even if they had no particular predisposition.

Frequencies of *Helicobacter* species reported here are not construed as prevalence: there is selection in the controls (for adequate documentation by above criterion) and of IP patients (for a clinic with particular interest in the gastroenterological aspects of PD). Any direct estimate of association between presence of *H. suis* and PD could be biased by cohort selection. We therefore obtain an estimate of the risk of *H. suis* in patients with IP compared with those from gastroenterological services by standardizing for the relative risk of having *H. pylori*, and by setting *H. suis* frequency in context of exposure to anti-*Helicobacter* therapy. No analysis of the relationship of *H. suis* status to gastric symptoms was planned: the request form did not contain a checklist.

***H. suis* detection**

The presence of *H. suis* DNA in the extracts was determined using a *H. suis*-specific quantitative real time (RT)-PCR, based on the *ureA* gene. For generating the standard, part of the *ureAB* gene cluster (1236 bp) from *H. suis* strain HS5 was amplified using primers U430F and U1735R, as described previously (O'Rourke et al., 2004). The standard consisted of 10-fold dilutions, starting at 10^8 PCR amplicons, for each 9 μ L of reaction mixture. One microliter of extracted DNA template was added to 9 μ L reaction mixture, consisting of 0.25 μ L of both primers located within the 1236 bp fragment (to yield a 150 bp PCR product), 3.5 μ L HPLC water and 5 μ L SensiMixTM SYBR No-ROX (Bioline Reagents Ltd, London, UK). Sense primer was BF_HsuisF1: 5'-AAA ACA MAG GCG ATC GCC CTG TA-3'. Anti-sense primer was BF_HsuisR1: 5'-TTT CTT CGC CAG GTT CAA AGC G-3'. Annealing temperature was 62 °C. Both standards and samples were run in duplicate on a CFX96TM RT-PCR System with a C1000 Thermal Cycler (Bio-Rad, Hercules, CA, USA). To confirm the presence of *H. suis* DNA, all positive samples were sequenced (Joosten et al., 2013). All 413 available extracts (including replicate biopsies at a given time and any follow-up biopsies) were assayed on two separate occasions to estimate between-assay agreement. To demonstrate that the *H. suis* DNA in the biopsy extracts was different from the *H. suis* strain HS5 DNA used as standard in the RT-PCR, multilocus sequence typing (MLST) was performed with seven housekeeping genes, as previously described (Liang et al., 2013).

Statistical analysis

Use of exact binomial confidence intervals provided an estimate of the uncertainty in estimates of proportions. The risk ratio, risk of having *H. suis* in IP patients to that in controls compared with relative risk of having *H. pylori*, was estimated from the paired data on the two species, using a conditional Poisson regression with robust standard errors (Cummings, 2011).

RESULTS

The distribution of age at time of first biopsy in controls [mean 52 (data interval 24, 81) years] encapsulated that in the IP patients [62 (45, 79) years]. There was no significant difference in the proportion of male patients: 57% of the IP patients were male (34/60), 47% (120/256) of the controls. Overall agreement (99.3%) between the two *H. suis* RT-PCR assay runs on separate occasions was strong ($n = 413$, Kappa = 0.95, $P < 0.001$: null hypothesis of no agreement rejected) and there was no significant difference between the first and second run {exact McNemar's test, ratio of paired proportions 1.11 [95% confidence interval (CI) 0.99, 1.24], $P = 0.25$ }. Thirty-one of the 413 extracts were positive in either run, 28 in both. Nucleotide sequencing of positive samples, and subsequent Basic Local Alignment Search Tool (BLAST) analysis, revealed 95–100% homology with

known *H. suis* strains. The frequency of detection of *H. suis* and *H. pylori* in the extract(s), held in the archive from the 60 IP patients and 256 controls, is shown in Table 1. In patients biopsied on more than one occasion, only the first occasion is considered. Overall, *H. suis* DNA was present in 27 (binomial exact 95% CI 15, 38)% of the IP patients and in 2 (0, 3)% of controls. It was present in 18 (10, 30)% of antral biopsies and 13 (6, 25)% of corporal biopsies from IP patients. Thus, the excess *H. suis* frequency was not an artefact due to sampling both regions in IP, but only one region (46% antral, 1% corporal, 2% duodenal, rest 'gastric') in all but four controls (2 had antral and corporal biopsies, 1 antral and duodenal, in 1 both were labelled 'gastric').

Table 1: Frequency of *H. suis* contrasted with *H. pylori* in gastric biopsy DNA extracts from IP patients and controls with and without known exposure to anti-*H. pylori* therapy.

Frequency	IP patients % (no. with species/total)	Controls % (no. with species/total)
<i>H. suis</i>		
Previous exposure to anti- <i>H. pylori</i> therapy	58 (11/19)*	3 (3/101)
No known exposure	12 (5/41)#	1 (1/155)
Total	27 (16/60)**	2 (4/256)**
<i>H. pylori</i>		
Previous exposure to anti- <i>H. pylori</i> therapy	0 (0/19)	22 (22/101)
No known exposure	41 (17/41)	13 (20/155)
Total	28 (17/60)	16 (42/256)

*: Anti-microbial prescription was guided by *in vitro* sensitivities:- Of the 11 with *H. suis*, 10 had received 1 week of amoxicillin, clarithromycin and proton pump inhibitor (PPI) (1 a further course of metronidazole, tetracycline, PPI and tripotassium dicitratobismuthate when he remained urea breath test-positive) and 1 had received 1 week of clarithromycin, tetracycline and PPI. Of 8 without *H. suis*, 7 had received 1 week of amoxicillin, clarithromycin and PPI and 1 had received 1 week of clarithromycin, tetracycline and PPI.

#: Three extracts also positive for *H. pylori*.

**: Mean age (gender) of the 16 IP patients with *H. suis* was 58 (range 47–68) years (10 male, 6 female), of the 4 controls, 54 (43–64) years (1 male, 3 female).

The relative risk of having *H. suis* in IP patients compared with controls was 9.9 times greater than that of having *H. pylori* [i.e. $[(16/60)/(4/256)]/[(17/60)/(42/256)] = 17.07/1.73$]. The 95% confidence interval (3.0, 32.7) did not include 1, indicating that the two relative risks are significantly different. This was despite a higher frequency of *H. pylori* in IP patients [28 (binomial exact 95% C.I. 17, 40)%] than in controls [16 (12, 21)%]. Of the 16 IP patients with *H. suis*, 11 were from the 19 with proven *H. pylori* eradication, 3 from the 17 pre-*H. pylori* eradication and 2 from the 24 who were *H. pylori*-negative. Frequency of *H. suis* detection differed significantly among these three groups (Fisher's exact test, $P = 0.001$), being greatest in those who had undergone anti-*H. pylori* therapy than in the remainder. There was no significant difference in the anti-*H. pylori* regimen

(clarithromycin/amoxicillin/proton pump inhibitor or other: footnote Table 1) between those with and without *H. suis* in its aftermath. Even in those without known previous exposure to anti-*H. pylori* therapy, *H. suis* was more frequent in IP patients (5/41) than in controls (1/155) ($P = 0.002$). The low frequency of *H. suis* in the 256 controls precluded estimation of any increased risk with exposure to anti-*H. pylori* therapy. Partial MLST results for the *H. suis* DNA found in extracts from 6 IP patients and 1 control are shown in Table 2. The allele pattern in these samples differed from that in the HS5 DNA standard: that is, the sample results were not due to contamination. It was not possible to obtain complete MLST sequence types, due to interference from human DNA and the low amount of *H. suis* DNA found in some extracts. None of the samples had been reported as positive for '*Helicobacter heilmannii*-like organisms' on the original assay 16S rDNA assay.

Table 2: Partial *H. suis* MLST results on DNA extracts from gastric biopsies of IP patients and a control

DNA extract	Allele no.						
	<i>atpA</i>	<i>efp</i>	<i>mutY</i>	<i>ppa</i>	<i>trpC</i>	<i>ureAB</i>	<i>yphC</i>
IP patient 1	4		11				
IP patient 2		1					
IP patient 3						1	
IP patient 4		1	11				
IP patient 5			5			1	1
IP patient 6		1	5				
Control patient		1	1			1	
HS5*	4	2	4	1	1	1	1

*: positive control in the quantitative RT-PCR assays.

DISCUSSION

Relative frequency of *H. suis* in IP

The frequency of finding *H. suis* DNA in extracts from gastric biopsies in a well-defined disease state, clinically definite IP, contrasts sharply with the apparently 'sporadic' nature in patients from gastroenterological services undergoing diagnostic endoscopy. The statistical significance of the finding withstood standardization for the relative frequency of *H. pylori* in the two patient groups. However, defining the strength of association between presence of *H. suis* DNA in gastric biopsies and PD will clearly require representative cohorts of IP probands and controls. As human infection with NHPH is characteristically sparse and patchy (Stolte et al., 1997), we may be underestimating the true frequency of *H. suis* in both patient groups. There are no previous studies of the frequency of NHPH in PD for comparison.

Proposed pathophysiological role of *Helicobacter* species in IP

The concept of 'brain-altering' remote infections is receiving increasing attention. Neuro-inflammation in IP may not merely be the adverse effect of microglial scavenging of degenerating neurons and reaction to aberrant protein (Tansey et al., 2007; Zipp et al., 2006). If PD is driven by systemic immuno-inflammatory processes (Dobbs et al., 2008; 2012), intervention against them, or their driving forces, could modify its course

(Dobbs et al., 2013; Malfertheiner et al., 2012). A U-turn in brady/hypokinesia and weight gain was described after eradicating a spiral *Helicobacter*, associated with antral gastritis, in a cachectic IP patient, who had been wheelchair-bound without assistance for over a year (Dobbs et al., 2005). The aetiopathogenic significance of zoonotic helicobacters in IP remains to be explored. Finding clinical correlates (e.g. epidemiological, such as mortality, and pathophysiological, such as circulating inflammatory markers) would suggest that *H. suis* DNA in gastric mucosa is of significance in IP. Autoimmunity is suggested as the mechanism of the effect of *H. pylori* eradication on hypokinesia (Dobbs et al., 2008; 2010; 2012). Response appears unrelated to infection load. Poor response is associated with anti-nuclear antibody seropositivity. There are HLA-DR risk loci for PD. If NHPH eradication has a similar effect, then the immune hypothesis could move to a pattern recognition response rather than classical HLA-restricted autoimmunity.

Optimizing diagnosis of human *H. suis* infection

Diagnosing NHPH in IP is problematic. Our experience is that the urea breath test is usually negative, except where *H. pylori* co-exists. A serological test for *H. suis* would provide a useful screen for infection, or any memory of it, in potentially high-risk patient or occupational groups, but none is currently available. The 16S rDNA assay for '*Helicobacter heilmannii*-like organisms' (Chisholm et al., 2003) did not detect the *H. suis* found on RT-PCR. It may, in part, be a question of sensitivity. *Helicobacter heilmannii*-like DNA (Chisholm et al., 2003) had been detected in a urea breath test-positive case (Dobbs et al., 2005): re-evaluation by the methodology used here showed this to *be H. suis*. There appears to be advantage in taking more than one biopsy to determine NHPH status by molecular microbiology. Low infection load (Stolte et al., 1997) will hamper histological confirmation. Sensitivity and specificity of the RT-PCR will need to be determined for introduction into routine clinical use. This is not as problematic for *H. suis* as for other NHPH, since it is the only gastric *Helicobacter* isolated from pigs in Europe, where infection is very common and load high. The gold standard of seeing spiral gastric helicobacters, in a histological section adjacent to the biopsy for DNA extraction, can be applied. Greater frequency of *H. suis* in IP where *H. pylori* has been eradicated points to need for post-treatment endoscopic biopsy. New infection in a particularly susceptible host cannot be excluded, although adult transmission of another gastric *Helicobacter*, *H. pylori*, is unusual. It is likely that the IP patients were co-infected with *H. pylori* and *H. suis* prior to eradicating *H. pylori*, and *H. suis* filled the niche afterwards. Indeed, susceptibility testing of isolates from sows suggests relative intrinsic insensitivity to amoxicillin and metronidazole, greater intrinsic susceptibility to tetracycline (Vermoote et al., 2011). Current use of tetracycline in first-line treatment strategies is limited (Malfertheiner et al., 2012). Information on *in vitro* anti-microbial susceptibility of *H. suis* in man is needed.

Transmission of *H. suis*

Contact with pigs is a risk factor for human gastric *H. suis* infection (Meining et al., 1998). It was demonstrated, using MLST, that a *H. suis* strain from the stomach of a pig veterinarian with gastric complaints was closely related to porcine strains (Joosten et al., 2013). This illustrates the zoonotic potential of this species: direct contact with pigs can be a source for human *H. suis* infection. Moreover, *H. suis* can be present, and survive, in minced pork (De Cooman et al., 2013): raw or undercooked porcine products might be another source of infection. Complete MLST sequence types were not obtained for any of the *H. suis* strains from patients,

precluding comparison with sequence types present in pigs. Although all alleles of the housekeeping genes detected in patients have been found in pig *H. suis* strains (Liang et al., 2013), the patients' strains could be *H. suis*-like, human-adapted bacteria. Isolation of strains from patients' stomachs could throw light on their origin. Challenges are the fastidious nature of *H. suis*, and obtaining fresh human biopsies from infected mucosa, where infection is not confluent and biopsies are small, non-targeted and limited in number (Haesebrouck et al., 2009). Horizontal and/or vertical transmission of NHPH might contribute to explaining the epidemiology of IP. Whilst we do not know if NHPH can be passed between humans, most *H. pylori* infections are transmitted where there is close contact, as between parent or sibling and infant. This fits with the siblings of IP patients sharing facets of the syndrome and increased frequency of *H. pylori* seropositivity (Dobbs et al., 2008). Adult transmission of a causative factor is suggested by spouses of IP patients being a short, but highly significant, 'distance-down-the-pathway' (Dobbs et al., 2008), with marked, multifarious, relevant differences between spouses and control couples. These are difficult to explain by selective mating or learnt/reactive behavior.

Future directions

Irrespective of association of *H. suis* with PD, its known association with gastric pathology requires its detection to be added to diagnostic microbiology on gastric biopsies from patients in high-risk groups. Feedback of results of the service evaluation to individual IP patients, who were traceable by the single clinic, has been made, with the option of re-biopsy in those with historical *H. suis* infection. A natural progression of the work is to examine the archived samples for zoonotic helicobacters commonly associated with cats and dogs.

AUTHORSHIP

Guarantor of the article: Dr Sylvia Dobbs. Author contributions: RJD, SMD, IB and AL requested FH's expertise, in re-evaluating the NHPH assay used at Public Health England. FH, RJD and SMD were joint principal investigators. CB & AS contributed equally. CB, AS, BF, AL performed the laboratory work. CB, AS, AL, FH, SMD, RJD, AC collected and analyzed the data. AS, AL, AC, RJD, SMD, FH designed the study. CB, BF, FP, RD, DT, CW, IB contributed to the design. CB, AS, FH, IB, AC, SMD, RJD wrote the paper. All authors have approved the final version of the manuscript.

ACKNOWLEDGEMENTS

Declaration of personal interests: None. Declaration of funding interests: This work was supported by the Research Fund of Ghent University, Belgium, Code GOA 01G00408, and the Psychiatry Research Trust (Registered Charity no. 284286), London, UK, which received grants from the Cyril Corden Trust and the Cecil Pilkington Charitable Trust, and donations from Brian Newman & Louise Barton. Caroline Blaecher is a PhD student at the Department of Pathology, Bacteriology and Avian Diseases, Faculty of Veterinary Medicine, Ghent University, Belgium, and a Visiting Researcher in Pharmaceutical Science at King's College London. The authors would like to thank Nathalie Van Rysselberghe and Sofie De Bruyckere for their excellent technical assistance.

References

- Andersen LP, Norgaards A, Holck S, Blom J, Elsborg L. 1996. Isolation of a "*Helicobacter heilmannii*"-like organism from the human stomach. *Eur J Clin Microbiol Infect Dis* 15: 95-96.
- Calne DB, Snow BJ, Lee C. 1992. Criteria for diagnosing Parkinson's disease. *Ann Neurol* 32 Suppl: 125-127.
- Charlett A, Dobbs RJ, Dobbs SM, Weller C, Ibrahim MAA, Dew T, Sherwood R, Oxlade NL, Plant JM, Bowthorpe J, Lawson AJ, Curry A, Peterson DW, Bjarnason IT. 2009. Blood profile holds clues to role of infection in a premonitory state for idiopathic parkinsonism and of gastrointestinal infection in established disease. *Gut Pathogens* 1: 20.
- Chisholm SA, Owen RJ, Teare EL, Saverymuttu S. 2001. PCR-based diagnosis of *Helicobacter pylori* infection and real-time determination of clarithromycin resistance directly from human gastric biopsy samples. *J Clin Microbiol* 39: 1217-1220.
- Chisholm SA, Owen RJ. 2003. Development and application of a novel screening PCR assay for direct detection of "*Helicobacter heilmannii*"-like organisms in human gastric biopsies in Southeast England. *Diagn Microbiol Inf Dis* 46: 1-7.
- Cummings P. 2011. Estimating adjusted risk ratios for matched and unmatched data: An update. *The Stata Journal* 11 (Number 2): 290-298.
- Debongnie JC, Donnay M, Mairesse J. 1995. *Gastrospirillum hominis* ('*Helicobacter heilmannii*') : a cause of gastritis, sometimes transient, better diagnosed by touch cytology ? *Am J Gastroenterol* 90: 411-416
- De Cooman L, Flahou B, Houf K, Smet A, Ducatelle R, Pasmans F, Haesebrouck F. 2013. Survival of *Helicobacter suis* bacteria in retail pig meat. *Int J of Food Microbiol* 166(1): 164-167.
- Dobbs RJ, Dobbs SM, Weller C, Bjarnason IT, Oxlade NL, Charlett A, Al-Janabi MA, Kerwin RW, Mahler RF, Price AB. 2005. Role of Chronic Infection and Inflammation in the Gastrointestinal Tract in the Aetiology and Pathogenesis of Idiopathic Parkinsonism. Part 1: Eradication of *Helicobacter* in the Cachexia of Idiopathic Parkinsonism. *Helicobacter* 10: 267-275.
- Dobbs RJ, Dobbs SM, Weller C, Charlet A, Bjarnason IT, Curry A, Ellis D, Ibrahim MAA, Mccrossan M, O'Donohue J, Owen RJ, Oxlade NL, Sanderson J, Sudhanva M, Williams J. 2008. *Helicobacter* hypothesis for Idiopathic Parkinsonism: before and beyond. *Helicobacter* 13: 309-322
- Dobbs SM, Dobbs RJ, Weller C, Charlett A, Bjarnason IT, Lawson AJ, Letley D, Harbin L, Price AB, Ibrahim MAA, Oxlade NL, Bowthorpe J, Leckstroem D, Smee C, Plant JM, Peterson DW. 2010. Differential effect of *Helicobacter pylori* eradication on time-trends in brady/hypokinesia and rigidity in idiopathic parkinsonism. *Helicobacter* 15: 279-294.
- Dobbs RJ, Charlett A, Dobbs SM, Weller C, Ibrahim MAA, Iguodala O, Smee C, Plant JM, Lawson AJ, Taylor D, Bjarnason I. 2012. Leukocyte-subset counts in idiopathic parkinsonism provide clues to a pathogenic pathway involving small intestinal bacterial overgrowth. A surveillance study. *Gut Pathogens* 4: 12.
- Dobbs SM, Charlett A, Dobbs RJ, Weller C, Iguodala O, Smee C, Lawson AJ, Taylor D, Bjarnason I. 2013. Antimicrobial surveillance in idiopathic parkinsonism: indication-specific improvement in hypokinesia following *Helicobacter pylori* eradication and non-specific effect of antimicrobials for other indications in worsening rigidity. *Helicobacter* 18: 187-196.
- Haesebrouck F, Pasmans F, Flahou B, Chiers K, Baele M, Meyns T, Decostere A, Ducatelle R. 2009. Gastric helicobacters in domestic animals and nonhuman primates and their significance for human health. *Clin Microbiology Reviews* 202-223
- Ho SA, Hoyle JA, Lewis FA, Secker AD, Cross D, Mapstone NP, Dixon MF, Wyatt JL, Tompkins DS, Taylor GR. 1991. Direct polymerase chain reaction test for detection of *Helicobacter pylori* in humans and animals. *J Clin Microbiol* 29: 2543-2549.
- Javala K, On SL, Harrington CS, Andersen LP, Hänninen ML, Vandamme P. 2001. A cultured strain of "*Helicobacter heilmannii*" a human gastric pathogen, identified as *H. bizzozeronii*: evidence for zoonotic potential of *Helicobacter*. *Emerg Infect Dis* 7: 1036-1038.
- Joosten M, Flahou B, Meyns T, Smet A, Arts J, De Cooman L, Pasmans F, Ducatelle R, Haesebrouck F. 2013. Case Report : *Helicobacter suis* infection in pig veterinarian. *Helicobacter*. 18(5): 392-396.
- Kivistö R, Linros J, Rossi M, Rautelin H, Hänninen ML. 2010. Characterization of multiple *Helicobacter bizzozeronii* isolates from a finnish patient with severe dyspeptic symptoms and chronic active gastritis. *Helicobacter* 15(1): 58-66.
- Lee E, Burnett CA, Lulich N, Lorraine L, Cameron, Sestito JP. 2002. Proportionate mortality of crop and livestock farmers in the United States, 1984-1993. *Am J Ind Med* 42: 410-420.

20. Liang J, Ducatelle R, Pasmans F, Smet A, Haesebrouck F, Flahou B. 2013. Multilocus sequence typing of the porcine and human gastric pathogen *Helicobacter suis*. J Clin Microbiol 51: 920-926.
21. Malfertheiner P, Megraud F, O'Morain CA, Atherton J, Axon AT, Bazzoli F, Gensini GF, Gisbert JP, Graham DY, Rokkas T, El-Omar EM, Kuipers EJ; European Helicobacter Study Group. 2012. European *Helicobacter* study group: management of *Helicobacter pylori* infection- the Maastricht IV/Florence consensus report. Gut 61(5): 646-664.
22. Meining A, Kroher G, Stolte M. 1998. Animal reservoirs in the transmission of *Helicobacter heilmannii*. Results of a questionnaire-based study. Scandinavian Journal of Gastroenterology 33: 795-798
23. Morgner A, Bayerdorffer E, Meining A, Stolte M, Kroher G. 1995. *Helicobacter heilmannii* and gastric cancer. Lancet 346: 551-552.
24. O'Rourke JL, Solnick JV, Neilan BA, Seidel K, Hayter R, Hansen LM, Lee A. 2004. Description of '*Candidatus Helicobacter heilmannii*' based on DNA sequence analysis of 16S rRNA and urease genes. Int J Syst Evol Microbiol 54: 2203-2211.
25. Stolte, M., Kroher, G., Meining, A., Morgner, A., Bayerdtirffer E., Bethke B. 1997. A Comparison of *Helicobacter pylori* and *H. heilmannii* Gastritis. A Matched Control Study Involving 404 Patients. Scand J Gastroenterol 32: 28-33.
26. Strang RR. 1965. The association of gastroduodenal ulceration with Parkinson's disease. Med J Austr 52: 842-843.
27. Tansey MG, McCoy MK, Frank-Cannon TC. 2007. Neuroinflammatory mechanisms in Parkinson's disease: potential environmental triggers, pathways, and targets for early therapeutic intervention. Exp Neurol 208: 1-25.
28. Trebesius K, Adler K, Vieth M, Stolte M, Haas R. 2001. Specific detection and prevalence of *Helicobacter heilmannii*-like organisms in the human gastric mucosa by fluorescent in situ hybridization and partial 16S ribosomal DNA sequencing. J Clin Microbiol. 39: 1510-1516.
29. Van den Bulck K, Decostere A, Baele M, Driessen A, Debonnie JC, Burette A, Stolte M, Ducatelle R, Haesebrouck F. 2005. Identification of non-*Helicobacter pylori* spiral organisms in gastric samples from humans, dogs and cats. J Clin Microbiol 43: 2256-2260.
30. Vermoote M, Pasmans F, Flahou B, Van Deun K, Ducatelle R, Haesebrouck F. 2011. Antimicrobial susceptibility pattern of *Helicobacter suis* strains. Veterinary Microbiol 153: 339-342.
31. von Campenhausen S, Bornschein B, Wick R, Bötzel K, Sampaio C, Poewe W, Oertel W, Siebert U, Berger K, Dodel R. 2005. Prevalence and incidence of Parkinson's disease in Europe. Eur Neuropsychopharmacol 15: 473-490.
32. Wirdefeldt K, Adami H-O, Cole P, Trichopoulos D, Mandel J. 2011. Epidemiology and etiology of Parkinson's disease: a review of the evidence. Eur J Epidemiol 26: 51-58.
33. Wüppenhorst N, von Loewenich F, Hobmaier B, Vetter-Knoll M, Mohadjer S, Kist M. 2013. Culture of a gastric non-*Helicobacter pylori Helicobacter* from the stomach of a 14 year-old girl. Helicobacter 18(1): 1-5.
34. Zipp F, Aktas O. 2006. The brain as a target of inflammation: common pathways link inflammatory and neurodegenerative diseases. Trends Neurosci 19: 518.

Chapter 2

The choroid plexus epithelium as a novel player in the stomach-brain axis during *Helicobacter* infection

Gorlé N^{*,1,2}, Blaecher C^{*,3}, Bauwens E³, Balusu S^{1,2}, Vandewalle J^{1,2}, Van Wonterghem E^{1,2}, Van Imschoot G^{1,2}, Liu C³, Ducatelle R³, Libert C^{1,2}, Haesebrouck F^{3,#}, Smet A^{3,#} and Vandenbroucke RE^{1,2,#}

¹VIB Center for Inflammation Research, Ghent, Belgium, VIB, B-9052 Ghent, Belgium

²Department of Biomedical Molecular Biology, Ghent University, B-9052 Ghent, Belgium

³Department of Pathology, Bacteriology and Avian Diseases, Ghent University, B-9820 Merelbeke, Belgium

^{*,#}Both authors contributed equally

Paper submitted

Abstract

Several studies suggest a link between shifts in gut microbiota and neurological disorders. Recently, we reported a high prevalence of *Helicobacter suis* in patients with Parkinson's disease. Here, we evaluated the effect of acute gastric *H. suis* infection on the brain in mice. One month of infection with *H. suis* resulted in increased brain inflammation, reflected in activated microglia. Two communication pathways between the gastrointestinal tract and the brain were shown to be involved. We detected choroid plexus inflammation and disruption of the epithelial blood-cerebrospinal fluid (CSF) barrier upon *H. suis* infection, while the endothelial blood-brain barrier (BBB) remained functional. These changes were accompanied by leakage of the gastrointestinal barrier and low-grade systemic inflammation, suggesting that *H. suis*-evoked gastrointestinal permeability and subsequent peripheral inflammation induces changes in brain homeostasis. Additionally, *H. suis* infection caused changes in the signaling of vasoactive intestinal peptide (VIP). Interestingly, systemic administration of VIP reversed the inflammation-induced disruption of the blood-CSF barrier integrity. In conclusion, this study shows for the first time that *H. suis* infection induces inflammation in the brain and that the choroid plexus is a novel player in the stomach-brain axis.

Introduction

The healthy human gastrointestinal tract harbors approximately 100 trillion commensal bacteria (Clemente et al., 2012; Foster and McVey Neufeld, 2013). An increasing number of recent studies indicate that the shifts in the gastrointestinal microbiome caused by a broad variety of triggers, including gastrointestinal infections, might have a significant effect on our health (Ochoa-Reparaz and Kasper, 2014; Walters et al., 2014; Zhang et al., 2015). Indeed, dysbiosis has been associated with a wide range of human health problems (Clemente et al., 2012; Zhang et al., 2015; Boulange et al., 2016), including neurological disorders such as depression, anxiety-like behavior, autism, Alzheimer's disease and Parkinson's disease (Parracho et al., 2005; Bercik et al., 2011; Bravo et al., 2011; Zhang et al., 2015; Hu et al., 2016; Mulak and Bonaz, 2015).

The genus *Helicobacter* is a group of gram-negative, spiral-shaped bacteria colonizing the gastrointestinal tract of humans and animals. *Helicobacter pylori* (*H. pylori*) is the best-known gastric human pathogen and is one of the major causes of gastric cancer (Kusters et al., 2006). Besides *H. pylori*, other *Helicobacter* species, referred to as gastric non-*Helicobacter pylori Helicobacter* (NHPH) species, have also been associated with severe human gastric malignancies (Haesebrouck et al., 2009). The most common NHPH species in humans is *H. suis* (Haesebrouck et al., 2009), which naturally colonizes the stomach of pigs (Liang et al., 2013) and non-human primates (Bosschem et al., 2016). Most likely, transmission of *H. suis* occurs through direct or indirect contact between animals and humans. Additionally, *H. suis* can be present and survive in minced pork, suggesting that raw or undercooked pork may be a source of *H. suis* infection of humans (De Cooman et al., 2013).

Recently, an exceptionally high frequency (27%) of *H. suis* DNA was found in gastric biopsies from human patients with Parkinson's disease compared to a control group with no clinical parkinsonism (2%) (Blaecher et al., 2013). Besides, *H. suis* DNA was also detected in a blood sample from a patient with both parkinsonism and Alzheimer's disease (unpublished results). The patient was a middle-aged male, who was treated with tetracycline, clarithromycin, bismuth and a proton pump inhibitor to eradicate *H. suis*. This therapy resulted in a temporary improvement of the well-being as well as the gastric and neurological symptoms of the patient (Dobbs et al., 2013). This finding suggests that the inflammatory effects caused by *Helicobacter* are not restricted to the stomach.

To prevent disturbance of brain homeostasis due to alterations in the systemic blood composition, the brain is protected by a series of barriers, including the blood–brain barrier (BBB) and the blood–cerebrospinal fluid (CSF) barrier (De Bock et al., 2014). It has been suggested that systemic inflammation may contribute to the impairment of the BBB (Kortekaas et al., 2005; Stolp and Dziegielewska, 2009) and the blood–CSF barrier (Vandenbroucke et al., 2012; Demeestere et al., 2015). Furthermore, several neurodegenerative disorders, including Parkinson's disease and Alzheimer's disease, have been associated with disruption of the BBB (Stolp and Dziegielewska, 2009; Lee and Pienaar, 2014; Li et al., 2015) and the blood–CSF barrier (Vawter et al., 1996; Pisani et al., 2012; Bergen et al., 2015; Balusu et al., 2016a; Gorle et al., 2016). Because it has been proposed that *Helicobacter* infections cause systemic inflammation (Blaecher et al., 2013; Alvarez-Arellano and Maldonado-Bernal, 2014), gastric infection with *H. suis* could influence the integrity of the brain barriers and thereby contribute to the pathophysiology of neurodegenerative disorders.

In this study, we investigated the effect of gastric *H. suis* infection on brain homeostasis. Our results show that *H. suis* induces inflammation not only in the stomach, but also in the brain. Moreover, we identified two possible pathways of communication between the gastrointestinal tract and the brain. On the one hand, leakage of the gastrointestinal barrier induced systemic inflammation and caused loss of the blood-CSF barrier integrity. On the other hand, *H. suis*-induced alterations in VIP signaling might affect brain homeostasis because we found that administration of VIP can reverse systemic inflammation-induced blood-CSF barrier leakage and brain inflammation.

Materials and methods

Mice. For the *Helicobacter* infection experiments, female C57BL/6 mice (5 weeks old) were purchased from Harlan (Harlan Laboratories, Horst, The Netherlands). The animals were housed in filtertop cages in groups of 5-10 mice/cage with free access to food and water and with a 12-hour light/12-hour dark cycle in a conventional animal house. All experiments were approved by the ethical committee of the Faculty of Veterinary Medicine of Ghent University. For the VIP experiment, female C57BL/6 mice (7 weeks old) were purchased from Janvier (Janvier Labs, Le Genest-Saint-Isle, France). Mice were housed with 4-6 mice/cage in a specific pathogen-free animal facility with free access to food and water and with a 14-hour light/10-hour dark cycle. This experiment was approved by the ethics committee of the Faculty of Sciences of Ghent University.

Cultivation of *H. suis* bacteria. *H. suis* bacteria (strain HS1) were grown biphasically on 1.5% *Brucella* agar (Immunosource, Halle-Zoersel, Belgium), supplemented with 1 ml skirrow (Oxoid, Hampshire, United Kingdom), 3 ml vitox (Oxoid, United Kingdom), 5 µg/ml amphotericin B (Sigma-Aldrich, Germany), 0.05% HCL (Sigma-Aldrich, Taufkirchen, Germany) and 20% inactivated fetal bovine serum (Perbio, Thermo Scientific Hyclone, Cramlington, United Kingdom) with on top *Brucella* broth (Immunosource, Belgium), supplemented with 0.05% HCL (Sigma-Aldrich, Germany) (Baele et al., 2008). The plates were incubated at 37°C under micro-aerobic conditions (10% CO₂, 5% O₂ and 85% N₂). Using a counting chamber (Neubauer Improved Assistant) the bacteria were counted and diluted to 10⁸ viable *H. suis* bacteria/ml.

In vivo infection procedure. At the age of 6 weeks, animals were anaesthetized with 2.5% isoflurane and intragastrically inoculated with either 300 µl of a stock solution containing 10⁸ viable *H. suis* bacteria/ml or with 300 µl of the growth medium of the bacteria (control group). One month after infection, mice were euthanized for further analysis by the administration of an overdose of ketamine/xylazine followed by decapitation. In order to perform the different analyses, mice were divided into 4 different groups: the first group of mice (10 controls and 10 *H. suis* infected mice) was used to determine blood-CSF barrier and BBB permeability, the second group (5 controls and 5 *H. suis* infected mice) was used for quantification of gastrointestinal permeability, the third (5 controls and 5 *H. suis* infected mice) and fourth group (10 controls and 10 *H. suis* infected mice) were used for immunohistochemical stainings and DNA, RNA and protein analyses of stomach and brain samples, respectively.

VIP experiment. Mice were intraperitoneally (ip) injected with lipopolysaccharide (LPS) from *Salmonella enterica* ser. Abortus equi (Sigma) dissolved in D-PBS. The dose was 120 µg/20 g body weight. Control animals received ip injections of D-PBS. Additionally, mice were injected twice with 5 nmol VIP (05-23-2101-1MG,

Millipore, Merck, Overijse, Belgium) intravenously (iv): together with and 2h after LPS injection. Six hours after LPS injection, CSF was isolated and used to determine blood-CSF permeability as described below and choroid plexus and hippocampus were isolated for gene expression analysis.

Tissue collection. For RNA and protein analysis, mice were transcardially perfused with D-PBS/heparin (0.2% heparin) supplemented with 0.5% bromophenol blue. The stomach (including a small part of the duodenum) was collected and cut open via the curvatura major. After rinsing with sterile Hank's Buffered Salt Solution (Gibco by Thermo Fisher Scientific, Gaithersburg, USA), samples from the antrum and the corpus of the stomach and duodenum were collected and stored in RNA*later* (Qiagen, Antwerpen, Belgium) at -70°C or snap frozen in liquid nitrogen for RNA and protein extraction, respectively. Brain tissue was dissected out. The choroid plexus from all four ventricles was collected and snap frozen in liquid nitrogen, while hippocampus samples were stored in RNA*later* (Qiagen).

For immunohistochemical analyses, mice were transcardially perfused with 4% paraformaldehyde (PFA). Brain tissue and samples from the antrum and corpus of the stomach and duodenum were collected. The brain was divided into the left and right hemisphere. The right hemisphere was successively fixated in 4% PFA and embedded in paraffin for paraffin sections. The stomach and duodenum samples were either embedded in cryoprotectant (Thermo Scientific™, Waltham, USA), underwent rapid freezing on dry ice and were stored at -70°C or were fixated overnight in a solution of 3.7% formaldehyde (Chem-lab, Zedelgem, Belgium) and subsequently embedded in paraffin (Labo-nord, VWR, Leuven, Belgium).

DNA extraction and quantification of colonizing *H. suis* bacteria in the stomach and duodenum. Stomach and duodenum samples stored in RNA*later* were homogenized using the MagNAlyser (Roche diagnostics, Vilvoorde, Belgium) and RNA and DNA were separated as described before (Joosten et al., 2013). The number of colonizing *H. suis* bacteria per mg tissue was determined in the DNA samples using Real-Time (RT)-quantitative (q) Polymerase Chain Reaction (PCR). RT-qPCR detection of *H. suis* DNA was performed using the *ureA*-based primers (**Table 1**) (Blaecher et al., 2013) and the standard was generated as described before (O'Rourke et al., 2004). Standard and samples were run in duplicate on a CFX384™ RT-qPCR System with a C1000 Thermal Cycler (Bio-Rad, Temse, Belgium) according to the manufacturer's instructions.

RNA extraction and RT-qPCR for gene expression. Using the RNeasy mini kit (Qiagen), total RNA was extracted from the antrum and the corpus of the stomach, the duodenum, hippocampus and choroid plexus according to the manufacturer's instructions and measured using Nanodrop equipment (Nanodrop ND-1000, Fisher Scientific). cDNA was synthesized using a cDNA Synthesis Kit (Bio-Rad). RT-qPCR analysis of the stomach and duodenum samples was performed on a CFX384™ RT-qPCR System with a C1000 Thermal Cycler (Bio-Rad), using the 2xSensimix (Sybr No-Rox kit; Bioline, London, UK), while for brain samples the RT-qPCR was performed on the Light Cycler 480 system (Roche) using the 2xSensimix (Sybr No-Rox kit; Bioline). Expression levels were normalized to the expression of 3 reference genes: *Hprt*, *H2afz* and *Ppia* for stomach and duodenum samples, *Hprt*, *Rpl* and *Ubc* for choroid plexus samples and *Gapdh*, *Rpl* and *Ubc* for hippocampus samples. The primer sequences are shown in **Table 1**.

Cytokine/chemokine measurements. The snap frozen gastrointestinal samples were lysed using lysis buffer containing 0.5% CHAPS (Sigma-Aldrich) and a protease inhibitor complete tablet (Roche Applied Science; 11

873 580 001). Protein concentration of the samples was measured using the Pierce BCA protein assay kit (ThermoFisher Scientific). Cytokines and chemokines in serum, CSF and protein lysates were measured using the Bio-Plex cytokine assays (Bio-Rad) according to the manufacturer's instructions.

Quantification of the gastrointestinal permeability. Gastrointestinal permeability was determined as previously described (Vandenbroucke et al., 2014; Van Hauwermeiren et al., 2015). Briefly, 4 kDa FITC-dextran (Sigma) was intragastrically administered five hours before collection of blood. Blood was collected in EDTA tubes (Sarstedt, Wexford, Ireland) by means of heart puncture followed by centrifugation. Subsequently, plasma was isolated and diluted twice in sterile D-PBS. Gastrointestinal leakage was determined by measurement of fluorescence at $\lambda_{\text{ex}}/\lambda_{\text{em}} = 488/520$ nm.

Quantification of blood-CSF barrier and BBB permeability. Blood-CSF barrier and BBB permeability were determined as previously described (Vandenbroucke et al., 2012; Brkic et al., 2015). Briefly, 4 kDa FITC-dextran (Sigma) was injected iv 1 h before CSF collection. CSF was obtained from the fourth ventricle using the cisterna magna puncture method (Vandenbroucke et al., 2012; Brkic et al., 2015). Subsequently, mice were perfused with D-PBS/heparin (0.2% heparin) and brain tissue was isolated. CSF samples were diluted 100-fold in sterile D-PBS, and blood-CSF barrier leakage was determined by measurement of fluorescence at $\lambda_{\text{ex}}/\lambda_{\text{em}} = 488/520$ nm. Brain samples were cut into small pieces, incubated overnight at 37°C in formamide while shaking and supernatant was collected after centrifugation for 15 min at maximum speed. Brain fluid was diluted 2-fold in sterile D-PBS, and BBB leakage was determined by measurement of fluorescence at $\lambda_{\text{ex}}/\lambda_{\text{em}} = 488/520$ nm.

Histopathology and immunohistochemistry. In order to evaluate inflammation in the stomach and duodenum, paraffin sections of 5 μm were cut using a microtome (Microm, Prosan, Thermo-Fisher). Sections were stained with Hematoxylin Gill III Prosan (Merck) and Eosine Yellow (VWR) according to standardized protocols. Gastric tissue was examined using a light microscope (Leica DM2000). The H&E stainings of the stomach were used to determine the grade of overall gastritis, using the modified visual analog scale similar to the adapted Updated Sydney System as previously described (Flahou et al., 2010).

For zonula occludens 1 (ZO1) immunostaining of the stomach, 30 μm cryosections were cut using a cryostat (Micron HM500) and mounted on slides. After air drying for 1 h, sections were fixated with 1% PFA for 10 min, washed for 5 min in PBS, and permeabilized for 10 min with 0.1% NP-40. After two rinses with PBS, samples were blocked for 1 h at room temperature with 5% goat serum and then incubated with primary antibody (Invitrogen, ZO1, 1:100, Merelbeke, Belgium). After incubation overnight at 4°C, they were washed with PBS-T and incubated with secondary antibody (Thermo Scientific; goat anti-rabbit-DyLight633, 1:300) diluted in 5% serum for 90 min at room temperature. Finally, a Hoechst staining was performed to counterstain the nuclei, the slides were dehydrated and xylene-based mounting medium was applied. Cells were visualized using a confocal microscope (Zeiss LSM780).

For the MUC13 staining of the stomach sections of 5 μm were cut from paraffin embedded tissue, using a microtome (Microm, Prosan, Thermo-Fisher), and placed on a microscope slide. Deparaffinization was performed by the addition of successively, xylene for 10 min, isopropyl alcohol for 5 min, 95% ethanol for 5 min, 50% ethanol for 5 min and aqua destillata for 5 min. Antigen retrieval was performed in citrate buffer (10 mM, pH6) using a pressure cooker, where after samples were rinsed with washing buffer (Dako, Santa-Clara,

USA). Endogenous peroxidase was blocked for 5 min with peroxidase blocking reagent (Dako) and rinsed with aqua destillata and washing buffer (Dako). Successively slides were incubated for 30 min with the primary in-house polyclonal mouse MUC13 antibody, diluted in antibody diluent (1/1000, Dako, S3022). After a rinse step with washing buffer (Dako), samples were incubated with labeled polymer-HRP anti-mouse for 30 min at room temperature (Evision+ system-HRP (DAB) (K4007)). Samples were rinsed 2 times with washing buffer (Dako), where after DAB + substrate (Dako) and DAB + chromogen (Dako) were added during 5 min. Finally, a counterstaining with hematoxyllin was performed during 10 min, followed by a rinse step of 1 min in running tap water and a dehydration step. Gastric tissue was visualized using a light microscope (Leica DM2000).

For IBA1 immunostaining, paraffin sections of the brains were processed as described before using the anti-IBA1 antibody (Wako, 019-19741, Neuss, Germany) (Brkic et al., 2015). Imaging of whole brain sections was done using Zeiss Axio Scan.Z1 followed by quantification using Fiji (<http://fiji.sc/Fiji>). Quantification of the percentage of brown color was done via color thresholding, with correction for the total amount of tissue, while microglia were classified into resting and activated in the subventricular zone according to adopted criteria (Hains and Waxman, 2006). Representative images were taken using an Olympus BX51 microscope.

For Claudin 1 (CLDN1) (Invitrogen, 51-9000) and VIP (Immunostar, 20077, Hudson, USA) immunostaining, paraffin-embedded brain and stomach tissues were sectioned at 5 μ m, dewaxed, and rinsed in water and PBS. Antigen retrieval was done using citrate buffer (Dako; S2031), followed by washing in PBS. For CLDN1 staining, endogenous peroxidase was blocked by incubating the slides in 3% H₂O₂ in methanol for 10 min, followed by washing in PBS. All samples were blocked with 5% goat serum in antibody diluent (Dako; S2022) for 30 min at RT, followed by incubation overnight at 4°C with primary antibody (CLDN1, Invitrogen, 51-9000, 1:100; VIP, Immunostar, 1/50). The next day, slides were washed with PBS. For CLDN1 immunostaining, slides were incubated with secondary antibody coupled to HRP (Dako; E-K4003), and washed with PBS. An amplification step was performed using tyramide (TSA kit, Perkin Elmer, Zaventem, Belgium) and visualization was done using ABC (Vector, Brussel, Belgium) and DAB. Slides were counterstained with haematoxylin, dehydrated and xylene-based mounting media was applied. Cells were visualized using an Olympus BX51 microscope. For VIP immunostaining, slides were incubated with secondary antibody (Thermo Scientific; goat anti-rabbit-DyLight633, 1:1000) diluted in 5% serum for 1 h at room temperature. Finally, a Hoechst staining was performed to counterstain the nuclei, the slides were dehydrated and xylene based mounting medium was applied. Cells were visualized using a confocal microscope (Leica TCS SP5).

Statistics. Data were analyzed by Mann-Whitney test using GraphPad Prism or SPSS and are presented as means \pm standard error of mean (SEM), unless stated otherwise. Significance levels are indicated on the graphs *, $0.01 \leq P < 0.05$; **, $0.001 \leq P < 0.01$; ***, $0.0001 \leq P < 0.001$; ****, $P < 0.0001$.


Results

Colonization by *H. suis* induces gastric inflammation


One month after infection, colonization by *H. suis* was confirmed by analyzing the stomach and small intestine. The mouse stomach consists of a non-glandular part and a glandular part. In this study, we analyzed only the glandular part, consisting mainly of the corpus and the antrum of the stomach. As shown in Figure 1a, the corpus and antrum of the stomach of *H. suis*-infected animals were colonized by an average of 4.51×10^5 and 7.30×10^5 bacteria per mg tissue, respectively. No *H. suis* DNA was detected in the stomach of control mice (Figure 1a). Additionally, *H. suis* DNA was also detected in the duodenum of infected mice but at a lower level than in the stomach (Figure 1a).

Analysis of H&E-stained stomach of control animals showed no or negligible infiltration of mononuclear and polymorphonuclear cells in the mucosa and the submucosa, which agrees with the normal histological image of the stomach (Figure 1b). In contrast, pronounced infiltration of mononuclear and polymorphonuclear cells was detected in the stomach of *H. suis*-infected animals (Figure 1c), indicating gastric inflammation as scored by the Updated Sydney System (Figure 1d). Furthermore, inflammation was also associated with changes in the expression levels of pro-inflammatory cytokines. Gene expression analyses revealed that the mRNA levels of the murine interleukin 8 (*IL8*) homologue keratinocyte chemoattractant (*Kc*) and lipopolysaccharide-induced CXC chemokine (*Lix*) were significantly up-regulated in corpus (Figure 1e) and in antrum (Figure 1f) of *H. suis*-infected mice. A significant increase in gene expression of *Il1 β* was observed only in the corpus of those animals (Figure 1e) and a significant decrease of *Il10* only in the antrum (Figure 1f). Additionally, a significant increase in protein levels of IL1 β and IL17 was observed in the corpus of the stomach of *H. suis*-infected mice (Figure 1h-i). In contrast, no significant changes in the expression of cytokines and chemokines were found in the duodenum of mice infected with *H. suis* (Figure 1g).


Figure 1



b



c







Figure 1. Analysis of colonization and inflammation in the stomach and duodenum after one month of *Helicobacter suis* (*H. suis*) infection. (a) *H. suis* colonization in stomach (corpus and antrum) and duodenum of C57BL/6 mice one month after *H. suis* infection compared to control mice (n = 7-10). (b-c) Representative H&E images of stomach sections of control (b) and *H. suis*-infected (c) C57BL/6 mice one month after infection. (d) Inflammation score based on H&E staining of stomach of *H. suis*-infected and control mice (n = 5). (e) Inflammatory gene expression analysis (represented as fold change) of interleukin 1 β (*Il1β*), lipopolysaccharide-induced CXC chemokine (*Lix*), keratinocyte chemoattractant (*Kc*), *Il10* and *Il17* in corpus of stomach in C57BL/6 mice after one month of infection with *H. suis* compared to control mice (n = 7-10). (f-g) Expression of the inflammatory genes interleukin 1 β (*Il1β*), lipopolysaccharide-induced CXC chemokine (*Lix*), keratinocyte chemoattractant (*Kc*), *Il10* and *Il17* in the antrum of the stomach (f) and in the duodenum (g) in C57BL/6 mice after one month of infection with *Helicobacter suis* (*H. suis*) compared to control mice, represented as fold change (n = 3-9). (h-i) Levels of IL1 β (h) and IL17A (i) in protein lysates determined using Bioplex of stomach (corpus and antrum) and duodenum after inoculation with *H. suis* or growth medium. Data were analyzed by Student's t-test (n = 5).

Gastric *H. suis* infection induces inflammation and activation of microglia in the brain

To determine whether *H. suis* affects the brain, cerebrospinal fluid (CSF) and brain tissue were analyzed for inflammation. Bioplex analysis of CSF revealed no significant changes in cytokines and chemokines (Figure 2a).

However, a significantly higher concentration of soluble tumor necrosis factor receptor 1 (sTNFR1) was found in the CSF of *H. suis*-infected mice (Figure 2b). Despite the limited changes in the CSF, a significantly stronger expression of *Il6* was detected in the brain hippocampus, while *Tnf*, *Il1 β* , *Kc* and inducible nitric oxide synthase (*iNos*) expression was unaffected (Figure 2c). Since microglia are important mediators of brain inflammation, we stained brain sections for ionized calcium binding adaptor molecule 1 (IBA1), a marker for microglia (Figure 2d-e). Automatic analysis showed a significant increase in the total IBA1-positive signal in the brain (Figure 2f). Additionally, manual quantification of the percentage of activated microglia in the subventricular zone, based on the morphology of IBA1-positive cells, revealed increased numbers of ramified microglia in *H. suis*-infected animals compared to control mice (Figure 2g). These data indicate that *H. suis* activates a pro-inflammatory response in the brain.

Figure 2



e





Figure 2. Analysis of brain inflammation in response to one month *Helicobacter suis* (*H. suis*) infection. (a) Levels of interleukin 1 β (IL1 β), IL6, tumor necrosis factor (TNF) and keratinocyte chemoattractant (KC) in cerebrospinal fluid (CSF) determined using Bioplex. Data were analyzed by Student's T-test (n = 4-5). (b) Concentration of TNFR1 in the cerebrospinal fluid (CSF) of control mice and mice infected with *H. suis* (n = 10). (c) Expression of inflammatory genes *Il1b*, *Il6*, *Tnf*, *Kc* and inducible nitric oxide synthase (iNos) in the hippocampus of C57BL/6 mice infected with *H. suis* compared to control mice, represented as fold change. Data were analyzed by Student's T-test (n = 3-5). (d-e) Representative images of ionized calcium-binding adapter molecule 1 (IBA1) staining of brain sections at the subventricular zone from C57BL/6 of control mice (d) and of mice after one month infection with *H. suis* (e). (f) Quantification of the percentage brown staining (IBA1) in the whole brain of *H. suis*-infected and negative control mice (n = 5). (g) Quantification of the percentage activated microglia in *H. suis*-infected and negative control mice (n = 5).

***H. suis* infection induces loss of the blood-CSF barrier but does not affect BBB integrity.**

The presence of tight brain barriers, including the BBB and the blood-CSF barrier, ensure brain homeostasis (De Bock et al., 2014), but several inflammatory triggers are known to disrupt barrier integrity (Nielsen et al., 2012; Vandenbroucke et al., 2012; Alvarez-Arellano and Maldonado-Bernal, 2014; Brkic et al., 2015; Gorle et al., 2016). Here, we studied whether the effect of *H. suis* on the brain is associated with loss of barrier integrity. No change in BBB permeability was observed (Figure 3a). In contrast to the BBB, a significant increase in blood-CSF barrier permeability was seen after one month of infection (Figure 3b). Gene expression analysis of the blood-CSF barrier located at the choroid plexus revealed that several tight-junction genes were affected by *H. suis*

infection. As shown in Figure 3c, gene expression of claudin 1 (*Cldn1*) and occludin (*Ocln*) was significantly down-regulated in the presence of *H. suis* infection. Other tight-junction genes, such as *Cldn5*, zona occludens 1 (*Zo1*) and *Zo3* decreased, but not significantly so. Finally, immunostaining was performed to determine the subcellular localization of CLDN1. The arrowheads in Figure 3d-e point to loss of apical localization of CLDN1 in the choroid plexus epithelial cells of *H. suis*-infected mice. These data indicate that *H. suis* infection compromises the functionality of the blood–CSF barrier but does not affect the integrity of the BBB.

Figure 3





Figure 3. Brain barrier analysis in response to one month of *Helicobacter suis* (*H. suis*) infection. (a) Relative blood-brain barrier (BBB) permeability one month after inoculation with growth medium or *H. suis* ($n = 5$). **(b)** Relative blood-cerebrospinal fluid (CSF) barrier permeability one month after inoculation with growth medium or *H. suis* ($n = 8-10$). **(c)** Expression of the tight junction genes claudin 1 (*Cldn1*), *Cldn5*, occludin (*Ocln*), zonula occludens (*Zo1*) and *Zo3* in choroid plexus tissue from C57BL/6 mice infected with *H. suis* compared to control mice ($n = 5$). **(d-e)** Representative images of choroid plexus tissue stained for CLDN1 of control mice (d) and *H. suis*-infected mice (e). CP: choroid plexus.

***H. suis* infection induces loss of gastric mucosal barrier integrity and subsequent systemic inflammation**

Increased gastrointestinal permeability, which could lead to leakage of pro-inflammatory mediators and bacterial metabolites from the gut into the systemic circulation, is considered one of the gut–brain axis pathways (Cryan and Dinan, 2012; Mayer et al., 2014). In our study, we used oral gavage of FITC-dextran to investigate the influence of gastric *H. suis* infection on the gastrointestinal barrier. Plasma fluorescence analysis revealed a significant increase in gastrointestinal permeability after one month of *H. suis* infection (Figure 4a). The gastric mucosal barrier is composed of the gastric mucosa (epithelium, lamina propria and muscularis mucosa), with a mucus layer on top (Linden et al., 2002; Kawakubo et al., 2004; Yamashiro, 2006; McGuckin, 2011 #321). This gastric mucus layer, consisting mainly of secreted mucins, is the first barrier pathogens encounter. The membrane-associated mucins, located underneath this mucus layer, also provide a barrier, thereby limiting access to the epithelial cell surface (McGuckin, 2011). Changes in mucin distribution and expression might affect the integrity of the gastric mucosal barrier (Kufe, 2009). Therefore, we analyzed gastric mucin expression. RT-qPCR analysis revealed a significantly increased gene expression of the transmembrane mucin 13 (*Muc13*) in the corpus of *H. suis*-infected mice, while the expression of *Muc5ac*, *Muc5b* and *Muc6* were unaffected (Figure 4b). This finding was confirmed histologically (Figure 4e-f): compared to control mice (Figure 4e), apical membrane and cytoplasmatic MUC13 staining (brown) was much more pronounced in the mucus-secreting epithelial cells of the stomach of *H. suis*-infected mice (Figure 4f). In contrast, *Muc13* gene expression did not change in antrum and was significantly down-regulated in duodenum of *H. suis*-infected animals (Figure 4c-d).

The gastric epithelium located underneath the mucus layer is formed by epithelial cells connected by tight junction proteins (Caron et al., 2015). We performed immunostaining to determine whether the *H. suis*-induced alterations in gastrointestinal barrier integrity were correlated with changes in junctional proteins in the stomach. Confocal imaging revealed that expression of ZO1 was disturbed. Compared to control mice, in which ZO1 was enriched mainly at the apical side of the stomach epithelial cells (Figure 4g), the ZO1 signal was much more diffuse in stomach samples from *H. suis*-infected animals (Figure 4h). The changes in the expression of both MUC13 and ZO-1 might thus explain the increased gastrointestinal leakage in the presence of *H. suis* infection.

Figure 4






Figure 4. Analysis of the gastrointestinal barrier after one month of *Helicobacter suis* (*H. suis*) infection. (a) Relative gastrointestinal permeability of control mice compared to mice infected with *H. suis* (n = 5). **(b)** Expression of mucins *Muc5ac*, *Muc5b*, *Muc13* and *Muc6* in the corpus of the stomach of C57BL/6 mice inoculated with *H. suis* and of control mice, represented as fold change (n = 7-10). **(c-d)** Expression of the mucin *Muc13* in antrum of the stomach (c) and in duodenum (d) in C57BL/6 mice one month after infection with *Helicobacter suis* (*H. suis*) compared to control mice, represented as fold change (n = 5-9). **(e-f)** Representative images of immunostaining of MUC13 in the stomach of control mice (e) and *H. suis* infected mice (f). **(g-h)** Representative confocal images of zonula occludens 1 (ZO1) staining in the stomach of control mice (g) and *H. suis*-infected C57BL/6 mice (h).

Disruption of the gastric mucosal barrier might result in leakage of pro-inflammatory cytokines, chemokines and bacterial metabolites from the stomach into the blood, leading to systemic inflammation (Cryan and Dinan, 2012). Additionally, recent studies indicated that alterations in circulating cytokine levels may directly influence the functionality of the brain barriers and the brain (Cryan and Dinan, 2012; Vandenbroucke et al., 2012; Balusu et al., 2016b). Therefore, the presence of systemic inflammation was evaluated by analyzing the serum of

control and *H. suis*-infected mice. As shown in Figure 5a, the levels of the pro-inflammatory cytokine IL1 β and chemokine MIP1 α increased significantly in the serum of *H. suis*-infected mice. These findings indicate that the loss of blood-CSF barrier integrity observed in our study might have been caused by a low grade, systemic inflammation induced by *H. suis*. In agreement with this finding, gene expression analysis of choroid plexus tissue revealed a significantly increased expression of pro-inflammatory cytokines *Il1b* and *iNos* (Figure 5b).

Figure 5





Figure 5. Analysis of systemic inflammation after one month of *Helicobacter suis* (*H. suis*) infection. (a) Levels of serum interleukin (IL) 1 β , IL6, macrophage inflammatory protein (MIP1 α), tumor necrosis factor (TNF) and keratinocyte chemoattractant (KC) determined using Bioplex after inoculation of mice with *H. suis* or growth medium,. Data were analyzed by Student's t-test (n = 3-4). **(b)** Expression of the inflammatory genes *Il1b*, *Il6*, *Tnf*, *Kc* and *Inos* in choroid plexus tissue of C57BL/6 mice inoculated with *H. suis* or growth medium, represented as fold change (n = 3-5).

H. suis infection disturbs VIP signaling

Another important link between the gastrointestinal system and the brain are neuropeptides, such as the vasoactive intestinal peptide (VIP) (De Vadder et al., 2015). Indeed, VIP seems not only to be involved in the maintenance of the gastrointestinal epithelial barrier integrity (Conlin et al., 2009) but might also play a role in brain homeostasis (Delgado et al., 2004). Additionally, VIP reactive neurons have been found in the enteric nervous system and in the choroid plexus (Lindvall et al., 1978; Delgado et al., 2004). One month after infection with *H. suis*, the expression of *Vip* in the corpus of the stomach was significantly decreased (Figure 6a), whereas expression of the receptor, *Vipr1*, was significantly up-regulated and *Vipr2* remained stable (Figure 6a). Additionally, immunofluorescent stainings revealed that VIP was retained in the lumen of the gastric glands of *H. suis*-infected animals (Figure 6b-c). To evaluate whether alterations in VIP levels could affect the blood-CSF barrier integrity and thereby brain homeostasis in a LPS-induced systemic inflammation model, mice were treated with VIP. As in the infection with *H. suis*, injection of LPS in mice resulted in increased leakage of the blood-CSF barrier as described before (Vandenbroucke et al., 2012) (Figure 6d); this was reversed by VIP treatment (Figure 6d). LPS-induced increased leakage of the blood-CSF barrier was associated with significant down-regulation of the tight junction proteins *Cldn5*, *Ocln* and *Zo3* in the choroid plexus (Figure 6e). Downregulation of *Ocln* was significantly less in LPS-injected mice treated with VIP (Figure 6e). Expression of *Cldn1* was increased significantly by LPS injection, whereas expression of *Zo1* decreased when LPS and VIP were combined (Figure 6e). LPS injection also increased inflammatory gene expression in the choroid plexus,

reflected by an increase of *Il1b*, *Il6*, *Tnf*, *Kc* and *iNos*; the increases in *Il6*, *Tnf*, *Kc* and *iNos* were significantly reduced by VIP treatment (Figure 6f). Together, these data indicate that VIP might play a protective role in the preservation of brain homeostasis in the presence of systemic inflammation, and it might be important in the *H. suis*-associated changes in the brain.

Figure 6






Figure 6. Analysis of vasoactive intestinal peptide (VIP) signaling after one month of *Helicobacter suis* (*H. suis*) infection. (a) Expression of vasoactive intestinal peptide (*Vip*), VIP receptor (*VipR1*) and *VipR2* in the corpus of the stomach of C57BL/6 mice inoculated with *H. suis* or growth medium, represented as fold change ($n = 6-10$). (b-c) Representative confocal images of VIP staining in the stomach of control (d) and *H. suis* infected (e) C57BL/6 mice. (d) Relative blood-cerebrospinal fluid (CSF) barrier permeability as a result of systemic inflammation induced by intraperitoneal (ip) injection of 120 $\mu\text{g}/20\text{ g}$ body weight lipopolysaccharide (LPS), followed by intravenous (iv) injection of PBS or VIP, compared to controls ($n = 5-8$). (e) Relative expression of the tight junction proteins Claudin 1 (*Cldn1*), *Cldn5*, Occludin (*Ocln*), Zonula occludens 1 (*Zo1*) and *Zo3* in the choroid plexus as a result of systemic inflammation, followed by iv injection of PBS or VIP, compared to controls ($n = 8-10$). (f) Relative expression of the inflammatory cytokines interleukin 1 β (*Il1β*), *Il6*, tumor necrosis factor (*Tnf*), keratinocyte chemoattractant (*Kc*) and inducible nitric oxide synthase (*iNos*) in the choroid plexus as a result of systemic inflammation, followed by iv injection of PBS or VIP, compared to controls ($n = 8-10$).

Discussion

Alterations in the gut microbiome are believed to contribute to many different human health problems (Parracho et al., 2005; Bercik et al., 2011; Bravo et al., 2011; Clemente et al., 2012; Zhang et al., 2015). Next to the direct effects *H. pylori* infection on the gastrointestinal system (Kusters et al., 2006; Mayer et al., 2014; Li et al., 2016), it has been associated with the occurrence of different neurodegenerative disorders, including Alzheimer's disease (Wirdefeldt et al., 2011; Alvarez-Arellano and Maldonado-Bernal, 2014; Wong et al., 2014) and Parkinson's disease (Wirdefeldt et al., 2011; Blaecher et al., 2013; Alvarez-Arellano and Maldonado-Bernal, 2014; Wong et al., 2014). We recently highlighted in a similar way the importance of gastric *H. suis* infections in Parkinson's disease (Blaecher et al., 2013). However, it remains unknown how gastric *Helicobacter* infection in general influences the brain.

The gastrointestinal system and the brain are connected through neural, endocrine and immune pathways. Here, we demonstrated that one month of infection with *H. suis* resulted in increased brain inflammation. In particular, we observed a moderate but significant up-regulation of *Il6* gene expression in the hippocampus, elevated levels of sTNFR1 in CSF (a biomarker for Alzheimer's disease) (Jiang et al., 2011), and an increased amount of activated microglia in the brain. Interestingly, elevated brain levels of IL6 have also been reported in traumatic brain injury (Goodman et al., 1990; Kossmann et al., 1995), depression (Brietzke et al., 2009), Parkinson's disease (Boka et al., 1994) and Alzheimer's disease (Angelopoulos et al., 2008), suggesting a role for IL6 in the above-mentioned neuropathologies. Furthermore, microglia activation has been observed in *H. pylori* infection (Lo et al., 2009), and priming of the microglia by prior pathology accelerates the pathology in the brain (Cunningham, 2013). To ensure homeostasis in the CNS, the brain is protected from fluctuations in the systemic blood composition by a series of brain barriers, including the BBB and the blood-CSF barrier (De Bock et al., 2014). Our results indicate that gastric *H. suis* infection can induce leakage of the blood-CSF barrier but not of the BBB. This disturbance was accompanied by loss of tight junctions, and specifically by a significant decrease in the expression of *Cldn1* and *Ocln* and disturbed localization of CLDN1 protein in the choroid plexus epithelium.

Although the observed *H. suis*-induced blood-CSF barrier leakage was rather limited, dysfunction of the choroid plexus epithelium is known to be destructive to the brain (Redzic et al., 2005; Marques et al., 2007; Johanson et al., 2011; Balusu et al., 2016a). Moreover, increased blood-CSF barrier permeability has been observed in several neuroinflammatory disorders, including Alzheimer's disease, aging and Parkinson's disease (Pisani et al., 2012; Marques et al., 2013; Brkic et al., 2015; Balusu et al., 2016a; Gorle et al., 2016). The choroid plexus epithelium is uniquely positioned between blood and brain, and we and others believe that this blood-CSF interface is an essential player in the communication between the periphery and the brain. Indeed, several studies reported that choroid plexus epithelial cells sense inflammatory changes in the periphery, which in turn induces changes in functionality of the choroid plexus epithelial cells (Thouvenot et al., 2006; Marques et al., 2007; Marques et al., 2009; Vandenbroucke et al., 2012; Balusu et al., 2016b). Also in our study, we observed *H. suis*-induced systemic inflammation characterized by increased levels of IL1 β and MIP1 α in the serum, which could have contributed to disruption of the blood-CSF barrier and subsequently to brain inflammation.

Disruption of the gastric mucosal barrier might lead to leakage of inflammatory cytokines, chemokines and bacterial components from the stomach into the systemic circulation (Cryan and Dinan, 2012; Mayer et al., 2014), thereby inducing systemic inflammation. In our study, the extent of colonization of the stomach by *H. suis* was similar to what has been reported (Flahou et al., 2010). DNA from *H. suis* was also found in the duodenum, but it is not clear whether this pathogen colonized the duodenum or, otherwise, the qRT-PCR picked up DNA from bacteria colonizing the stomach. Furthermore, *H. suis* infection evoked a pro-inflammatory immune response in the stomach characterized by infiltration of innate immune cells and up-regulation of IL1 β and murine IL8 homologue genes *Lix* and *Kc*; these observations are in line with previous studies (Flahou et al., 2010; Flahou et al., 2012; Bosschem et al., 2016; Bosschem et al., In submission). Inflammation in the stomach was also accompanied by increased gastrointestinal barrier permeability and changes in the localization of the tight junction protein zonula occludens 1 (ZO1). Concordant with our study, co-culture of primary human gastric epithelial cells with *H. pylori* resulted in disruption of membrane ZO1 (Wroblewski, 2011).

Aberrant expression of transmembrane mucins has also been shown to play a role in disturbance of mucosal barrier integrity (Kufe, 2009). In our study, an inappropriate overexpression of the membrane-associated mucin MUC13 was seen. This mucin, which is predominantly expressed in the intestine and only at very low level in the stomach, seems to be induced by IL1 β in the inflamed stomach during *Helicobacter* infection (Cheng et al., 2016). Inappropriate overexpression of transmembrane mucins can affect gastric mucosal barrier integrity by disrupting cell polarity and cell-cell interactions (Kufe, 2009), but further investigations are necessary to confirm the role of MUC13 in gastric mucosal barrier dysfunction. Taken together, gastric *H. suis* infection induced inflammation in the stomach, which was associated with gastric mucosal damage resulting in leakage of inflammatory mediators into the blood and leading to systemic inflammation.

The gastrointestinal tract and the brain communicate through the systemic pathway, but they might also interact via neural communication (Mayer et al., 2014). Vasoactive intestinal peptide (VIP) is a neuropeptide with diverse functions and was recently identified as a component of the gut-brain axis (De Vadder et al., 2015). VIP has been shown to promote homeostasis of the mucosal barrier of the gastrointestinal tract and has been implicated in tight junction regulation (Conlin et al., 2009; Chen et al., 2015). In our study, VIP expression was changed in the stomach of *H. suis* infected animals, which agrees with the previously described altered density of VIP immunoreactive nerves in mice in response to *H. pylori* infection (Bercik et al., 2002). Alterations in VIP expression might thus also contribute to dysfunction of the gastric mucosal barrier seen in our study.

In the brain, VIP can prevent overactivation of microglia in LPS-induced inflammation (Kim et al., 2000; Delgado and Ganea, 2003). VIP has been described as protective in Parkinson's disease (Delgado and Ganea, 2003) and Alzheimer's disease (Song et al., 2012). To determine whether alterations in the peripheral VIP levels affect the blood-CSF barrier during peripheral inflammation, mice were injected in the peritoneum with LPS and then treated with VIP. Our results indicate that administration of VIP in LPS-induced disruption of brain homeostasis could reverse inflammation in the choroid plexus. VIP reactive nerves had been demonstrated in the choroid plexus (Lindvall et al., 1978) and their contribution to the regulation of choroid plexus activity had been

described (Nilsson et al., 1991). These reports are in line with our finding that administration of VIP reduces LPS-induced leakage of the blood–CSF barrier.

In conclusion, we have identified the pivotal role of the choroid plexus in the communication between the gastrointestinal tract and the brain. Our results indicate that disruption of the blood-CSF barrier is induced by *H. suis*-associated systemic inflammation and/or dysregulated VIP signaling. Although further research is required to unravel the specific mechanisms of gastrointestinal–brain communication, VIP is a potential candidate for treatment of neurodegenerative diseases because it can restore the blood-CSF barrier and thereby reduce the neuroinflammation associated with gastrointestinal infections.

Acknowledgments

Declaration of funding interests: This work was supported by the Research Fund of Ghent University, Belgium (01G00408 and 01IO071) and by the Research Foundation Flanders (FWO Vlaanderen). The authors thank Sofie De Bruyckere for her excellent technical assistance and the VIB Bio Imaging Core for suggestions concerning immunohistochemistry and imaging.

References

1. Alvarez-Arellano L, Maldonado-Bernal C (2014) *Helicobacter pylori* and neurological diseases: Married by the laws of inflammation. *World J Gastrointest Pathophysiol* 5:400-404.
2. Angelopoulos P, Agouridaki H, Vaiopoulos H, Siskou E, Doutsou K, Costa V, Baloyiannis SI (2008) Cytokines in Alzheimer's disease and vascular dementia. *Int J Neurosci* 118:1659-1672.
3. Baele M, Decostere A, Vandamme P, Ceelen L, Hellemans A, Mast J, Chiers K, Ducatelle R, Haesebrouck F (2008) Isolation and characterization of *Helicobacter suis* sp. nov. from pig stomachs. *Int J Syst Evol Microbiol* 58:1350-1358.
4. Balusu S, Brkic M, Libert C, Vandenbroucke RE (2016a) The choroid plexus-cerebrospinal fluid interface in Alzheimer's disease: more than just a barrier. *Neural Regen Res* (in press).
5. Balusu S, Van Wonterghem E, De Rycke R, Raemdonck K, Stremersch S, Gevaert K, Brkic M, Demeestere D, Vanhooren V, Hendrix A, Libert C, Vandenbroucke RE (2016b) Identification of a novel mechanism of blood-brain communication during peripheral inflammation via choroid plexus-derived extracellular vesicles. *EMBO Mol Med*.
6. Bercik P, De Giorgio R, Blennerhassett P, Verdu EF, Barbara G, Collins SM (2002) Immune-mediated neural dysfunction in a murine model of chronic *Helicobacter pylori* infection. *Gastroenterology* 123:1205-1215.
7. Bercik P, Denou E, Collins J, Jackson W, Lu J, Jury J, Deng Y, Blennerhassett P, Macri J, McCoy KD, Verdu EF, Collins SM (2011) The intestinal microbiota affect central levels of brain-derived neurotrophic factor and behavior in mice. *Gastroenterology* 141:599-609, 609 e591-593.
8. Bergen AA, Kaing S, ten Brink JB, Netherlands Brain B, Gorgels TG, Janssen SF (2015) Gene expression and functional annotation of human choroid plexus epithelium failure in Alzheimer's disease. *BMC Genomics* 16:956.
9. Blaecher C, Smet A, Flahou B, Pasmans F, Ducatelle R, Taylor D, Weller C, Bjarnason I, Charlett A, Lawson AJ, Dobbs RJ, Dobbs SM, Haesebrouck F (2013) Significantly higher frequency of *Helicobacter suis* in patients with idiopathic parkinsonism than in control patients. *Aliment Pharmacol Ther* 38:1347-1353.
10. Boka G, Anglade P, Wallach D, Javoy-Agid F, Agid Y, Hirsch EC (1994) Immunocytochemical analysis of tumor necrosis factor and its receptors in Parkinson's disease. *Neurosci Lett* 172:151-154.
11. Bosschem I, Flahou B, Van Deun K, De Koker S, Volf J, Smet A, Ducatelle R, Devriendt B, Haesebrouck F (In submission) Species-specific immunity to *Helicobacter suis*.
12. Bosschem I, Flahou B, Bakker J, Heuvelman E, Langermans J, De Bruybe E, Joosten M, Smet A, Ducatelle R, Haesebrouck F (2016) Comparative virulence of in vitro cultures primate- and pig-associated *Helicobacter suis* strains in a BALB/c mouse and Mongolian gerbil model. *Helicobacter* in press.
13. Boulange CL, Neves AL, Chilloux J, Nicholson JK, Dumas ME (2016) Impact of the gut microbiota on inflammation, obesity, and metabolic disease. *Genome Med* 8:42.
14. Bravo JA, Forsythe P, Chew MV, Escaravage E, Savignac HM, Dinan TG, Bienenstock J, Cryan JF (2011) Ingestion of *Lactobacillus* strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proc Natl Acad Sci U S A* 108:16050-16055.
15. Brietzke E, Stertz L, Fernandes BS, Kauer-Sant'anna M, Mascarenhas M, Escosteguy Vargas A, Chies JA, Kapczinski F (2009) Comparison of cytokine levels in depressed, manic and euthymic patients with bipolar disorder. *J Affect Disord* 116:214-217.

16. Brkic M, Balusu S, Van Wonterghem E, Gorle N, Benilova I, Kremer A, Van Hove I, Moons L, De Strooper B, Kanazir S, Libert C, Vandenbroucke RE (2015) Amyloid beta Oligomers Disrupt Blood-CSF Barrier Integrity by Activating Matrix Metalloproteinases. *J Neurosci* 35:12766-12778.
17. Caron TJ, Scott KE, Fox JG, Hagen SJ (2015) Tight junction disruption: *Helicobacter pylori* and dysregulation of the gastric mucosal barrier. *World J Gastroenterol* 21:11411-11427.
18. Chen L, Yuan W, Chen Z, Wu S, Ge J, Chen J, Chen Z (2015) Vasoactive intestinal peptide represses activation of tumor-associated macrophages in gastric cancer via regulation of TNFalpha, IL-6, IL-12 and iNOS. *Int J Oncol* 47:1361-1370.
19. Liu C, Rossi M, Linden S, Padra M, Blaecher C, Bauwens E, Joosten M, Flahou B, Vandenbrouck W, Ducatelle R, Haesebrouck F, Smet A (2016) The *Helicobacter heilmannii* hofE and hofF genes are essential for colonization of the gastric mucosa and play a role in IL-1beta-induced gastric MUC13 expression. *Helicobacter* 21:504-522.
20. Clemente JC, Ursell LK, Parfrey LW, Knight R (2012) The impact of the gut microbiota on human health: an integrative view. *Cell* 148:1258-1270.
21. Conlin VS, Wu X, Nguyen C, Dai C, Vallance BA, Buchan AM, Boyer L, Jacobson K (2009) Vasoactive intestinal peptide ameliorates intestinal barrier disruption associated with *Citrobacter rodentium*-induced colitis. *Am J Physiol Gastrointest Liver Physiol* 297:G735-750.
22. Cryan JF, Dinan TG (2012) Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour. *Nat Rev Neurosci* 13:701-712.
23. Cunningham C (2013) Microglia and neurodegeneration: the role of systemic inflammation. *Glia* 61:71-90.
24. De Bock M, Vandenbroucke RE, Decrock E, Culot M, Cecchelli R, Leybaert L (2014) A new angle on blood-CNS interfaces: a role for connexins? *FEBS Lett* 588:1259-1270.
25. De Cooman L, Flahou B, Houf K, Smet A, Ducatelle R, Pasmans F, Haesebrouck F (2013) Survival of *Helicobacter suis* bacteria in retail pig meat. *Int J Food Microbiol* 166:164-167.
26. De Vadder F, Plessier F, Gautier-Stein A, Mithieux G (2015) Vasoactive intestinal peptide is a local mediator in a gut-brain neural axis activating intestinal gluconeogenesis. *Neurogastroenterol Motil* 27:443-448.
27. Delgado M, Ganea D (2003) Neuroprotective effect of vasoactive intestinal peptide (VIP) in a mouse model of Parkinson's disease by blocking microglial activation. *FASEB J* 17:944-946.
28. Delgado M, Pozo D, Ganea D (2004) The significance of vasoactive intestinal peptide in immunomodulation. *Pharmacol Rev* 56:249-290.
29. Demeestere D, Libert C, Vandenbroucke RE (2015) Clinical implications of leukocyte infiltration at the choroid plexus in (neuro)inflammatory disorders. *Drug Discov Today* 20:928-941.
30. Dobbs SM, Dobbs RJ, Smet A, Weller C, Charlett A, Bjarnason I, Lawson AJ, Ibrahim MAA, Taylor D, Blaecher C, Pasmans F, Ducatelle R, Flahou B, Haesebrouck F (2013) Clinical significance of *Helicobacter suis* in idiopathic Parkinsonism. *Helicobacter* 18:149-149.
31. Flahou B, Deun KV, Pasmans F, Smet A, Volf J, Rychlik I, Ducatelle R, Haesebrouck F (2012) The local immune response of mice after *Helicobacter suis* infection: strain differences and distinction with *Helicobacter pylori*. *Vet Res* 43:75.

32. Flahou B, Haesebrouck F, Pasmans F, D'Herde K, Driessen A, Van Deun K, Smet A, Duchateau L, Chiers K, Ducatelle R (2010) *Helicobacter suis* causes severe gastric pathology in mouse and mongolian gerbil models of human gastric disease. PLoS One 5:e14083.
33. Foster JA, McVey Neufeld KA (2013) Gut-brain axis: how the microbiome influences anxiety and depression. Trends Neurosci 36:305-312.
34. Goodman JC, Robertson CS, Grossman RG, Narayan RK (1990) Elevation of tumor necrosis factor in head injury. J Neuroimmunol 30:213-217.
35. Gorle N, Van Cauwenberghe C, Libert C, Vandenbroucke RE (2016) The effect of aging on brain barriers and the consequences for Alzheimer's disease development. Mamm Genome.
36. Haesebrouck F, Pasmans F, Flahou B, Chiers K, Baele M, Meyns T, Decostere A, Ducatelle R (2009) Gastric helicobacters in domestic animals and nonhuman primates and their significance for human health. Clin Microbiol Rev 22:202-223, Table of Contents.
37. Hains BC, Waxman SG (2006) Activated microglia contribute to the maintenance of chronic pain after spinal cord injury. J Neurosci 26:4308-4317.
38. Hu X, Wang T, Jin F (2016) Alzheimer's disease and gut microbiota. Sci China Life Sci.
39. Jiang H, Hampel H, Prvulovic D, Wallin A, Blennow K, Li R, Shen Y (2011) Elevated CSF levels of TACE activity and soluble TNF receptors in subjects with mild cognitive impairment and patients with Alzheimer's disease. Mol Neurodegener 6:69.
40. Johanson C, Stopa E, McMillan P, Roth D, Funk J, Krinke G (2011) The distributional nexus of choroid plexus to cerebrospinal fluid, ependyma and brain: toxicologic/pathologic phenomena, periventricular destabilization, and lesion spread. Toxicol Pathol 39:186-212.
41. Joosten M, Blaecher C, Flahou B, Ducatelle R, Haesebrouck F, Smet A (2013) Diversity in bacterium-host interactions within the species *Helicobacter heilmannii sensu stricto*. Vet Res 44:65.
42. Kawakubo M, Ito Y, Okimura Y, Kobayashi M, Sakura K, Kasama S, Fukuda MN, Fukuda M, Katsuyama T, Nakayama J (2004) Natural antibiotic function of a human gastric mucin against *Helicobacter pylori* infection. Science 305:1003-1006.
43. Kim WK, Kan Y, Ganea D, Hart RP, Gozes I, Jonakait GM (2000) Vasoactive intestinal peptide and pituitary adenylyl cyclase-activating polypeptide inhibit tumor necrosis factor- α production in injured spinal cord and in activated microglia via a cAMP-dependent pathway. J Neurosci 20:3622-3630.
44. Kortekaas R, Leenders KL, van Oostrom JC, Vaalburg W, Bart J, Willemsen AT, Hendrikse NH (2005) Blood-brain barrier dysfunction in parkinsonian midbrain in vivo. Ann Neurol 57:176-179.
45. Kossmann T, Hans VH, Imhof HG, Stocker R, Grob P, Trentz O, Morganti-Kossmann C (1995) Intrathecal and serum interleukin-6 and the acute-phase response in patients with severe traumatic brain injuries. Shock 4:311-317.
46. Kufe DW (2009) Mucins in cancer: function, prognosis and therapy. Nat Rev Cancer 9:874-885.
47. Kusters JG, van Vliet AH, Kuipers EJ (2006) Pathogenesis of *Helicobacter pylori* infection. Clin Microbiol Rev 19:449-490.
48. Lee H, Pienaar IS (2014) Disruption of the blood-brain barrier in Parkinson's disease: curse or route to a cure? Front Biosci (Landmark Ed) 19:272-280.

49. Li J, Sung CY, Lee N, Ni Y, Pihlajamäki J, Panagiotou G, El-Nezami H (2016) Probiotics modulated gut microbiota suppresses hepatocellular carcinoma growth in mice. *Proc Natl Acad Sci U S A* 113:E1306-1315.
50. Li JC et al. (2015) Increased permeability of the blood-brain barrier and Alzheimer's disease-like alterations in slit-2 transgenic mice. *J Alzheimers Dis* 43:535-548.
51. Liang J, Ducatelle R, Pasmans F, Smet A, Haesebrouck F, Flahou B (2013) Multilocus sequence typing of the porcine and human gastric pathogen *Helicobacter suis*. *J Clin Microbiol* 51:920-926.
52. Linden S, Nordman H, Hedenbro J, Hurtig M, Boren T, Carlstedt I (2002) Strain- and blood group-dependent binding of *Helicobacter pylori* to human gastric MUC5AC glycoforms. *Gastroenterology* 123:1923-1930.
53. Lindvall M, Alumets J, Edvinsson L, Fahrenkrug J, Hakanson R, Hanks J, Owman C, Schaffalitzky de Muckadell OB, Sundler F (1978) Peptidergic (VIP) nerves in the mammalian choroid plexus. *Neurosci Lett* 9:77-82.
54. Lo YC, Shih YT, Wu DC, Lee YC (2009) In vitro effects of *Helicobacter pylori*-induced infection in gastric epithelial AGS cells on microglia-mediated toxicity in neuroblastoma SH-SY5Y cells. *Inflamm Res* 58:329-335.
55. Marques F, Sousa JC, Sousa N, Palha JA (2013) Blood-brain-barriers in aging and in Alzheimer's disease. *Mol Neurodegener* 8:38.
56. Marques F, Sousa JC, Correia-Neves M, Oliveira P, Sousa N, Palha JA (2007) The choroid plexus response to peripheral inflammatory stimulus. *Neuroscience* 144:424-430.
57. Marques F, Sousa JC, Coppola G, Geschwind DH, Sousa N, Palha JA, Correia-Neves M (2009) The choroid plexus response to a repeated peripheral inflammatory stimulus. *BMC Neurosci* 10:135.
58. Mayer EA, Knight R, Mazmanian SK, Cryan JF, Tillisch K (2014) Gut microbes and the brain: paradigm shift in neuroscience. *J Neurosci* 34:15490-15496.
59. Mulak A, Bonaz B (2015) Brain-gut-microbiota axis in Parkinson's disease. *World J Gastroenterol* 21:10609-10620.
60. Nielsen HH, Qiu J, Friis S, Wermuth L, Ritz B (2012) Treatment for *Helicobacter pylori* infection and risk of Parkinson's disease in Denmark. *Eur J Neurol* 19:864-869.
61. Nilsson C, Lindvall-Axelsson M, Owman C (1991) Simultaneous and continuous measurement of choroid plexus blood flow and cerebrospinal fluid production: effects of vasoactive intestinal polypeptide. *J Cereb Blood Flow Metab* 11:861-867.
62. O'Rourke JL, Solnick JV, Neilan BA, Seidel K, Hayter R, Hansen LM, Lee A (2004) Description of '*Candidatus Helicobacter heilmannii*' based on DNA sequence analysis of 16S rRNA and urease genes. *Int J Syst Evol Microbiol* 54:2203-2211.
63. Ochoa-Reparaz J, Kasper LH (2014) Gut microbiome and the risk factors in central nervous system autoimmunity. *FEBS Lett* 588:4214-4222.
64. Parracho HM, Bingham MO, Gibson GR, McCartney AL (2005) Differences between the gut microflora of children with autistic spectrum disorders and that of healthy children. *J Med Microbiol* 54:987-991.
65. Pisani V, Stefani A, Pierantozzi M, Natoli S, Stanzione P, Franciotta D, Pisani A (2012) Increased blood-cerebrospinal fluid transfer of albumin in advanced Parkinson's disease. *J Neuroinflammation* 9:188.
66. Redzic ZB, Preston JE, Duncan JA, Chodobski A, Szymdynger-Chodobska J (2005) The choroid plexus-cerebrospinal fluid system: from development to aging. *Curr Top Dev Biol* 71:1-52.

67. Song M, Xiong JX, Wang YY, Tang J, Zhang B, Bai Y (2012) VIP enhances phagocytosis of fibrillar beta-amyloid by microglia and attenuates amyloid deposition in the brain of APP/PS1 mice. *PLoS One* 7:e29790.
68. Stolp HB, Dziegielewska KM (2009) Review: Role of developmental inflammation and blood-brain barrier dysfunction in neurodevelopmental and neurodegenerative diseases. *Neuropathol Appl Neurobiol* 35:132-146.
69. Thouvenot E, Lafon-Cazal M, Demettré E, Jouin P, Bockaert J, Marin P (2006) The proteomic analysis of mouse choroid plexus secretome reveals a high protein secretion capacity of choroidal epithelial cells. *Proteomics* 6:5941-5952.
70. Van Hauwermeiren F, Vandenbroucke RE, Grine L, Lodens S, Van Wonterghem E, De Rycke R, De Geest N, Hassan B, Libert C (2015) TNFR1-induced lethal inflammation is mediated by goblet and Paneth cell dysfunction. *Mucosal Immunol* 8:828-840.
71. Vandenbroucke RE, Vanlaere I, Van Hauwermeiren F, Van Wonterghem E, Wilson C, Libert C (2014) Pro-inflammatory effects of matrix metalloproteinase 7 in acute inflammation. *Mucosal Immunol* 7:579-588.
72. Vandenbroucke RE, Dejonckheere E, Van Lint P, Demeestere D, Van Wonterghem E, Vanlaere I, Puimege L, Van Hauwermeiren F, De Rycke R, Mc Guire C, Campestre C, Lopez-Otin C, Matthys P, Leclercq G, Libert C (2012) Matrix metalloprotease 8-dependent extracellular matrix cleavage at the blood-CSF barrier contributes to lethality during systemic inflammatory diseases. *J Neurosci* 32:9805-9816.
73. Vawter MP, Dillon-Carter O, Tourtellotte WW, Carvey P, Freed WJ (1996) TGFbeta1 and TGFbeta2 concentrations are elevated in Parkinson's disease in ventricular cerebrospinal fluid. *Exp Neurol* 142:313-322.
74. Walters WA, Xu Z, Knight R (2014) Meta-analyses of human gut microbes associated with obesity and IBD. *FEBS Lett* 588:4223-4233.
75. Wirdefeldt K, Adami HO, Cole P, Trichopoulos D, Mandel J (2011) Epidemiology and etiology of Parkinson's disease: a review of the evidence. *Eur J Epidemiol* 26 Suppl 1:S1-58.
76. Wong F, Rayner-Hartley E, Byrne MF (2014) Extraintestinal manifestations of *Helicobacter pylori*: a concise review. *World J Gastroenterol* 20:11950-11961.
77. Wroblewski LE, Peek RM Jr (2011) "Targeted disruption of the epithelial-barrier by *Helicobacter pylori*". *Cell Commun Signal*. 9: 29
78. Yamashiro S (2006) Dellmann's Textbook of Veterinary Histology. *The Canadian Veterinary Journal* 48:414.
79. Zhang YJ, Li S, Gan RY, Zhou T, Xu DP, Li HB (2015) Impacts of gut bacteria on human health and diseases. *Int J Mol Sci* 16:7493-7519.

Tables

Table 1. Overview of used qRT-PCR primer sequences.

	Forward primer sequence	Reverse primer sequence
<i>H2afz</i>	GGTATCACCCCTCGTCACTT	TCAGCGATTTGTGGATGTGT
<i>Hprt</i> (stomach)	CAGGCCAGACTTTGTTGGAT	TTGCGCTCATCTTAGGCTTT
<i>Ppi</i>	AGCATACAGGTCTGGCATC	TTACACCTCCCAAAGACCAC
<i>Il18</i>	CACCTCACAAGCAGAGCACAAG	GCATTAGAAACAGTCCAGCCCATAC
<i>Kc</i>	GCTGGGATTCACCTCAAGAA	TCTCCGTTACTTGGGGACAC
<i>Mip2</i>	TGCCTGAAGACCCTGCCAAGG	GTTAGCCTTGCCTTTGTTTCTAG
<i>Lix</i>	CTCAGTCATAGCCGCAACCGAGC	CCGTTCTTTCCACTGCGAGTGC
<i>Ocln</i>	CCAGGCAGCGTGTCTCT	TTCTAAATAACAGTCACCTGAGGGC
<i>Cldn1</i>	TCTACGAGGGACTGTGGATG	TCAGATTGAGCAAGGAGTCG
<i>Cldn5</i>	GCAAGGTGTATGAATCTGTGCT	GTCAAGGTAACAAAGAGTGCCA
<i>Muc5ac</i>	Purchased from Qiagen (Qt01196006)	
<i>Muc5b</i>	CAGATCCATCCATCCCATTCT	TATCTGACTACCACTTGTGATGTGACT
<i>Muc6</i>	TGCTCCCAGAATGAGTACTTCGA	CAGAGGTGGAAGTGTGAACTCAGT
<i>Muc13</i>	GCCAGTCCTCCACACGGTA	CTGGGACCTGTGCTTCCACCG
<i>Vip</i>	AGTGTGCTGTTCTCTCAGTCG	GCCATTTTCTGCTAAGGGATTCT
<i>VipR1</i>	GATGTGGGACAACCTCACCTG	TAGCCGTGAATGGGGGAAAAC
<i>VipR2</i>	GACCTGCTACTGCTGGTTG	CAGCTCTGCACATTTTGTCTCT
<i>Tnf</i>	ACCCTGGTATGAGCCCATATAC	ACACCCATTCCCTTCACAGAG
<i>iNos</i>	CAGCTGGGCTGTACAAACCTT	CATTGGAAGTGAAGCGTTTCG
<i>Zo1</i>	AGGACACCAAAGCATGTGAG	GGCATTCTGCTGGTTACA
<i>Zo3</i>	ACCCTATGGCCTGGGCTTC	CCCGGGTACAACGTGTCC
<i>Hprt choroid plexus</i>	AGTGTGGATACAGGCCAGAC	CGTGATTCAAATCCCTGAAGT
<i>Rpl</i>	CCTGCTGCTCTCAAGGTT	TGGTTGTCACTGCCTGGTACTT
<i>Ubc</i>	AGGTCAAACAGGAAGACAGACGTA	TCACACCCAAGAACAAGCACA

<i>Gapdh</i>	TGAAGCAGGCATCTGAGGG	CGAAGGTGGAAGAGTGGGAG
<i>H. suis</i>	AAAACAMAGGCGATCGCCCTGTA	TTTCTTCGCCAGGTTCAAAGCG

Chapter 3

A novel isolation protocol and probe-based RT-PCR for diagnosis of gastric infections with the zoonotic pathogen *Helicobacter suis*

Blaecher Caroline¹, Bauwens Eva^{1*}, Tay Alfred^{2*}, Peters Fanny², Dobbs Sylvia^{3,4,5}, Dobbs John^{3,4,5}, Charlett
André^{3,6}, Ducatelle Richard¹, Haesebrouck Freddy^{1*}, Smet Annemieke^{1*#}

¹Department of Pathology, Bacteriology and Avian Diseases, Faculty of Veterinary Medicine, Ghent University,

²The Marshall Centre for Infectious Diseases Research and Training, University of Western Australia, Nedlands,
Perth, Western Australia, Australia,

³Institute of Pharmaceutical Science, King's College London, London, UK.,

⁴The Maudsley Hospital, London, UK.,

⁵Department of Gastroenterology, King's College Hospital, London, UK.,

⁶Statistics Unit, Centre for Infectious Disease Surveillance and Control, Public Health England, London, NW9
5EQ, UK.

* equal contribution.

Adapted from: *Helicobacter* 2016, doi: 10.1111/hel.12369

Abstract

Background: *Helicobacter suis* is a very fastidious micro-organism associated with gastritis, gastric ulcers and mucosa-associated lymphoid tissue (MALT) lymphoma in humans. *In vitro* isolation of this agent from human patients has so far been unsuccessful.

Materials and methods: A probe-based real-time (RT-) PCR for the rapid detection of *H. suis* in gastric biopsies was developed. Secondly, a mouse-passage-based protocol was optimized for isolation of low numbers of viable *H. suis* bacteria. Mice were inoculated with different numbers of viable *H. suis* (10^2 - 10^8) and kept for 4 weeks to allow multiplication of this pathogen.

Results: The probe-based real-time (RT)-PCR exhibited a high degree of diagnostic specificity and analytical sensitivity, high linear correlations (r^2 between 0.995 and 0.999) and high amplification efficiencies (> 90%) for *H. suis*. No cross-reactivity was detected with human, porcine, non-human primate and murine DNA nor with DNA from other bacteria including *Helicobacter* spp. and *Campylobacter* spp.

H. suis was successfully re-isolated from the stomach of mice inoculated with at least 10^4 viable *H. suis*, using a biphasic medium (pH 5), consisting of *Brucella* agar with *Brucella* broth on top, both supplemented with vitox supplement, Campylobacter-selective supplement, amphotericin (5 µg/ml), HCl (0.05 %), fetal bovine serum (20 %) and linezolid (5 µg/ml). Linezolid was necessary to inhibit proliferation of contaminants, including lactobacilli.

Conclusion: The methods described in this paper can be implemented for detection or isolation of *H. suis* from human gastric biopsies.

Introduction

Helicobacter pylori (*H. pylori*) is associated with a wide range of gastric disorders, including gastritis, ulcers and cancer (Montecucco and Rappuoli, 2001). However, gastric disease in humans has also been associated with other *Helicobacter* species. These are collectively called non-*H. pylori Helicobacter* spp (NHPH). They are long, spiral-shaped organisms most commonly colonizing the stomachs of domestic animals (Haesebrouck et al., 2009). Reported estimates of prevalence of NHPH in human populations range from 0.2 to 6 % (Haesebrouck et al., 2009). However, due to difficulties in the diagnosis of NHPH infections these estimates are certainly biased and most probably underestimate the true prevalence. Moreover, it cannot be excluded that infections with these agents sometimes remain asymptomatic or cause only mild symptoms which are not rigorously examined. The prevalence of *H. pylori* in the Western world is decreasing, leaving a niche for infection with NHPH (Haesebrouck et al., 2009; Overby et al., 2016).

H. suis appears to be the most important amongst the NHPH. Not only does it represent the most common of the NHPH detected in the stomach of people suffering from gastric disorders, but it has also been detected in 27 % of Parkinson's disease (PD) patients and only in 2 % of controls from gastroenterology-services using an *ureA*-based SYBR green RT-PCR assay on gastric biopsies with positivity on amplicon sequencing (Blaecher et al., 2013).

Pigs are the main reservoir of *H. suis* (Haesebrouck et al., 2009; De Bruyne et al., 2012), but *H. suis* has also been found in the stomach of rhesus and crab-eating macaques (O'Rourke et al., 2004; Martin et al., 2013; Bosschem et al., 2016). *H. suis* is transmitted to humans most likely through direct or indirect contact with pigs (Joosten et al., 2013). Additionally, this bacterium can be present and persist in minced pork, suggesting that raw or undercooked pork may also constitute a source of *H. suis* infections in humans (De Cooman et al., 2013). *H. suis* has been associated with gastritis, peptic and duodenal ulcers, and low-grade mucosa-associated lymphoid tissue (MALT) lymphoma in humans. Compared to *H. pylori*, gastritis caused by *H. suis* appears less active and less severe, but the risk of developing MALT lymphoma is higher (Nakamura et al., 2007; Nakamura et al., 2008; Matsui et al., 2014).

The diagnosis of gastric *Helicobacter* infections is extremely difficult and usually involves endoscopic sampling of gastric biopsies. Commercial non-invasive tests to specifically diagnose infections with *H. suis* are currently unavailable. Attempts to isolate *H. suis* from human gastric biopsies, either directly, or after mouse-passage (Matsui et al., 2014) have been unsuccessful so far. Direct isolation from biopsies may be hampered by the low colonization density of this agent (Blaecher et al., 2013), whilst isolation after mouse-passage has been hampered by overgrowth of *Lactobacillus* spp. (Matsui et al., 2014; Aiba et al., 2015). Analysis of the gastric biopsies by molecular methods currently appears to be the only feasible way to identify infections with *H. suis*. One available molecular method makes use of RT-PCR based on the *ureA* gene for identification and quantification of *H. suis* bacteria (Blaecher et al., 2013; Flahou et al., 2010). In this assay, additional sequencing of the positive PCR amplicons is necessary, since cross-reactivity with host DNA and DNA from other gastric NHPH may occur (unpublished results). Recently, a new *H. suis*-specific PCR has been developed, using primer pairs targeting the *carR* gene. Compared to *ureA*, this *carR* gene is a good candidate for the identification of *H. suis*, since it is not present in other gastric *Helicobacter* genomes (Matsui et al., 2013). However, this novel

assay required further investigation, as it is not known whether the *carR*-specific primers react with other bacterial or host DNA.

The purpose of the present study is to further improve diagnostic tests for the detection of *H. suis*. We first developed and validated a *carR* probe-based RT-PCR that allows detection and quantification of *H. suis* in gastric tissue samples. Secondly, a mouse-passage-based protocol was further optimized for the isolation of low numbers of *H. suis* bacteria. Since mouse-passage of *H. suis* might influence strain characteristics, we also sequenced the genome of a porcine *H. suis* strain before and after passage in mice.

Materials and methods

1) *CarR* probe-based RT-PCR

Bacterial strains and *in vitro* culture conditions

H. suis, *H. heilmannii*, *H. ailurogastricus* and *Lactobacillus* spp. (*L. reuteri* and *L. gasseri*) (Table 1) were cultivated biphasically on *Brucella* agar (Immunosource, Halle-Zoersel, Belgium) plates, supplemented with 1 ml Skirrow (Oxoid, Hampshire, United Kingdom), 3 ml Vitox (Oxoid), 5 µg/ml amphotericin B (Sigma-Aldrich, Steinheim, Germany), 0.05 % HCL (Sigma-Aldrich) and 20 % inactivated fetal bovine serum (Perbio, Thermo Scientific Hyclone, Erembodegem; Belgium) (*H. suis* growth medium) (Baele et al., 2008). *H. bizzozeronii*, *H. felis*, *H. salomonis*, *H. pylori*, *H. cynogastricus*, *H. baculiformis*, *H. cetorum*, *H. cholecystus*, *H. trogonum*, *H. acinonychis*, *H. mustellae*, *H. pullorum*, *H. marmotae*, *H. mesocricetorum*, *H. bilis*, *H. equorum*, *H. canadensis* and *Campylobacter* spp. (*C. jejuni*, *C. coli*, *C. fetus* and *C. upsaliensis*) (Table 1) were grown on brain heart infusion (BHI) agar (Bio-rad, Nazareth, Belgium) plates supplemented with 10 % horse blood (E&O Laboratories, Bonnybridge, UK), Skirrow, Vitox, and 5 µg/ml amphotericin B. All *Helicobacter* spp. and *Campylobacter* spp. were cultivated under micro-aerobic conditions (10 % CO₂, 5 % O₂ and 85 % N₂; 37°C). *Staphylococcus aureus*, *Staphylococcus pseudintermedius*, *Streptococcus suis*, *Streptococcus equi* subsp. *equi*, *Streptococcus equi* subsp. *zooepidemicus*, *Pasteurella multocida*, *Mannheimia haemolytica*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Proteus vulgaris*, *Bacillus licheniformis*, *Salmonella* Typhimurium, *Trueperella pyogenes*, *Escherichia coli*, *Aeromonas hydrophila*, *Enterococcus faecalis* and *Actinobacillus equuli* were grown aerobically on commercially available Columbia agar plates supplemented with 5 % sheep blood (Oxoid) at 37°C. *Clostridium perfringens* and *Fusobacterium gastrosuis* (Table 1), grown on Columbia agar plates (with 5% sheep blood), were incubated anaerobically at 37°C, whereas *Yersinia pseudotuberculosis* was grown on Columbia agar plates (with 5 % sheep blood) and incubated aerobically at 30°C (Table 1).

Primer and probe design

Analysis and comparison of the *carR*-gene sequence from *H. suis* strains HS1 (Genbank accession no. ADGY000000000), HS5 (Genbank accession no. HDAO000000000), TKY (EMBL/Genbank/DDBJ database accession no. AB849014), SNTW101 (EMBL/Genbank/DDBJ database accession no. AB849015), HS8 (EMBL/Genbank/DDBJ database accession no. AB849016) and HS10 (EMBL/Genbank/DDBJ database accession no. AB849017) resulted in the selection of a specific 149 bp region. Based on this region (149 bp), primers and a probe (20 bp) were developed: sense primer *carR*-FQ1 (5'-TGT GGC TAG RTG TTT GCC -3'), antisense primer

carR-RQ1 (5'-CT GCA GGC AAY GCT GTT GTT -3') and *carR* probe (5'- ACC ACG CCT GCA GGC AAC AA-3'). The melting temperature and GC% of the primers and probe were calculated and optimized using the OligoAnalyser3.1 (Integrated DNA technologies, Leuven, Belgium). Subsequently, the primers and probe were verified using BLASTN to ensure that amplification of genes from other organisms or species was unlikely. Finally, the probe was combined with the dye 6-carboxyfluorescein (FAM) and the quencher ZEN/3IABkFQ.

Bacterial DNA extraction and quantitation of standards

DNA was extracted from the different bacterial species using the PrepmanTM reagent (Life Technologies, Ledeberg, Belgium) according to the manufacturer's instructions. For generation of the standard for the RT-PCR assay, part of the *carR* gene (423bp) from *H. suis* HS5 was amplified using primers cbcAR-fq (5'- ATG TAC AAG SGC TCT TGA GGG -3') and cbcAR-rq (5'- GAT GGC TGC TTG GAA ACT AGC -3') with an annealing temperature of 57°C. Subsequently, the DNA concentration of the PCR products was measured and diluted to a concentration of 10¹⁰ Genomic equivalents (GE). The standard consisted of 10-fold-dilution series starting at 10⁹ GE for each 10 µl of reaction mixture.

***H. suis* ureA-based SYBR green RT-PCR**

The *H. suis* SYBR green assay based on the amplification of part of the gene encoding the urease A subunit was carried out as described before (Blaecher et al., 2013).

***H. suis* carR SYBR green RT-PCR**

The *H. suis* *carR* RT-PCR was first evaluated using SYBR green. Two µl of DNA template (samples and standard) was suspended in a 10 µl reaction mixture consisting of 3.5 µl HPLC water, 6 µl SensiMixTM SYBR No-ROX (Bioline Reagents Ltd, London, UK) and 0.25 µl of primers *carR*-FQ1 and *carR*-RQ1 located within the 423 bp fragment to yield a 149 bp PCR product. The amplification conditions consisted of 95°C for 15 min followed by 47 cycles of denaturation (30 seconds at 95°C), annealing (30 sec at 56°C) and elongation (30 sec at 72°C). The final elongation step took place at a temperature of 95°C for 15 min. Melting curve data were obtained starting at 65°C to 95°C with a velocity of 0.5°C per 5 sec. The standard and bacterial DNA samples were run in duplicate on a CFX384TM RT-PCR System or a CFX96TM RT-PCR System with a C1000 Thermal Cycler (Bio-Rad).

***H. suis* carR probe-based RT-PCR and optimization**

Assays were performed on a CFX96 real-time system (Bio-rad). Primer concentrations were determined in a checkerboard system with a standard probe concentration of 250 nM. Subsequently, the probe concentration was optimized with the previously determined optimal primer concentrations (Bloo et al., 2013). The reaction mixture of 12 µl consisted of 6 µl of IQTM Supermix (Bio-rad), 3.25 µl of HPLC water (Merck Millipore, Overijse, Belgium), 0.3 µl of both primers (*carR*-FQ and *carR*-RQ), 0.15 µl of the probe and 2 µl DNA template (standards and samples). The amplification conditions consisted of 95°C for 10 min followed by 47 cycles of denaturation (20 sec at 95°C), annealing (30 sec at 62.5°C) and elongation (30 sec at 73°C). The final elongation step took place at a temperature of 95°C for 15 min. Both standards and samples were run in duplicate. The precision, i.e.

the reproducibility and repeatability of the developed *H. suis* *carR* probe-based RT-PCR was evaluated by determining inter- and intra-assay variability, both expressed by the standard deviation (SD) (Bustin et al., 2009). Intrarun variability was ascertained by comparing the obtained cq-values for the standard ten-fold dilution series (starting from 10^9 GE) within one RT-PCR run, whereas interr run variability was evaluated by comparing those values between 3 RT-PCR runs. The diagnostic specificity of the *carR*-based SYBR green RT-PCR, the *carR* probe-based RT-PCR and the *ureA*-based SYBR green RT-PCR was evaluated by assaying DNA extracts of different *Helicobacter* spp., *Campylobacter* spp. and other bacteria (Table 1). The amplification efficiency, slope and r^2 were calculated by linear regression analysis using the Bio-Rad CFX Manager v1.6 software (Bio-Rad, Hercules, CA,USA).

Gastric tissue samples for validation of the *carR* probe-based RT-PCR

To validate the use of the *carR* probe-based RT-PCR to detect *H. suis* in gastric tissue samples and exclude reactivity with host DNA, we applied the optimized protocol to analyze 80 human biopsies in which *H. suis* infection was or was not detected by *ureA*-based SYBR green RT-PCR with positivity on amplicon sequencing (Blaecher et al., 2013; Table 2), 9 gastric samples from pigs with or without known natural *H. suis* infection (Table 2), 9 gastric samples from non-human primates with or without known natural *H. suis* infection (Bosschem et al., 2016; Table 2) and 8 gastric samples from mice experimentally inoculated with *H. suis* or with the growth medium of the bacterium (Table 2). The human gastric biopsies originated from a previous experiment with approval from the ethics committee of Ghent University (EC UZG 2011/393) and from the diagnostic service of the Gastrointestinal Reference Unit, Public Health England. The non-human primate gastric samples originated from rhesus macaques and crab-eating macaques, residing at the Biomedical Primate Research Centre (Rijswijk, The Netherlands), accredited by the association for Assessment and Accreditation of Laboratory Animal Care (EU Directive 63/2010), and which were euthanized for reasons unrelated to the study. The stomach samples from control mice and mice experimentally infected with *H. suis* originated from an experiment with approval from the ethics committee of the faculty of Veterinary Medicine, Ghent University (approval no. EC 2014/73). The gastric tissue samples from pigs, used in the study by De Witte and colleagues (De Witte et al., 2016), were included in the analyses as well.

DNA from all gastric tissue samples was extracted using the Qiagen Blood and Tissue kit (Qiagen, Venlo, The Netherlands), according to the manufacturer's instructions.

2) Optimization of a mouse-passage-based isolation protocol for *Helicobacter suis*

Mice and experimental procedure

Six-week-old, female specific pathogen free (SPF) C57BL/6JolaHsd mice were purchased from Harlan NL (Horst, The Netherlands). The animals were housed in individual filter top cages, had free access to water and food (an autoclaved commercial diet: TEKLAD 2018S, containing 18 % protein; Harlan) throughout the experiment and were monitored daily. The experimental protocol was approved by the ethics committee of the faculty of Veterinary Medicine, Ghent university, Belgium (approval no. EC 2012/58).

H. suis HS1, grown biphasically as described above, was used as inoculum. After incubation under micro-aerobic conditions, the bacteria were harvested. Ten-fold dilution series were prepared in *Brucella* broth. The final concentrations ranged from 10^8 to 10^2 viable *H. suis* bacteria/ml. For each bacterial concentration, 2 mice were anaesthetized on two successive days with 2.5% isoflurane and then intragastrically inoculated with 300 μ l using a feeding canula. At 4 weeks post-inoculation, animals were euthanized by cervical dislocation under deep isoflurane anaesthesia (5%). The stomach of each mouse was resected for isolation purposes.

Isolation of porcine *H. suis* strain from the murine stomach after mouse-passage

The *H. suis* isolation protocol, which has successfully been applied on porcine and non-human primate stomachs highly colonized with *H. suis* (Baele et al., 2008; Bosschem et al., 2016), was further optimized for use on murine gastric tissue with low numbers of *H. suis* bacteria. We found that supplementation with linezolid at 5 μ g/ml to both phases of the *H. suis* growth medium did not inhibit *H. suis* cultivation, whereas the growth of contaminants present in the murine stomach was prevented (unpublished results).

This 5 μ g/ml linezolid containing selective medium was evaluated in the experimentally infected mice. The murine stomachs were opened via the greater curvature. The gastric content was removed and the stomachs were washed and submerged in a 1 % HCl solution for 45 min. Each stomach was homogenized and half of it was inoculated onto the linezolid-supplemented *H. suis* growth medium, whereas the other half was inoculated onto the *H. suis* growth medium without linezolid supplementation (control sample). Subsequently, approximately 150 μ l *Brucella* broth with or without (control sample) supplementation of 5 μ g/ml linezolid was added on top of the inoculum. The plates were incubated at 37°C under micro-aerobic conditions, daily checked for growth and humidified with the linezolid-supplemented *Brucella* broth or the *Brucella* broth without linezolid (control samples). Finally, pure isolates were transferred to *H. suis* growth medium without linezolid and stored at -70°C until further use.

Whole genome sequencing of *H. suis* isolates obtained after mouse-passage

The genomic DNA of 4 isolates obtained after mouse-passage of the porcine *H. suis* HS1 strain (isolate HS1_output (O)1, isolate HS1_O2, isolate HS1_O3 and isolate HS1_O4, originating from 4 different mice) and the original porcine HS1_input (I)1 isolate, retrieved from the pig's stomach and used for mouse-passage, was extracted using the Qiagen Blood and Tissue kit (Qiagen) according to the manufacturer's guidelines. Whole genome sequencing was performed as described before (Joosten et al., 2015). Briefly, 1 ng genomic DNA was used for library generation. Sequencing libraries were then prepared using Nextera XT chemistry (Illumina Inc., San Diego, CA, USA) in accordance with the manufacturer's recommendations. Subsequently, libraries were sequenced for a 250-300 bp paired-end sequencing run using the Miseq personal sequencer (Illumina). All genomes were assembled with the CLC Genomics Workbench, version 7 and annotated using the RAST server (Aziz et al., 2008).

Identification of recombination events in the genomes of *H. suis* isolates obtained after mouse-passage

Homologous recombination in 4 *H. suis* isolates obtained after mouse-passage of the porcine *H. suis* strain HS1 was assessed using an iterative application of ClonalFrameML as follows (Didelot & Wilson, 2015). From the obtained genome assemblies, a core genome alignment was first created using the genome profiler (GeP) software tool (Zhang et al., 2015). Subsequently, a maximum-likelihood input phylogeny was inferred using RaxML v7.0.43 (Stamakis, 2006) under the General Time-Reversible model. Both the RaxML tree and the core genome alignment were then analyzed by ClonalFrameML. This assay was rerun several times until the number of recombination events and phylogeny remained stable. Estimates of recombination parameters and the set of recombination events were recorded from the final run.

Nucleotide sequence accession numbers

The genome sequences determined in this study have been deposited in the EMBL database under BioProject record numbers: PRJEB15615, PRJEB15616, PRJEB15617, PRJEB15618, PRJEB15619 and received the following accession numbers: FMSU01000000, FMSQ01000000, FMSS01000000, FMST01000000 and FMSR01000000.

RESULTS AND DISCUSSION

1) *CarR* probe-based RT-PCR

CarR probe-based RT-PCR assay optimization

The primer and probe concentrations used for this assay were the lowest concentrations that yielded the lowest quantitation cycle (cq) value and the highest ΔR_n . This latter is the R_n value (the fluorescent emission of the reporter dye, normalized to the background fluorescence) of the complete reaction mixture (including template) minus the R_n value of a negative control (Bloo et al., 2013). This resulted in a final reaction mixture of 12 μ l composed of 6 μ l of IQTM Supermix (Bio-rad, Belgium), 3.25 μ l of HPLC water (Merck Millipore, Belgium), *H. suis*-specific *carR* forward primer *carR*-FQ1 at a concentration of 0.5 μ M, *H. suis*-specific *carR* reverse primer *carR*-RQ1 at a concentration of 0.5 μ M, *H. suis*-specific *carR* FAM-labeled probe at a concentration of 0.25 μ M and 2 μ l template. By application of a temperature gradient for analysis of the standard curve, the optimal annealing temperature for the combination of *carR*-FQ1, *carR*-RQ1 and the *carR* probe was set at 62.5°C, leading to a run amplification efficiency between 90-100%, the square of the linear correlation (r^2) between 0.995 and 0.999 and a slope of -3.53 (Figure 1). With the exception of a change in the annealing temperature, the amplification conditions of the *carR* probe-based RT-PCR were identical to the conditions used in the *H. suis ureA*-based SYBR green RT-PCR.

In 2004, Schabereiter-Gurtner and co-workers (Schabereiter-Gurtner et al., 2004) developed an *ureA* probe-based RT-PCR and a 23S rRNA probe-based RT-PCR for the detection of *H. pylori* in stool samples. In contrast to our developed *carR* probe-based RT-PCR assay, no information on the linear correlation of the *ureA*/23S rRNA probe-based RT-PCR assays for the detection of *H. pylori* was provided in this study. However for these latter assays, a slope of -3.7 was reported for both standard curves, consisting of tenfold dilution series. Using the formula, $E=10^{-(1/\text{slope})}-1$, an efficiency of 86 % could be calculated for both the *ureA* and 23S rRNA probe-

based RT-PCR assays. Since the *carR* probe-based RT-PCR in our study resulted in a slope of -3.53 and the optimal slope for a tenfold dilution series is considered -3.32 (Real-time PCR application guide, Bio-Rad Laboratories, 2006), the *carR* probe-based RT-PCR is characterized by a higher amplification efficiency compared to the *ureA*/23S rRNA probe-based RT-PCRs. As amplification efficiency is a measure for the percentage of DNA template amplified in each cycle and a slope of -3.32 for a tenfold dilution series equals an amplification efficiency of 100% (Real-time PCR application guide, Bio-Rad Laboratories, 2006), the amplification of DNA template is more efficient in the *carR* probe-based RT-PCR assay for the detection of *H. suis* compared to the *H. pylori*-specific *ureA*/23S rRNA probe-based RT-PCR assays.

Figure 1




Figure 1: Standard curve along with efficiency and linear correlation of *carR* probe-based RT-PCR. Standards were run in duplicate. The *carR* probe-based RT-PCR is characterized by a high efficiency (>90 %) and a linear correlation (r^2) between 0.995 and 0.999.

Analytical sensitivity and diagnostic specificity

The analytical sensitivity of the *carR* probe-based RT-PCR was tested with the described quantitation standard of *H. suis*. Triplicates of serial dilutions ranging from 10^0 to 10^9 GE were assayed. In all replicates, cq-values were generated, suggesting that the limit of detection (LOD) of *H. suis* DNA using the *carR* probe-based RT-PCR assay is 1 GE per PCR. In theory, a DNA dilution of 10^0 will contain 1 GE per PCR. However, Bustin (Bustin et al., 2009) highlighted that the most sensitive LOD is 3 GE per PCR assuming a Poisson distribution, a 95% chance of including at least 1 GE per PCR and 1GE detection (Bustin et al., 2009). This means that in case of a dilution of 10^0 GE, 37 % of the aliquots will contain no GE, another 37% will contain 1 GE and the final 26 % will contain 2 or more GE (Bustin et al., 2009). A similar detection limit has been reported for the *H. suis*-specific *ureA*-based SYBR green RT-PCR (Blaecher et al., 2013; Bosschem et al., 2015).

To verify the diagnostic specificity of the *carR* probe-based RT-PCR, DNA samples from several bacteria (Table 1), including different *Helicobacter* spp., *Campylobacter* spp. and members of the *Enterobacteriaceae* family (Schabereiter-Gurtner et al., 2004; Moyaert et al., 2008; Matsui et al., 2014), were analyzed. Of all 77 different bacterial non-*H. suis* samples and in contrast to the *ureA*-based SYBR green RT-PCR (Table 1), the *carR* probe-based RT-PCR amplified only *H. suis* DNA and not the DNA from the other 77 bacterial non-*H. suis* samples (95 % confidence interval: 95%-100%), indicating a high degree of diagnostic specificity. No aspecific reactions with DNA from gastric and enterohepatic *Helicobacter* spp., *Campylobacter* spp. and other bacteria were observed (Table 1). Identical results were obtained using the *carR*-based SYBR green RT-PCR, highlighting the diagnostic specificity of the *carR* primers for *H. suis*.

Assay performance and precision

To determine the assay performance and precision of the *carR* probe-based RT-PCR with the described quantitation standards of *H. suis*, intrarun and interrune variability was defined as described before (Blooi et al., 2017). Both inter- and intrarun assays of the *carR* probe-based RT-PCR were characterized by high squared linear correlation ($r^2 > 0.995$) and amplification efficiency ($\geq 90\%$) values, together with low intra- and inter-assay variabilities ($SD < 1$; Figures 2 and 3). The *cq*-values for the dilution series of *H. suis* bacteria in a run and between 3 different runs were similar (Figure 2 and 3). Within a run the highest variation was observed at 10^9 GE, resulting in a standard deviation of 0.97 (Supplementary Table 1), whereas between different runs the highest variation was observed at 10^0 and 10^9 GE, resulting in a standard deviation of 0.76 and 0.62, respectively (Supplementary Table 2). The separate *cq*-values of each inter- and intrarun assay are presented in Supplementary Table 1 and 2. The above results demonstrate that the probe-based RT-PCR has a high reproducibility as well as a good performance over the tested quantitation range.

Figure 2




Figure 2: Overview of the *cq*-values of the standard dilution series in 1 run. The *cq*-values for the dilution series of *H. suis* bacteria within 1 run were similar. Inside 1 run the highest variation was observed at 10^9 GE, resulting in a standard deviation of 0.97. The separate *cq*-values of one run are demonstrated in Supplementary Table 2.


Figure 3

Figure 3: Overview of the mean cq-values of the standard dilution series between 3 different runs. Between 3 different runs, the cq-values were likewise the same as in 1 run. At the dilution of 10^9 and 10^0 GE, the highest standard deviation of the cq-values between the 3 runs amounted 0.76 and 0.62, respectively. The separate cq-values of the 3 different runs are demonstrated in Supplementary Table 3.

Validation on gastric tissue samples

To validate the developed *H. suis*-specific *carR* probe-based RT-PCR, DNA from human, non-human primate, murine and porcine gastric tissue samples was assayed (Table 2 and Supplementary Table 3). All tested samples were run in the *H. suis*-specific *ureA*-based SYBR green, the *carR*-based SYBR green and the *carR* probe-based RT-PCR assay to compare variability in cq-values between the assays. Out of the 80 human gastric biopsy samples, the *ureA*-based SYBR green RT-PCR identified 7 positive human gastric samples. Six of these also reacted positive using the *carR* probe-based RT-PCR. None of the human gastric samples tested positive for *H. suis* DNA using the *carR*-based SYBR green RT-PCR (Table 2 and Supplementary Table 3). The number of *H. suis* positive non-human primate samples identified by the *ureA*-based SYBR green RT-PCR and the *carR* probe-based RT-PCR were 9 and 8, respectively, whereas only 6 of the 9 samples tested positive for *H. suis* DNA using the *carR*-based SYBR green RT-PCR (Table 2 and Supplementary Table 3). Finally, for both murine and porcine gastric tissue samples, all known negative and positive *H. suis* samples were properly identified using both *carR* probe-based RT-PCR and *ureA*-based SYBR green RT-PCR. However, none of the known positive murine and porcine gastric samples were identified using the *carR*-based SYBR green RT-PCR (Table 2 and Supplementary Table 3).

Additional sequencing was performed for the samples reacting positive in *ureA*-based SYBR green RT-PCR. One human and one non-human primate sample, which tested negative for *H. suis* using the *carR* probe-based RT-PCR, were false positive in the *ureA*-based SYBR green RT-PCR (Table 2 and Supplementary Table 3). The sequence of the amplicon obtained within the human gastric sample corresponded to that of human DNA. That

of the non-human primate gastric sample corresponded to the *ureA/B* gene sequence of an unknown *Helicobacter* sp.

In contrast, none of the human, porcine and murine samples tested positive for the presence of *H. suis* DNA using the *carR*-based SYBR green RT-PCR. Although the melting curve obtained for these human, porcine and murine samples did not correspond with that of *H. suis*, high *cq*-values were detected for these samples using the *carR*-based SYBR green RT-PCR. Sequencing of the amplicons obtained in the SYBR green RT-PCR with *carR*-based primers revealed that human, porcine, murine and primate DNA was amplified, indicating that these primers do not only bind to the *H. suis carR* gene, but also to host DNA which is present in large quantities in gastric samples. This renders the *carR*-based SYBR green RT-PCR detection of amplicons inefficient and a method that specifically detects DNA sequences amplified from the *H. suis carR* gene is required. In the present study, this was realized using a probe that detects a specific part of the *carR* gene. Indeed, the SYBR green RT-PCR assay emits a fluorescent signal when the fluorophore binds to any amplified double stranded DNA. In contrast, the probe binds on a specific region within the amplicon of single stranded DNA. The emission of fluorescence only takes place when the reporter dye is separated from the quencher by the DNA polymerase. Since the true positive samples, derived from the human, non-human primate, murine and porcine stomachs, tested false negative using the *carR*-based SYBR green RT-PCR assay, they are not reliable and therefore not included in the evaluation of the variability between the different assays. Compared to the results from the *ureA*-based SYBR green RT-PCR, the *cq*-values from the *H. suis* positive samples obtained by the *carR* probe-based RT-PCR were less variable resulting in a lower *cq*-mean and standard deviation (Table 3 and Supplementary Table 3). This finding clearly indicates a higher diagnostic specificity and analytical sensitivity for the *carR* probe-based RT-PCR.

In our study, no aspecific reactions with human, non-human primate, murine and porcine DNA nor with DNA from other bacteria could be observed when using the *carR* probe-based RT-PCR. As only relatively low amounts of *H. suis* DNA have been detected in human gastric biopsies using RT-PCR and large amounts of host DNA are present in these samples, the use of the target-specific *carR* probe may be very useful to detect *H. suis* DNA in human gastric biopsies.

2) Optimization of a mouse-passage-based isolation protocol for *Helicobacter suis*

Optimization of a mouse-passage *H. suis* isolation method

Overgrowth with *Lactobacillus* spp. has been reported as one of the main reasons for failure of *H. suis* isolation from the murine and human stomach (Matsui et al., 2014; unpublished data). In order to inhibit the growth of contaminants, including *Lactobacillus* spp., and allow cultivation of *H. suis*, linezolid was added to the biphasic *H. suis* isolation medium. Linezolid is a synthetic oxazolidinone, inhibiting the protein synthesis by binding to the 50S ribosomal subunit of the peptidyltransferase center of the bacterial ribosomes. It is active against a broad spectrum of Gram-positive bacteria, including many *Lactobacillus* species (Agar and Gould, 2012), and has a relatively low activity (MIC₉₀ 8 µg/ml) against *H. pylori* (Sanchez et al., 2000).

The linezolid-supplemented biphasic growth medium and the *H. suis* growth medium without linezolid (control sample) were inoculated with homogenized gastric tissue samples from mice, infected for 4 weeks with a

dilution series of viable *H. suis*. All plates with growth medium that did not contain linezolid, were overgrown by contaminants, making isolation of *H. suis* impossible. On linezolid-supplemented biphasic growth medium inoculated with stomach homogenate of mice infected with 10^8 , 10^7 , 10^6 , 10^5 and 10^4 bacteria, pure *H. suis* isolates were obtained. *H. suis* bacteria could be observed after 2-3 days of *in vitro* cultivation. On the contrary, after two weeks of incubation, attempts to isolate *H. suis* from the stomach of mice infected with fewer than 10^4 viable bacteria remained unsuccessful, indicating that the minimal infection dose of *H. suis* in mice is at least 10^4 viable *H. suis* bacteria/ml. This finding has also been confirmed in other reports (Matsumoto et al., 2009; Matsui et al., 2014).

Low recombination events and maintenance of integrity in the *H. suis* genome

Whole genome analysis of the 4 isolates obtained after mouse-passage of porcine *H. suis* HS1 strain (HS1_O1, HS1_O2, HS1_O3 and HS1_O4) and the original porcine HS1_I1 isolate, showed that mouse-passage of *H. suis* did not result in major recombination events within the *H. suis* genome (Figure 4). Similar findings were reported for *H. pylori* after mouse-passage (Lundin et al., 2005). This may indicate that *H. suis* strains obtained using this mouse-passage-based isolation protocol can be used to further characterize the agent.

Figure 4




Figure 4: Phylogenetic tree in combination with an overview of the recombination events. Shown are the different recombination events in combination with their position detected after whole genome comparison of the different *H. suis* strains, successively HS2, HS1_I1, HS1_o4, HS1_O3, HS1_O1, HS1_O2, HS7, HS3, HS5, HS10, HS6, HS4, HS8, HS9.

Conclusions

The developed *carR* probe-based RT-PCR appears to be a reliable and accurate diagnostic method that can be applied for the detection and quantitation of *H. suis* in gastric samples. Since no interactions with other NHPH, nor with host DNA could be observed, the use of this technique is recommended for the screening of gastric biopsies, derived from human patients and animals. Using the optimized isolation protocol, we succeeded in isolating relatively low numbers of *H. suis* bacteria from the mouse stomach. Application of this technique may help to obtain and characterize pure human-derived *H. suis* isolates after mouse-passage of *H. suis*-infected human gastric biopsies.

Acknowledgements

This work was supported by the Research Fund of Ghent University, Belgium, Code GOA 01G00408 and by the Research Foundation Flanders (FWO Vlaanderen). The authors would like to thank Nathalie Van Rysselberghe and Sofie De Bruyckere for their excellent technical assistance.

References:

1. Ager S and Gould K. 2012. Clinical update on linezolid in the treatment of Gram-positive bacterial infections. *Infection and drug resistance* 5: 87-102.
2. Aiba Y, Nakano Y, Koga Y, Takahashi K, Komatsu Y. 2015. A highly acid-resistant novel strain of *Lactobacillus johnsonii* NO.1088 has antibacterial activity, including that against *Helicobacter pylori*, and inhibits gastrin-mediated acid production in mice. *MicrobiologyOpen* 4(3):465-474.
3. Aziz RK, Bartels D, Aaron A Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST Server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75
4. Baele M, Decostere A, Vandamme P, Ceelen L, Hellemans A, Mast J, Chiers K, Ducatelle R, Haesebrouck F. 2008. Isolation and characterization of *Helicobacter suis* sp.nov. from pig stomach. *Int J Syst Evol Microbiol*. 58(Pt 6):1350-8.
5. Blaecher C, Smet A, Flahou B, Pasmans F, Ducatelle R, Taylor D, Weller C, Bjarnason I, Charlett A, Lawson AJ, Dobbs RJ, Dobbs SM, Haesebrouck F. 2013. Significantly high frequency of *Helicobacter suis* in patients with idiopathic parkinsonism in contrast to control patients. *Alim.Pharm and Therap*. 38(11-12): 1347–1353
6. Blooi M, Pasmans F, Longcore JE, Spitzen-van der Sluijs A, Vercammen F, Martel A. 2013. Duplex Real-Time PCR for Rapid Simultaneous Detection of *Batrachochytrium dendrobatidis* and *Batrachochytrium salamandrivorans* in Amphibian Samples. *J. Clin. Microbiol*. 51(12): 4173-4177
7. Bosschem I, Bayry J, De Bruyne E, Van Deun K, Smet A, Vercauteren G, Ducatelle R, Haesebrouck F, Flahou B. 2015. Effect of Different Adjuvants on Protection and Side-Effects Induced by *Helicobacter suis* Whole-Cell Lysate Vaccination. *PLoS ONE* 10(6): e0131364
8. Bosschem I, Flahou B, Bakker J, Heuvelman E, Langermans JAM, De Bruyne E, Joosten M, Smet A, Ducatelle R, Haesebrouck F. 2016. Comparative virulence of *in vitro* cultures primate- and pig-associated *Helicobacter suis* strains in a BALB/c mouse and Mongolian gerbil model. *Helicobacter*, in press
9. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. 2009. The MIQE Guidelines: Minimum information for publication of quantitative Real-Time PCR experiments. *Clin Chem* 55: 4
10. De Bruyne E, Flahou F, Chiers K, Meyns T, Kumar S, Vermoote M, Pasmans F, Millet S, Dewulf J, Haesebrouck F. 2012. An experimental *Helicobacter suis* infection causes gastritis and reduced daily weight gain in pigs. *Veterinary Microbiology* . 160(3-4). p.449-454
11. De Cooman L, Flahou B, Houf K, Smet A, Ducatelle R, Pasmans F, Haesebrouck F. 2013. Survival of *Helicobacter suis* bacteria in retail pig meat. *Int. J. Food Microbiol*. 166:164-167
12. De Witte C, Ducatelle R, De Bruyne E, Smet A, Haesebrouck F, Flahou B. 2015. Detection, isolation and characterization of a highly prevalent new *Fusobacterium* sp. colonizing the stomach of pigs. *Proceeding De Witte EHSG congress Cyprus*. *Helicobacter* 20 (S1): abstract noW3.1s
13. Didelot X and Wilson DJ. 2015. ClonalFrameML: Efficient Inference of Recombination in Whole Bacterial Genomes. *PLoS Comput Biol* 11(2): e1004041
14. Flahou B, Haesebrouck F, Pasmans F, D'Herde K, Driessen A, Van Deun K, Smet A, Duchateau L, Chiers K, Ducatelle R. 2010. *Helicobacter suis* Causes Severe Gastric Pathology in Mouse and Mongolian Gerbil Models of Human Gastric Disease. *PLoS ONE* 5(11): e14083.
15. Haesebrouck F, Pasmans F, Flahou B, Chiers K, Baele M, Meyns T, Decostere A, Ducatelle R. 2009. Gastric helicobacters in domesticated animals and non-human primates and their significance for human health. *Clin Microbiol Rev* 22:202-223.
16. Joosten M, Flahou B, Meyns T, Smet A, Arts J, De Cooman L, Pasmans F, Ducatelle R, Haesebrouck F. 2013. Case report: *Helicobacter suis* infection in a pig veterinarian. *Helicobacter* 18: 392-396.
17. Joosten M, Lindén S, Rossi M, Tay ACY, Skoog E, Padra M, Peters F, Perkins T, Vandamme P, Van Nieuwerburgh F, D'Herde K, Van den Broeck W, Flahou B, Deforce D, Ducatelle R, Marshall B, Haesebrouck F, Smet A. 2015. Divergence between the Highly Virulent

- Zoonotic Pathogen *Helicobacter heilmannii* and Its Closest Relative, the Low-Virulence "*Helicobacter ailurogastricus*" sp. nov. *Infect Immun* 84:293–306.
18. Lundin A, Björkholm B, Kupersmidt I, Unema M, Nilsson P, Andersson DI, Engstrand L. 2005. Slow genetic divergence of *Helicobacter pylori* strains during long-term colonization. *Infection and Immunity* 73(8): 4818-4822.
19. Martin ME, Bhatnagar S, George MD, Paster BJ, Canfield DR, Eisen JA, Solnick JV. 2013. The impact of *Helicobacter pylori* infection on the gastric microbiota of the Rhesus Macaque. *PLoS One* 8 (10):e76375.
20. Matsui H, Takahashi T, Murayama SY, Uchiyama I, Yamaguchi K, Shigenobu S, Matsumoto T, Kawakubo M, Horiuchi K, Ota H, Osaki T, Kamiya S, Smet A, Flahou B, Ducatelle R, Haesebrouck F, Takahashi S, Nakamura S, Nakamura M. 2014. Development of new PCR primers by comparative genomics for the detection of *Helicobacter suis* in gastric biopsy specimens. *Helicobacter* 19:260-271.
21. Matsumoto T, Kawakubo M, Shiohara M, Kumagai T, Hidaka E, Yamauchi K, Oana K, Matsuzawa K, Ota H, Kawakami Y. 2009. Phylogeny of a novel "*Helicobacter heilmannii*" organism from a Japanese patient with chronic gastritis based on DNA sequence analysis of the 16S rRNA and urease genes. *J Microbiol.* 47(2):201-207.
22. Montecucco C and Rappuoli R. 2001. Living dangerously: how *Helicobacter pylori* survives in the human stomach. *Nature Reviews Molecular Cell Biology* 2: 457-466
23. Moyaert H, Pasmans F, Ducatelle R, Haesebrouck F, Baele M. 2008. Evaluation of 16SrRNA gene-based PCR assays for genus level identification of *Helicobacter* species. *J Clin Microbiol* 46 (5): 1867-1869.
24. Nakamura M, Murayama SY, Serizawa H, Sekiya Y, Eguchi M, Takahashi S, Nishikawa K, Takahashi T, Matsumoto T, Yamada H, Hibi T, Tsuchimoto K, Matsui H. 2007. "*Candidatus Helicobacter heilmannii*" from a Cynomolgus Monkey Induces Gastric Mucosa-Associated Lymphoid Tissue Lymphomas in C57BL/6 Mice. *Infect. Immun.* 75 (3): 1214-1222
25. Nakamura T, Seto M, Tajika M, Kawai H, Yokoi T, Yatabe Y, Nakamura S. 2008. Clinical Features and Prognosis of Gastric MALT Lymphoma With Special Reference to Responsiveness to *H. pylori* Eradication and API2-MALT1 Status. *The American Journal of Gastroenterology* 103, 62–70.
26. O'Rourke JL, Solnick JV, Neilan BA, Seidel K, Hayter R, Hansen LM, Lee A. 2004. Description of '*Candidatus Helicobacter heilmannii*' based on DNA sequence analysis of 16S rRNA and urease genes. *Int J of Syst and Evol Microbiol* 54:2203-2211.
27. Overby A, Murayama SY, Matsui H, Nakamura M. 2016. In the aftermath of *Helicobacter pylori*: other Helicobacters rising up to become the next gastric epidemic? *Digestion* 93: 260-265
28. Sanchez JEG, Saenz NG, Rincon MR, Martin IT, Sanchez EG, Martinez MJF. 2000. Susceptibility of *Helicobacter pylori* to mupirocin, oxazolidinones, quinupristin/dalfopristin and new quilonoles. *Journal of antimicrobial chemotherapy* 46: 283-285.
29. Schabereiter-Gurtner C, Hirschl AM, Dragosics B, Hufnagl P, Puz S, Kovach Z, Rotter M, Markistathis A. 2004. Novel Real-Time assay for the detection of *Helicobacter pylori* infection and simultaneous clarithromycin susceptibility testing of stool and biopsy specimens. *J Clin Microbiol* 42(10): 4512-4518.
30. Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics.* 22 (21): 2688-2690.
31. Zhang J, Halkilahti J, Hänninen ML, Rossi M. 2015. Refinement of whole-genome multilocus sequence typing analysis by addressing gene paralogy. *Journal of clin microciol.* 53(5): 1765-1767.

LEGENDS OF SUPPLEMENTARY TABLES

Supplementary Table 1 Overview of the separate cq-values, means and standard deviations for the standard dilution series within one run using the *carR* probe-based RT-PCR

Supplementary Table 2 Overview of the separate cq-values, means and standard deviations for the standard dilution series between 3 different runs using the *carR* probe-based RT-PCR

Supplementary Table 3 Detailed overview on the *H. suis* status and the obtained cq mean values after analysis of the gastric samples from different hosts with the *ureA*-based SYBR green RT-PCR, the *carR* probe-based RT-PCR and the *carR*-based SYBR green RT-PCR

TABLES

Table 1 Overview of the bacteria used to verify the diagnostic specificity of the probe-based RT-PCR for *H. suis* .

Group	Species	Isolate(s)	Class	Order	Amplification <i>ureA</i> -based SYBR green RT-PCR	Amplification <i>carR</i> probe- based RT-PCR	Amplification <i>carR</i> -based SYBR green RT-PCR
<i>Helicobacter</i> spp.	<i>H. suis</i>	HS1-2,HS4-9	Epsilonproteobacteria	Campylobacterales	Yes	Yes	Yes
	<i>H. heilmannii</i>	ASB1.4/11/9.4/7. 1/6.3	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. ailurogastricus</i>	ASB21_2/_7/_11	Epsilonproteobacteria	Campylobacterales	Yes	No	No
	<i>H. bizzozeronii</i>	r53/t10/ASB22 kol5/Yryla/m20/ Heydar	Epsilonproteobacteria	Campylobacterales	Yes	No	No
	<i>H. felis</i>	ds1/m26/m29/m 38- 39/dog7/jkm3/cs 5-7/106	Epsilonproteobacteria	Campylobacterales	Yes	No	No
	<i>H. salomonis</i>	m45/Yvira2/alma 0595/Inninen/	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. pylori</i>	Qir3/stm0138/J9 9/PMSS1/SS1	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. cynogastricus</i>	JKM4	Epsilonproteobacteria	Campylobacterales	Yes	No	No
	<i>H. baculiformis</i>	m50	Epsilonproteobacteria	Campylobacterales	Yes	No	No
	<i>H. cetorum</i>	015903/016202	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. cholecystus</i>	r3555	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. trogonum</i>	r3554	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. acinonychis</i>	Hacino1-3	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. mustellae</i>	img8776	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. pullorum</i>	cb103	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. marmotae</i>	cb55	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. mesocricetorum</i>	ccug45420	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. bilis</i>	atcc49320	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. equorum</i>	EqF1	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>H. canadensis</i>	H7163	Epsilonproteobacteria	Campylobacterales	No	No	No
<i>Campylobacter</i> spp.	<i>C. jejuni</i>	KC40/TB4	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>C. coli</i>	Grp028	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>C. upsaliensis</i>	img19529	Epsilonproteobacteria	Campylobacterales	No	No	No
	<i>C. fetus</i>	V24-6	Epsilonproteobacteria	Campylobacterales	No	No	No
Other bacteria	<i>Lactobacillus reuteri/ L. gasseri</i>	Cb110/Cb108	Bacilli	Lactobacillales	No	No	No
	<i>Clostridium</i>	LR3869	Clostridia	Clostridiales	No	No	No

<i>perfringens</i>						
<i>Staphylococcus aureus</i> / <i>S. pseudintermedius</i>	LR3843/LR3821	Bacilli	Bacillales	No	No	No
<i>Streptococcus suis</i> / <i>S. equi</i> subsp. <i>equi</i> / <i>S. equi</i> subsp. <i>Zooepidemicus</i>	LR3841/LR3876/LR3883	Bacilli	Lactobacillales	No	No	No
<i>Pasteurella multocida</i>	LR3892	Gammaproteobacteria	Pasteurellales	No	No	No
<i>Mannheimia haemolytica</i>	LR3870	Gammaproteobacteria	Pasteurellales	No	No	No
<i>Yersinia pseudotuberculosis</i>	LR3798	Gammaproteobacteria	Enterobacteriales	No	No	No
<i>Pseudomonas aeruginosa</i>	LR3861	Gammaproteobacteria	Pseudomonadales	No	No	No
<i>Klebsiella pneumoniae</i>	LR3806	Gammaproteobacteria	Enterobacteriales	No	No	No
<i>Proteus vulgaris</i>	LR1537	Gammaproteobacteria	Enterobacteriales	No	No	No
<i>Bacillus licheniformis</i>	DMS347	Bacilli	Bacillales	No	No	No
<i>Salmonella Typhimurium</i>	LR3862	Gammaproteobacteria	Enterobacteriales	No	No	No
<i>Trueperella pyogenes</i>	LR3863	Actinobacteria	Actinomycetales	No	No	No
<i>Escherichia coli</i>	LR3889	Gammaproteobacteria	Enterobacteriales	No	No	No
<i>Aeromonas hydrophila</i>	LR3758	Gammaproteobacteria	Aeromonadales	No	No	No
<i>Fusobacterium gastrosuis</i>	CDW01	Fusobacteria	Fusobacteriales	No	No	No
<i>Enterococcus faecalis</i>	LR3751	Bacilli	Lactobacillales	No	No	No
<i>Actinobacillus equuli</i>	ccug 19799T	Gammaproteobacteria	Pasteurellales	No	No	No

Table 2 Overview of the number of positive gastric samples, using the *ureA*-based RT-PCR, the *carR* probe-based RT-PCR and the *carR*-based SYBR green RT-PCR. Gastric samples were derived from pigs (natural host), humans, non-human primates (natural host) and mice (experimental infections).

Host gastric samples	Number of samples	Number of <i>H. suis</i> positive samples	Number of positive samples using <i>ureA</i> -based RT-PCR	Number of positive samples using <i>carR</i> probe-based RT-PCR	Number of positive samples using <i>carR</i> -based SYBR green RT-PCR
Human (with or without natural <i>H. suis</i> infection)	80	6	7	6	0
Non-human primates (with or without natural <i>H. suis</i> infection)	9	8	9	8	6
Mice (experimentally infected animals and control animals)	8	4	4	4	0
Pigs (with or without natural <i>H. suis</i> infection)	9	5	5	5	0

Table 3 Overview of the mean number and standard deviation of *H. suis* bacteria/μl (cq-mean) in the positive gastric samples per host, using *ureA*-based RT-PCR and *carR* probe-based RT-PCR respectively. Gastric samples were derived from pigs, humans, non-human primates and mice.

	Cq-mean <i>ureA</i> -based SYBR green RT-PCR	Cq-mean <i>carR</i> probe-based RT-PCR	Stdev <i>ureA</i> -based SYBR green RT-PCR	Stdev <i>carR</i> probe-based RT-PCR
Human positive gastric samples	9.68E+02	2.68E+01	2.13E+03	3.49E+01
Non-human primate positive gastric samples	1.38E+06	7.79E+04	3.95E+06	1.55E+05
Murine positive gastric samples	1.14E+04	4.20E+03	3.85E+03	2.41E+03
Porcine positive gastric samples	5.82E+02	2.50E+02	7.94E+02	3.44E+02

SUPPLEMENTARY TABLES

Supplementary Table 1

	1,00E+00	1,00E+01	1,00E+02	1,00E+03	1,00E+04	1,00E+05	1,00E+06	1,00E+07	1,00E+08	1,00E+09
Standard 1	37.57	33.47	29.07	25.38	21.98	18.49	15.16	11.92	7.53	3.26
Standard 2	35.03	32.32	29.1	25.92	22.11	18.99	15.21	11.86	7.93	3.59
Standard 3	36.49	33.11	29.08	26.03	21.93	18.42	15.25	11.98	7.67	3.08
Standard 4	35.86	32.59	29.28	25.52	22.01	18.98	15.1	11.84	7.52	3.48
Standard 5	36.99	32.75	28.85	25.5	22.05	18.74	15.04	11.75	7.68	3.57
Standard 6	35.37	33.03	29.31	25.58	22.09	18.96	15.43	11.85	7.9	3.56
Mean	36.2183	32.878	29.115	25.655	22.028	18.7633	15.1983	11.8667	7.705	3.42333
SD	0.97534	0.4094	0.1667	0.25859	0.0682	0.25696	0.13615	0.07789	0.17627	0.20772
Efficiency run (%)	89.9									
Linear correlation (r²)	0.998									

Supplementary Table 2

	1,00E+00	1,00E+01	1,00E+02	1,00E+03	1,00E+04	1,00E+05	1,00E+06	1,00E+07	1,00E+08	1,00E+09
Run 1	37.07	32.43	28.48	25.09	21.58	18.25	14.92	11.57	6.99	2.47
	36.41	31.98	28.42	24.96	21.59	18.28	14.93	11.73	7.11	1.79
	36.39	31.66	28.46	25.07	21.56	18.28	14.86	11.46	7.19	2.35
Run 2	34.66	31.71	28.47	25.15	21.92	18.43	14.8	11.67	6.96	2.97
	35.51	32.6	28.54	25.18	21.83	18.45	14.92	11.76	7.76	3.44
	36.03	32	28.62	25.17	21.82	18.41	14.93	11.65	7.29	3.07
Run 3	no value (nv)	32.37	29.18	26.11	22.25	18.6	15.57	12.09	7.83	3.24
	36.14	32.62	29.3	25.99	22.37	18.66	15.62	12.13	8.01	3.8
	nv	32.78	29.26	25.55	22.24	18.61	15.53	11.97	7.8	3.24
Mean	36.03	32.239	28.7478	25.3633	21.907	18.4411	15.12	11.7811	7.43778	2.93
SD	0.7654	0.4125	0.3796	0.42205	0.3131	0.15463	0.34329	0.23267	0.40862	0.62028
Mean efficiency run 1,2 and 3	90.03									
Mean linear correlation run 1,2 and 3	0.998									

Supplementary Table 3

Samples	<i>H. suis</i> status	<i>UreA</i> -based SYBR green RT-PCR	<i>CarR</i> probe-based RT-PCR	<i>CarR</i> -based SYBR green RT-PCR
Preserve Human DNA (n=80)				
human 1	Absent	Negative	Negative	Negative
human 2	Absent	Negative	Negative	Negative
human 3	Absent	Negative	Negative	Negative
human 4	Absent	Negative	Negative	Negative
human 5	Absent	Negative	Negative	Negative
human 6	Absent	Negative	Negative	Negative
human 7	Absent	Negative	Negative	Negative
human 8	Absent	Negative	Negative	Negative
human 9	Absent	Negative	Negative	Negative
human 10	Absent	Negative	Negative	Negative
human 11	Absent	Negative	Negative	Negative
human 12	Absent	Negative	Negative	Negative
human 13	Absent	Negative	Negative	Negative
human 14	Absent	Negative	Negative	Negative
human 15	Absent	Negative	Negative	Negative
human 16	Absent	Negative	Negative	Negative
human 17	Absent	Negative	Negative	Negative
human 18	Absent	Negative	Negative	Negative
human 19	Absent	Negative	Negative	Negative
human 20	Absent	Negative	Negative	Negative
human 21	Absent	Negative	Negative	Negative
human 22	Absent	Negative	Negative	Negative
human 23	Absent	Negative	Negative	Negative
human 24	Absent	Negative	Negative	Negative
human 25	Absent	Negative	Negative	Negative
human 26	Absent	Negative	Negative	Negative
human 27	Absent	Negative	Negative	Negative
human 28	Absent	Negative	Negative	Negative
human 29	Absent	Negative	Negative	Negative
human 30	Absent	Negative	Negative	Negative
human 31	Absent	Negative	Negative	Negative
human 32	Absent	Negative	Negative	Negative
human 33	Absent	Negative	Negative	Negative
human 34	Absent	Negative	Negative	Negative

human 35	Absent	Negative	Negative	Negative
human 36	Absent	Negative	Negative	Negative
human 37	Absent	Negative	Negative	Negative
human 38	Absent	Negative	Negative	Negative
human 39	Absent	Negative	Negative	Negative
human 40	Absent	Negative	Negative	Negative
human 41	Absent	Negative	Negative	Negative
human 42	Absent	Negative	Negative	Negative
human 43	Absent	Negative	Negative	Negative
human 44	Absent	Negative	Negative	Negative
human 45	Absent	Negative	Negative	Negative
human 46	Absent	Negative	Negative	Negative
human 47	Absent	Negative	Negative	Negative
human 48	Absent	Negative	Negative	Negative
human 49	Absent	Negative	Negative	Negative
human 50	Absent	Negative	Negative	Negative
human 51	Absent	Negative	Negative	Negative
human 52	Absent	Negative	Negative	Negative
human 53	Absent	Negative	Negative	Negative
human 54	Absent	Negative	Negative	Negative
human 55	Absent	Negative	Negative	Negative
human 56	Absent	Negative	Negative	Negative
human 57	Absent	Negative	Negative	Negative
human 58	Absent	Negative	Negative	Negative
human 59	Absent	Negative	Negative	Negative
human 60	Absent	Negative	Negative	Negative
human 61	Absent	Negative	Negative	Negative
human 62	Absent	Negative	Negative	Negative
human 63	Absent	Negative	Negative	Negative
human 64	Absent	Negative	Negative	Negative
human 65	Absent	Negative	Negative	Negative
human 66	Absent	Negative	Negative	Negative
human 67	Absent	Negative	Negative	Negative
human 68	Absent	Negative	Negative	Negative
human 69	Absent	Negative	Negative	Negative
human 70	Absent	Negative	Negative	Negative
human 71	Absent	Negative	Negative	Negative
human 72	Absent	Negative	Negative	Negative
human 73	Absent	Negative	Negative	Negative
human 74	Present	2.51E+01	2.05E+00	Negative
human 75	Present	1.07E+02	3.39E+01	Negative

human 76	Present	5.78E+03	9.21E+01	Negative
human 77	Present	4.65E+02	1.74E+00	Negative
human 78	Absent	2.12E+01	Negative	Negative
human 79	Present	3.32E+02	2.74E+01	Negative
human 80	Present	4.39E+01	3.69E+00	Negative
Non-human primate DNA (n=9)				
JI 503 biopt a	Present	1.49E+02	4.96E+00	Negative
Ro2019 bio a	Present	2.89E+04	2.71E+03	7.14E+02
r08041 bio a	Present	1.74E+04	1.28E+04	Negative
JI 190 bio a	Absent	3.96E+02	Negative	Negative
HSM1Ro40522 dna	Present	1.19E+07	4.54E+05	7.60E+05
baviaan 9 bio a	Present	1.03E+05	9.08E+04	2.02E+03
baviaan 9 bio f	Present	2.34E+04	1.34E+04	1.08E+03
rhesus aap bio a	Present	9.36E+04	1.92E+04	2.72E+03
rhesus aap bio f	Present	2.53E+05	3.04E+04	4.79E+03
Mouse DNA (n=8)				
1m2 c1 c	Absent	Negative	Negative	Negative
1m2 c2 c	Absent	Negative	Negative	Negative
1m2 c3 c	Absent	Negative	Negative	Negative
1m2 c4 c	Absent	Negative	Negative	Negative
1m2 s1 c	Present	1.22E+04	3.66E+03	Negative
1m2 s2 c	Present	1.60E+04	7.61E+03	Negative
1m2 s3 c	Present	1.08E+04	3.63E+03	Negative
1m2 s4 c	Present	6.68E+03	1.91E+03	Negative
Pig DNA (n=9)				
p17 c	Absent	Negative	Negative	Negative
p18 c	Absent	Negative	Negative	Negative
p20 c	Absent	Negative	Negative	Negative
p11 c	Present	1.72E+03	5.14E+02	Negative
p27 c	Present	4.03E+01	8.42E+00	Negative
p28 c	Present	2.22E+01	5.05E+00	Negative
p33 c	Absent	Negative	Negative	Negative
p49 c	Present	7.15E+00	2.25E+00	Negative
p50 c	Present	1.12E+03	7.22E+02	Negative

GENERAL DISCUSSION

...“all the diseases begin in the gut”... by Hippocrates

Besides the well-known human pathogen *Helicobacter pylori* (*H. pylori*), non-*Helicobacter pylori* *Helicobacter* (NHPH) species from domestic animals have also been associated with gastric disease in humans. Many of these very fastidious bacteria have recently been *in vitro* isolated from their natural hosts, which has led to progress in understanding the pathogenesis of NHPH-related pathologies. A distinct feature is their association with gastric mucosa-associated lymphoid tissue (MALT) lymphoma in humans, but this is most probably a tip of the iceberg...

Increased frequency of gastric *H. suis* infections in human patients with Parkinson’s disease (PD): coincidence or an underlying cause in the pathogenesis of this disease?

H. pylori may interfere with many biological processes, both inside and outside the stomach, possibly influencing or determining the occurrence and/or outcome of many extragastric diseases (Alvarez-Arellano et al., 2014; Budzynski et al., 2014; Smyk et al., 2014; Testerman et al., 2014; Wong et al., 2014). While its role in idiopathic thrombocytopenic purpura (Radic et al., 2014; Hasni et al., 2011; Yamanishi et al., 2006) and sideropenic anemia (Kang et al., 2011; Monzon et al., 2013) has already been acknowledged, emerging evidence suggests that *H. pylori* may be associated with neurodegenerative disorders as well (Csoti et al., 2016; Alvarez-Arellano et al., 2014; Budzynski et al., 2014; Dobbs et al., 2008; Dobbs et al., 2010; Dobbs et al., 2016; Nielsen et al., 2012; Rees et al., 2011). PD is a degenerative disorder of the central nervous system (CNS) characterized by cell death of dopamine-generating neurons in the substantia nigra, a region of the basal ganglia of the brain (Wirdefeldt et al., 2011; Mhyre et al., 2012; Kannarkat et al., 2013). The familial form of PD is caused by mutations in several different genes (including the α -synuclein gene and the *parkin* gene), whereas the aetiology of non-heritable idiopathic Parkinsonism (IP) remains unknown, although several risk factors, such as rural residence, livestock farming, head trauma, well-water drinking, infectious diseases and exposure to pesticides, have been associated with the appearance of this disease state (Lee et al., 2002; Schiesling et al., 2008; Dobbs et al., 2008; Hardy et al., 2006). Besides aberrations in motor functions, increased gastrointestinal dysfunction is well-acknowledged in patients with parkinsonism and usually precedes the diagnosis of PD by many years (Edwards et al., 1992; Charlett et al., 1997; Abbott et al., 2001; Pfeiffer, 2011; Dobbs et al., 2016). The finding of prodromal peptic ulcers in patients with PD in 1965 (Strang, 1965) paved the way to explore any link between (the then not yet described (Warren et al., 1983)) *H. pylori* and PD. Infection with *H. pylori* has been shown to be an arbiter for progression of the poverty of movement (hypokinesia) (Dobbs et al., 2008; Dobbs et al., 2010; Dobbs et al., 2012; Dobbs et al., 2013).

In some cases, decreased absorption of levodopa, which is the main drug used to treat PD, may be involved (Pierantozzi et al., 2001; Hashim et al., 2014). Indeed, eradicating *H. pylori* in PD patients by applying antibiotic treatment results in an increased absorption of levodopa and decreased motor fluctuations (Hashim et al., 2014). Although it remains unknown how *H. pylori* may interfere with levodopa absorption, several hypotheses have been proposed. Since the solubility of levodopa is pH dependent, *H. pylori*-induced gastric pH alterations

might influence absorption (Annibale et al., 2002). Furthermore, levodopa absorption may also be altered by *H. pylori*-associated gastric myoelectric dysfunctioning resulting in delayed gastric emptying (Miyaji et al., 1999). Finally, it has been suggested that *H. pylori* might utilize levodopa for growth (Doherty et al., 2009).

Whilst influence of *H. pylori* on levodopa absorption may be relevant to symptomatic treatment of PD, it does not provide evidence on aetio-pathogenesis. This was shown in a randomized, double-blind, placebo-controlled trial. Objectively-measured gait hypokinesia improved over the year following biopsy-proven (culture and polymerase chain reaction (PCR)) *H. pylori* eradication, and did not deteriorate over the next two years. The effect on hypokinesia was seen in patients never given symptomatic treatment for PD, and was no greater in those receiving background anti-parkinsonian medication. Thus, it was not due to improved absorption of parkinsonian medication (Dobbs et al 2010). This was confirmed in a longitudinal observational study of the effects of routine antimicrobial use on PD. Hypokinesia again improved after treating *H. pylori* infection, but antimicrobials given for other indications did not affect it (Dobbs et al 2013).

Activation of microglia cells, resulting in neuroinflammation and apoptosis of nerve cells as a consequence of chronic gastric inflammation may be a possible explanation for the association that was found between *H. pylori* infection and neurodegenerative disorders (Dobbs et al., 2000; Dobbs et al., 2008; Dobbs et al., 2010; Dobbs et al., 2012; Kountouras et al., 2012; Dobbs et al., 2016). This is described in more detail in the next heading, dealing with the stomach-brain axis.

Molecular mimicry between *H. pylori* antigens and human antigens, presented by dopamine producing neurons has also been proposed (Dobbs et al., 2000; Schulz et al., 2006; Dobbs et al., 2008; Dobbs et al., 2010; Dobbs et al., 2012; Dobbs et al., 2016). There is molecular mimicry between *H. pylori* and gastric antigens (Appelmek et al., 1997) and other studies reported the presence of auto-antibodies against dopamine-producing neurons in blood and the cerebrospinal fluid in patients with PD (Poupard and Emilie, 1984; Fiszer et al., 1996; Schulz et al., 2006; Dobbs et al., 2008; Dobbs et al., 2010; Dobbs et al., 2012; Dobbs et al., 2016).

It has been suggested that living or working on livestock farms may increase the risk of PD (Gorell et al., 1998; Lee et al., 2002; Wirdefeldt et al., 2011), raising the possibility that zoonotic transmission of animal-associated gastric helicobacters might also contribute to the etiology of this disease. There has been much attention to the possible dangers of herbicides, but arable farmers have a reduced risk of dying from PD (Lee et al., 2002). *H. suis* is highly prevalent in the stomach of pigs and is the most prevalent NHPH species in humans (Haesebrouck et al., 2009). A case report published in 2013 described a *H. suis* infection in a pig veterinarian suffering from reflux esophagitis and general dyspeptic symptoms (Joosten et al., 2013). Using multilocus sequence typing (MLST), it was demonstrated that the *H. suis* strain, detected in the stomach of the pig veterinarian, was closely related to porcine strains confirming the zoonotic potential of this *Helicobacter* species (Liang et al., 2013). It has been suggested that direct contact with pigs might be an important source for *H. suis* infection in humans (Haesebrouck et al., 2009; De Groote et al., 2005; Van den Bulck et al., 2005; Baele et al., 2008). Additionally, the consumption of raw or undercooked pork might also constitute a source of *H. suis* infection in humans (De Cooman et al., 2013).

Only few information about a possible association between gastric *H. suis* infections and PD was available at the onset of this thesis. There was a report of improvement in PD following eradication of a spiral gastric NHPH, reported at that time to be *H. heilmannii* (Dobbs et al., 2005), but which we showed to be *H. suis*. Therefore, **in the first chapter** of this thesis, we reinvestigated deoxyribonucleic acid (DNA), extracted from human gastric biopsy samples and archived at a *Helicobacter* reference laboratory, for the presence of *H. suis*. An exceptional high frequency (27%) of *H. suis* DNA was found in extracts from patients with PD, compared with those from a routine gastroenterology service referrals, in whom no PD had been documented (2%). *H. suis* DNA was more often detected in patients with biopsy proven *H. pylori* eradication. Based on this finding, it can be hypothesized that these patients were co-infected with *H. pylori* and *H. suis* as described before (Liu et al., 2015). Triple therapy, usually consisting of amoxicillin, clarithromycin and a proton pump inhibitor, may have been ineffective to eradicate *H. suis* giving this bacterium the opportunity to multiply in the human stomach and to cause gastric disease. Results of previous studies gave evidence of intrinsic reduced susceptibility to aminopenicillins of porcine *H. suis* isolates compared to *H. pylori* (Vermoote et al., 2011). In addition, recent *in vitro* antimicrobial susceptibility testing showed evidence of acquired resistance to clarithromycin among *H. suis* isolates (De Witte Chloë, 2017, personal communication). Both findings strengthen the above mentioned assumption. Indeed, in the case referred to above (Dobbs et al 2005), there was short-lived improvement in PD following standard triple therapy, which relapsed with recrudescence of *H. suis*. Successful (biopsy-proven) *H. suis* eradication resulted in long-term improvement. Nevertheless, successful eradication of *H. suis* has been described after treatment with amoxicillin, clarithromycin and a proton pump inhibitor (Joosten et al., 2013; Hellemans et al., 2005). These differences in treatment outcome, might support previous suggestions on the occurrence of acquired antimicrobial resistance (Vermoote et al., 2011; De Witte et al., 2017, personal communications).

We additionally described the presence of *H. suis* DNA in a blood sample of a patient with PD and unpublished results from recent *in vivo* experiments in mice showed the presence of *Helicobacter* antigens in macrophages at the level of mesenteric lymph nodes during chronic infection. Together, all these findings suggest that the effects caused by *H. suis* are not restricted to the stomach. Moreover, *H. pylori* DNA, and even culturable organisms have been found in gastric lymph nodes from human surgical specimens (Ito et al., 2008). Clearly, the role of *Helicobacter* in general and *H. suis* in particular, in PD requires further investigation.

The stomach-brain axis in relation to a *H. suis* infection: does *H. suis* affect neuroinflammation?

Evidence is accumulating that neuroinflammation plays an important role in the pathophysiology of PD. Activated microglia can be detected in the brain of PD patients, reflected by a change in morphology from ramified to a more round shape (Wang et al., 2015; Tansey et al., 2007; Hirsch and Hunot, 2009). Microglia activation results in the release of pro-inflammatory cytokines which can act on neuronal integrity (Wang et al., 2015; Fonseca et al., 2014). The intermediate stage, “primed microglia”, describes the atypical microglial stage, which precedes a further neurotoxic microglial activation as a consequence of a secondary pro-inflammatory stimulus which can come from the periphery (Perry et al., 2013). Indeed, peripheral inflammation can activate microglia, in particular when the brain barriers are disturbed (Wang et al., 2015). Several specialized CNS

barriers, such as the blood-brain barrier (BBB) and the blood-cerebrospinal fluid (CSF) barrier, play a crucial role in the maintenance of brain homeostasis, since they determine what will (and what will not) enter the brain (De Bock et al., 2014). The BBB, formed by capillary endothelial cells which are connected by tight junctions, separates the circulating blood from the extracellular fluid in the CNS. The blood-CSF barrier is established by the choroid plexus epithelial cells, which secrete the CSF and have an essential function in the communication between the peripheral immune system and the CNS (De Bock et al., 2014). Dysfunction of these brain barriers has been implicated in the pathogenesis of PD (Vawter et al., 1996; Pisani et al., 2012; Stolp et al., 2009; Lee et al., 2014).

Interactions between the gastrointestinal tract and the brain are well recognized (Cryan et al., 2012) and in particular, a bidirectional communication between the CNS and the gastrointestinal tract via the gut-brain axis has been considered. Moreover, it has been suggested that neuroinflammation could emanate from infection in the gastrointestinal tract (Dobbs et al., 2000; Dobbs et al., 2008; Dobbs et al., 2010; Ferrari et al., 2011; Dobbs et al., 2013; Dobbs et al., 2016). There is evidence that intestinal infection or dysbiosis plays a part in the pathogenesis. There are biological gradients between leukocyte subset counts and facets on PD (Dobbs et al. 2012). The same subsets are associated with hydrogen-breath-test positivity for small-intestinal bacterial overgrowth. In PD, successive courses of antimicrobials are associated with cumulative increase in flexor rigidity (Dobbs et al. 2013), which may suggest progressively increasing colonic dysbiosis (Carethers, 2011). There still appears to be a specific effect of *Helicobacter*: it had an effect on hypokinesia in addition to that on natural-killer cell count (Dobbs et al. 2012), and routine treatment of *Helicobacter* produced an indication specific improvement in hypokinesia (Dobbs et al. 2013). Gastric *Helicobacter* spp. generally cause a life-long gastric infection since the induced host immune response is not able to eradicate the infection, resulting in persistent inflammation and gastric epithelial cell damage (Montecucco & Rapuoli, 2001; Flahou et al., 2011; Flahou et al., 2012).

A recently proposed hypothesis is summarized in **Figure 1**: *Helicobacter* might influence the brain through the blood, since injury of the gastric mucosa might lead to leakage of bacterial components and metabolites, and even pro-inflammatory factors into the blood circulation resulting in systemic inflammation (Fukuda et al., 2001; Sun et al., 2004; Fedwick et al., 2005; Guttman and Finlay, 2009; Wroblewski et al., 2011; Caron et al., 2015; Cryan et al., 2014; Mayer et al., 2014). These inflammatory mediators may then contribute to disturbance of the brain homeostasis leading to neurodegenerative diseases (Dobbs et al., 2000; Dobbs et al., 2008; Dobbs et al., 2010; Dobbs et al., 2012; Kontouras et al., 2015; Dobbs et al., 2016; Charlett et al. 1998; Dobbs et al. 1999; Reale et al., 2009). Furthermore, several reports suggested the involvement of systemic inflammation in the impairment of the BBB (Kortekaas et al., 2005; Stolp et al., 2009) and the blood-CSF barrier (Vandenbroucke et al., 2012; Demeestere et al., 2015; Villaran et al., 2010; Perry et al., 2013).




Figure 1: Stomach-brain axis via systemic circulation. A gastric infection with *Helicobacter* may evoke inflammation resulting in a disruption of the gastric mucosal barrier. In turn this could lead to leakage of bacterial components and pro-inflammatory factors from the stomach into the systemic circulation, resulting in systemic inflammation. These inflammatory mediators may then contribute to disturbance of the brain homeostasis leading to neurodegenerative diseases. Figure was adapted from Baele et al., 2008 (*H. suis*), Cryan et al., 2012 (vessel), Mayer et al., 2014 (brain), Servier Medical Art (stomach).

Elaborating on the high frequency of *H. suis* infections in patients with PD as shown in **chapter 1** and keeping in mind the above mentioned hypothesis that *Helicobacter* might influence the brain through the blood, in the **second chapter** of this thesis we examined whether a gastric *H. suis* infection has an impact on the integrity of the gastric mucosal barrier and the brain barriers through the induction of inflammation. For this purpose, mice were infected with a virulent *H. suis* strain as shown before (Flahou et al., 2010). At 4 weeks post-infection, the inflammatory response induced by *H. suis* was characterized by an infiltration of immune cells in the gastric mucosa and submucosa as well as by the secretion of interleukin (IL)-1 β , IL-17 and the human IL-8-related murine cytokines, lipopolysaccharide (LPS)-induced CXC chemokine (LIX) and keratinocyte-derived chemokine (KC), at the site of infection. These findings were in line with previous *H. suis* experimental infection studies as well as with *in vivo* trials with other gastric helicobacters (Flahou et al., 2012; Flahou et al., 2010; Bosschem et al., 2016; Peek et al., 2010; Alvarez-Arellano et al., 2014; Hashim et al., 2014). Since the pro-inflammatory cytokine IL-1 β is a potent inhibitor of gastric acid secretion, it facilitates *Helicobacter* colonization of the gastric mucosa (Joosten et al., 2013; Semper et al., 2014). It also plays a crucial role in the development of gastric

pathologies (Joosten et al., 2013). Its secretion depends on the activation of the inflammasome, an intracellular multiprotein oligomer implicated in the activation of inflammatory processes (Semper et al., 2014). *H. pylori* induced inflammasome activation in dendritic cells (DCs) requires the nucleotide-binding oligomerization domain, leucine rich repeat and pyrin domain containing 3 (NLRP3) (Semper et al., 2014) and depends on the presence of the vacuolating cytotoxin (VacA) and cag pathogenicity island (cagPAI) virulence factors (Semper et al., 2014; Guo et al., 2015). In contrast to *H. pylori*, *H. suis* lacks both VacA and cagPAI (Vermoote et al., 2011). It is currently unknown how this bacterium triggers secretion of IL-1 β . Screening of a *H. suis* transposon mutant library, as previously developed for *H. pylori* (Odenbreit et al., 1996), for its ability to activate the inflammasome may help to identify the virulence factors involved in this process.

Besides IL-1 β , expression of the IL-8-related murine pro-inflammatory cytokines, LIX and KC were also significantly increased in the gastric mucosa exposed to *H. suis*. IL-8 is an important chemotactic and activating factor for neutrophils (Crabtree and Lindley, 1994) and can be upregulated by IL-1 β and tumour necrosis factor (TNF) through the nuclear factor (NF)- κ B signalling pathway (Liu et al., 2016; Crabtree and Lindley, 1994; Kim et al., 2001; Torok et al., 2005). Additionally, IL-8 secretion can be stimulated by the presence of IL-17 via the extracellular signal-regulated kinase (ERK) 1/2 mitogen-activating protein (MAP) kinase pathway (Kabir, 2011; Bagheri et al., 2015). The *H. pylori* virulence factor cagPAI and several *H. pylori* outer membrane proteins (OMPs) acting as adhesins (e.g. blood group antigen binding adhesion A (BabA), sialic acid binding adhesion A (SabA), outer inflammatory protein A (oipA), etc.), have been shown to be involved in the increased IL-8 production by gastric epithelial cells (White et al., 2015). Both cagPAI and these *H. pylori* adhesins are absent in *H. suis* (Vermoote et al., 2011). Recently, the *Helicobacter* outer membrane protein family (Hof)E and HofF proteins of *H. heilmannii* have been shown to be partly involved in the induction of IL-8 expression in the gastric epithelium (Liu et al., 2016). These OMPs are also present in *H. suis* and it remains to be elucidated whether they are involved in this process as well.

The extensive secretion of IL-8 or the related murine cytokines, LIX and KC, by gastric epithelial cells upon *Helicobacter* infection is probably a key factor in host defenses at the mucosal site. Since the immune mechanisms usually fail to eradicate *H. suis*, resulting in chronic infection (Flahou et al., 2010), continued upregulation of these pro-inflammatory cytokines could lead to gastric mucosal damage (Crabtree and Lindley, 1994). Indeed, in our *in vivo* experiment, the *H. suis*-evoked gastric immune response was accompanied by a disruption of the gastric mucosal barrier. This was characterized by an increased gastrointestinal barrier permeability and alterations in gastric tight junction proteins. A disruption of the gastric epithelial barrier, characterized by a loss of claudin-4, claudin-5 and occludin (Fedwick et al., 2005; Wroblewski et al., 2011) in combination with a reorganization of ZO-1 and claudin-1 (Guttman et al., 2009; Caron et al., 2015) has also been described during *H. pylori* infections.

Tight junctions regulate cell polarity and their formation is dependent on the pseudo-autosomal region (PAR)3-PAR6-atypical protein kinase C (aPKC) (PAR) complex (Vermeer et al., 2003). Loss of tight junctions can be mediated by receptor tyrosine kinases, which are members of the epidermal growth factor receptor (ErbB) family. Activation of ErbB2 has been shown to disrupt polarity by associating with PAR6 and aPKC and blocking the interaction with PAR3 (Aranda et al., 2006). Transmembrane mucins, including mucin MUC13, can

participate in receptor tyrosine kinase activation and can thereby contribute to the disruption of the PAR complex (Ren et al., 2006). Since MUC13 is aberrantly expressed upon *H. suis* infection and inflammation (Liu et al., 2016), it remains to be elucidated whether MUC13 plays a role in gastric mucosal barrier disruption. In order to determine whether MUC13 interacts with one of the members of the ErbB family present on the gastric epithelial cells, co-immunoprecipitation can be performed. Additionally, the role of the MUC13 signaling pathway in the PAR complex and the disruption of tight junctions in the stomach can be investigated using *in vitro* transfection assays. If the transfection experiments indicate that MUC13 plays a role in the tight junction regulation, it is recommended to evaluate whether these changes are correlated with a loss of barrier functionality using an electrical cell impedance system (ECIS) or transepithelial electrical resistance (TEER) measurements in transwells.

Apart from the γ -glutamyl transpeptidase (GGT) enzyme and urease (Flahou et al., 2011), no other information is currently available on *H. suis* virulence factors involved in gastric mucosal damage. Unpublished results from a whole transcriptome analysis of *H. heilmannii* showed that GGT is significantly upregulated when this bacterium is attached to gastric epithelial cells. Other virulence factors which are altered in expression in relation to adherence can thus be potential candidates involved in gastric mucosal barrier disruption and should be further investigated. This investigation can be facilitated by chromosomal insertional mutagenesis approaches and *in vitro* and *in vivo* infection assays using the developed mutants.

Disruption of the gastric mucosal barrier may result in leakage of pro-inflammatory cytokines, chemokines and bacterial metabolites and components, such as LPS, into the blood circulation, thereby inducing systemic inflammation (Cryan et al., 2012; Mayer et al., 2014). In our study, significantly increased levels of IL-1 β and the IL-8-related murine cytokine, macrophage inflammatory protein (MIP)1 α , were found in the serum of *H. suis*-infected mice resulting in systemic inflammation. This might thus lead to brain barrier dysfunction and subsequent alterations in brain homeostasis (Alvarez-Arellano et al., 2014; Mulak et al., 2015), as might the elevated serum cortisol and IL-6 found in PD (Charlett et al., 1998; Dobbs et al., 1999). Indeed, the integrity of the brain barriers in the *H. suis*-infected mice was compromised. Infection with *H. suis* resulted in changes in tight junction expression of the BBB and the blood-CSF barrier, but increased permeability was only detected at the blood-CSF barrier of *H. suis*-infected animals. Also neuroinflammation, characterized by the activation of microglia cells and the upregulation of pro-inflammatory cytokines, in particular IL-1 β and inducible nitric oxide synthase (iNos) in the choroid plexus and IL-6 in the hippocampus of the brain, was detected. Downregulation of tight junction genes in the hippocampus did not result in increased BBB leakage, suggesting that *H. suis* infection has a higher impact on the blood-CSF barrier compared to the BBB. Intraperitoneal injection of mice with LPS, a component which is also present in *H. suis* and other Gram-negative bacteria, also resulted in increased leakage of the blood-CSF barrier as shown before (Vandenbroucke et al., 2012).

Up to date, it remains unknown whether *H. suis* LPS might be involved in the disruption of the integrity of the brain barriers and brain homeostasis. The administration of the antibiotic polymyxin B, which inhibits different biological activities induced by LPS, might be useful to answer this question (Cavaillon et al., 1986). Another possibility to investigate the role of *H. suis* LPS is by creating an LPS mutant of *H. suis*, as already described for *Escherichia coli* (Liu et al., 2015), and repeat the experiment using the LPS-mutant of *H. suis*.

In our studies, fluorescein isothiocyanate (FITC)-dextran was used to determine the BBB and blood-CSF barrier permeability. The application of FITC-dextran is considered a valuable method for the characterization of alterations in permeability of both the BBB and the blood-CSF barrier (Hoffmann et al., 2011; Vandenbroucke et al., 2012). The advantages of the use of FITC dextran as a tool is that FITC dextrans can be obtained in different molecular sizes and that they can be used to determine both solute and ion permeability (small molecular size) and protein permeability (high molecular size) (Hoffmann et al., 2011). However, it is not recommended to use samples from FITC treated animals in combination with paraformaldehyde (PFA) for histological examination, because FITC dextran dissolves in PFA, leading to false negative results (Hoffmann et al., 2011).

An important finding throughout our study was the presence of the pro-inflammatory cytokine IL-1 β not only in the inflamed stomach but also in the serum and choroid plexus of the *H. suis*-infected animals, highlighting its role in the induction of neuroinflammation via systemic communication. The role of IL-1 β in the integrity of the stomach-brain axis and induction of neuroinflammation requires further investigation. The usage of *H. suis*-infected IL-1 $\beta^{-/-}$ mice might shed more light on this.

Besides systemic communication, neural communication via neuropeptides, such as VIP, Substance P (SP), calcitonin-gene related peptide (CGRP), somatostatin and neuropeptide Y, may also play a role (Delgado et al., 2004; Erin et al., 2012; Mayer et al., 2014; De Vadder et al., 2015). Therefore we have also investigated the effect of an *H. suis* infection on VIP signaling. A remarkable finding in the stomach was the altered expression of the vasoactive intestinal peptide (VIP) and its receptor *VIPR1* in *H. suis*-infected animals compared to controls. Neuropeptides, such as VIP, have been shown to promote homeostasis of the mucosal barrier of the gastrointestinal tract and have thus been implicated in tight junction regulation (Chen et al., 2015; Anlauf et al., 2003; Mulak et al., 2015). Alterations in mucosal VIP levels have also been reported in biopsies derived from human patients with chronic gastritis and ulcers (Erin et al., 2012; Chen et al., 2015; Islek et al., 2016; Li et al., 2009; Bercik et al., 2002). Changes in VIP expression might thus induce dysfunction of the gastrointestinal barrier and in this way contribute to the formation of gastric pathologies.

Up to date, the mechanisms by which *H. suis* might induce changes in VIP expression remain to be elucidated. However since alterations in VIP and its receptors have been reported in combination with gastric pathology (Erin et al., 2012), *H. suis*-induced gastric inflammation might be a possible explanation for the observed alterations in *VIP* and *VIPR1* expression upon *H. suis* infection.

VIP does not only contribute to maintenance of the gastrointestinal epithelial barrier integrity (Conlin et al., 2009), but it might also be involved in brain homeostasis. Indeed, fibers containing VIP have been identified close to the choroid plexus and it has been described that VIP regulates CSF secretion of the choroid plexus epithelial cells (Delgado et al., 2003; Nilsson et al., 1991; Lindvall et al., 1978). In addition, the protective role of VIP in neurodegenerative disorders, such as PD, has been reported (Delgado et al., 2003). Furthermore, it has been described that VIP is able to prevent overactivation of microglia in LPS-induced inflammation in the brain (Delgado et al., 2003; Kim et al., 2000). In our study, administration of VIP reduced LPS-induced blood-CSF barrier leakage and LPS-induced upregulation of *IL-6*, *TNF*, *KC* and *iNos* in the choroid plexus. Since VIP is present and produced in the CNS and the gastrointestinal system, its potential protective role in the

preservation of integrity of the epithelial barriers of gut and brain should not be neglected. The use of a VIP-knock-out mouse model might provide information on the role of VIP in *H. suis*-induced disruption of the integrity of the gut and brain barriers.

Taken together, our results suggest that increased gastrointestinal leakage and disturbance of the blood-CSF barrier along with neuroinflammation may be induced by *H. suis*-associated inflammation and dysregulation of the VIP signalling. Furthermore, the pivotal role of the choroid plexus in the communication between the stomach and the brain via systemic inflammation and dysregulation of VIP signalling has been identified. Nevertheless, further research is required to elucidate the underlying mechanisms by which an infection with *H. suis* may influence gastrointestinal–brain communication.

Optimization of diagnostic tools for the detection of *H. suis* in human samples

In general, diagnosing gastric NHPH infections in humans is problematic. Non-invasive commercial tests are currently unavailable. The urea breath test, frequently used to diagnose *H. pylori* infections, is usually negative in patients infected with NHPH, except when *H. pylori* co-exists (Matsui et al., 2014). The patchiness and sparseness of the gastric colonization by NHPH is probably the underlying cause for these false-negative results (Stolte et al., 1997; Solnick and Schauer, 2001). Serologic tests to screen for NHPH infections are lacking. Up until the present day, infection with these fastidious pathogenic agents is diagnosed by the analysis of gastric biopsies using histological, molecular and microbiological techniques (Debongnie et al., 1995; De Groote et al., 2001; Baele et al., 2004; O'Rourke et al., 2004; Matsui et al., 2014).

PCR has been considered as an analytically sensitive and specific diagnostic method for the identification of *Helicobacter* in human gastric biopsies (Ndip et al., 2003; Ricci et al., 2007). Moreover, real-time (RT)-PCR allows quantitation of bacteria in gastric biopsies (He et al., 2002). To identify *H. suis* in the gastric biopsy samples from the study performed in **chapter 1**, a *H. suis*-specific *ureA*-based SYBR green RT-PCR was used. For this latter molecular assay, additional sequencing of the positive amplicons was necessary, since cross-reactivity with host DNA and DNA from other gastric helicobacters occurred, questioning the biological specificity of this test. In SYBR green RT-PCR assays, a signal will be emitted when the fluorophore binds to any double stranded DNA (Valones et al., 2008). In our assay, aspecific binding of the *ureA*-based primers with other DNA than the *H. suis* target DNA will thus result in a fluorescent signal and in false-positive results. Furthermore, UreA is essential for all gastric helicobacters to resist the acidic environment and to colonize the stomach. This virulence factor is, based on its high sequence level identity among the different gastric helicobacters, well conserved (Smet et al., unpublished results) explaining the occurrence of cross-reactivity with DNA from closely related members in our assay. It can thus be concluded that the *ureA* gene is actually not the optimal candidate target for molecular assays in gastric *Helicobacter* diagnostics if identification to species level is required.

Recently, a new PCR has been developed, using primer pairs targeting the *H. suis carR* gene (Matsui et al., 2014). Based on BLASTN searches in the NCBI database, this gene is absent in other gastric and enterohepatic *Helicobacter* spp., *Campylobacter* spp. and several other bacteria including members of the *Enterobacteriaceae* family and might thus be, compared to *ureA*, a better candidate PCR amplicon for the identification of *H. suis*

by reducing the possibility for cross-reactivity with DNA from other bacteria (Matsui et al., 2014). Sensitivity and specificity of this *carR*-based molecular assay was at the onset of this thesis lacking and determination of both parameters is essential for its introduction into routine clinical use. Therefore, in the **third chapter** of this thesis, we developed a *carR* probe-based RT-PCR that allows detection and quantification of *H. suis* in gastric tissue samples. The introduction of a *H. suis*-specific *carR* probe increased the specificity by excluding cross-reactivity with host DNA and DNA from other bacteria. Such probe will bind to a specific region within the single-stranded DNA. In contrast to SYBR green assays, in probe-based assays fluorescence will only be emitted when the DNA polymerase separates the reported dye from the quencher during amplification, rendering the probe-based RT-PCRs more specific (Valones et al., 2008). Our assay validation showed a high degree of diagnostic specificity and sensitivity compared to the *ureA*-based SYBR green RT-PCR. However, further application on large numbers of clinical samples is recommended. The optimization of such assay is a major progress in *H. suis* diagnostics in humans and similar assays should be developed for the other zoonotic NHPH species as well. Screening and comparison of the recently available NHPH genome sequences (Vermoote et al., 2011; Schott et al., 2011; Arnold et al., 2011; Smet et al., 2013; Smet et al., unpublished results) might facilitate the search for and identification of potential *Helicobacter* species-specific genes. Subsequently, PCR-based molecular diagnostic tools based on the amplification of specific regions within these potential *Helicobacter* species-specific genes may be developed.

Another method to identify gastric *Helicobacter* infections in gastric biopsies is isolation and cultivation of *Helicobacter*. For *H. pylori*, the success rate for cultivation reaches up to 100% (Ndip et al., 2003), but it depends on the quality, the quantity and the transportation conditions of the gastric biopsies (Ricci et al., 2007). Because of the high specificity and sensitivity, cultivation is considered as the gold standard for the diagnosis of *H. pylori* infections (Kusters et al., 2006). For NHPH, however, this method has not yet been successful, with *H. felis* and *H. bizzozeronii* being the only 2 species cultivated so far from the human stomach (Kivisto et al., 2010; Wüppenhorst et al., 2013). This can mainly be attributed to their very fastidious nature (Haesebrouck et al., 2009). For *H. suis*, it requires a highly enriched biphasic medium at pH 5 and a microaerobic atmosphere (Baele et al., 2008). Isolation of this agent from the gastric mucosa of pigs is possible but is a long and laborious process due to its slow growth. The risk of overgrowth with other bacteria makes frequent subculturing necessary in order to obtain pure cultures and it usually takes several weeks to obtain a single *H. suis* isolate (Baele et al., 2008). Efforts to isolate *H. suis* from human gastric biopsies, either directly (unpublished results) or after mouse-passage (Matsui et al., 2014), remained unsuccessful. The low colonization density of this pathogenic agent in the human stomach (Matsui et al., 2014) as well as the relative fast overgrowth of the isolation plates with other members of the gastric microbiome, including *Lactobacillus* spp., have no doubt contributed to this failure (Sgouras et al., 2004; Vilaichone et al., 2002; Chenoll et al., 2011; Matsui et al., 2014). Furthermore, previous studies have described the inhibitory effects on the growth and colonization capacity of *H. pylori* by lactobacilli and bifidobacteria in the stomach (Vilaichone et al., 2002; Sgouras et al., 2004; Francavilla et al., 2008; Medouakh et al., 2010; Aiba et al., 2015; Chenoll et al., 2011). This may also be the case for *H. suis in vitro*. To overcome these obstacles, we improved the mouse-passage-based protocol for the isolation of low numbers of *H. suis* bacteria. Mouse-passage of *H. suis*-infected gastric biopsies

might allow *H. suis* to multiply and to increase in numbers which might increase the chances for successful isolation. In our study, *H. suis* was successfully re-isolated from the stomach of mice on a biphasic medium, specified for this bacterium as described before (Baele et al., 2008), and additionally supplemented with 5 µg/ml linezolid. This antimicrobial agent is active against a broad spectrum of Gram-positive bacteria, including many *Lactobacillus* species (Yagi and Zurenko, 1997; Agar and Gold, 2012), and has a relatively low activity (MIC₉₀ 8µg/ml) against *Helicobacter* (Sanchez et al., 2000). Indeed, in the presence of linezolid, *H. suis* was able to grow whereas lactobacilli proliferation was clearly inhibited. This optimized isolation protocol can now be applied to isolate *H. suis* from human gastric biopsies. A possible drawback is that mice should be inoculated with homogenized gastric tissue containing 10⁴ bacteria/ml. Indeed, in our study, the minimal infection dose of *H. suis* in mice was 10⁴ viable *H. suis* bacteria/ml. The same infection dose has been described before and has also been reported for *H. pylori* infection in rhesus monkeys (Matsui et al., 2014; Solnick et al., 2001). Nevertheless, due to the low colonization density and patchy distribution of *H. suis* in the human stomach, this bacterium can still be missed when collecting biopsies. Pooling gastric biopsies obtained from several regions in the stomach or taking gastric brushes might counter this problem.

Isolation of human-derived *H. suis* would be a major breakthrough in *H. suis*-related research. First, the availability of such isolates would allow *in vitro* antimicrobial susceptibility testing. Because the increasing development of antibiotic resistance in *Helicobacter* species has been suggested (Van de Bulck et al., 2005; Kusters et al., 2001; Megraud et al., 2004; Megraud et al., 2007; Vermoote et al., 2011; Thung et al., 2016; Regnath et al., 2016), antimicrobial susceptibility testing of human *H. suis* isolates would allow to determine an appropriate patient-specific eradication therapy, thereby minimizing the occurrence of treatment failures and antibiotic resistance (Thung et al., 2016). This has already been applied to eradicate *H. pylori* infections in humans (Draeger et al., 2015). DNA prepared from pure human-derived *H. suis* isolates could also be used to further characterize this agent and to identify the origin and the evolution of these strains by comparing their genomes with those from porcine and non-human primate *H. suis* strains (Vermoote et al., 2011; Liang et al., 2015; Joosten et al., 2013; Bosschem et al., 2016; Flahou et al., unpublished data). Finally, to study the pathogenesis of gastric *H. suis* infections, experimental infection studies are indispensable. Such *in vivo* trials have already been performed in rodent models using porcine and non-human primate strains (Flahou et al., 2010; Bosschem et al., 2016). Also infection studies with human-derived *H. suis* have been carried out in the past (O'Rourke et al., 2004; Nakamura et al., 2007; Park et al., 2008; Matsui et al., 2014). However, in those mouse experiments, homogenized human gastric tissue was used as inoculum. This implies that other microorganisms were inoculated together with *H. suis*, which might have influenced the results (Flahou et al., 2010). Infection of laboratory animals with pure human-derived *H. suis* isolates might therefore provide a more accurate image of the pathogenesis of infections with this fastidious bacterium (O'Rourke et al., 2004; Nakamura et al., 2007; Flahou et al., 2010; Matsui et al., 2014).

Taken together, the developed methods may be implemented for diagnosis of *H. suis* in human-derived gastric biopsies. The probe-based RT-PCR can be useful for the detection of *H. suis* bacteria in the human stomach, whereas the application of the optimized mouse-passage-based isolation protocol might enhance the chance to obtain and characterize human *H. suis* isolates and to study their pathogenesis *in vivo*.

Final conclusions and future perspectives

In conclusion, we observed an increased frequency of gastric *H. suis* infections in human patients with PD, suggesting a possible link between *H. suis* infection and PD. This was further substantiated by demonstrating that an infection with this microorganism may influence the stomach-brain axis via the systemic circulation and may affect VIP signaling. Furthermore, the availability of optimized and accurate diagnostic tools to identify *H. suis* infections opens new perspectives for the study of its true prevalence in the human population.

It should be further investigated whether the observed effects of a gastric *H. suis* infection on the brain barriers will sensitize to or enhance the development of PD. This can be determined by the use of different *in vivo* models. PD can be mimicked in mice by intracranial injection of the neurotoxin 6-hydroxydopamine or 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridin (MPTP) or by using A30P α -synuclein transgenic mouse models, overexpressing the human A30P α -synuclein gene (Godoy et al., 2008; Plaas et al., 2008; Antony et al., 2011). However, both 6-hydroxydopamine and MPTP have been reported to cause impairment of the BBB (Carvey et al., 2005; Chen et al., 2008), suggesting that the use of the A30P α -synuclein transgenic mouse models is more reliable for the evaluation of the effect of *H. suis* on the course of PD. On the other hand, Sampson et al. (2016) recently described the importance of the gut microbiota for the induction of motor deficits, microglia activation and α -synuclein pathology in α -synuclein overexpressing transgenic mouse. Indeed, antibiotic treatment improved pathophysiology in adult animals, whereas the pathophysiology was increased by microbial re-colonization of the animals.

It might also be interesting to perform similar experimental trials to evaluate the influence of *H. suis* infections on development and/or progression of other neurodegenerative disorders. For example, the influence of *H. suis* on Alzheimer's disease, the most common neurodegenerative disorder in the current human population (Wirdefelt et al., 2011), may be studied by using rodent models. Alzheimer's disease can be mimicked by intrahippocampal injection with the neurotoxin amyloid β , which induces the formation of A β -plaques in the brain, or by using APP-PS1 transgenic mouse models, overexpressing human amyloid β (Lok et al., 2013; Jean et al., 2015).

Besides studying the integrity of the stomach-brain barriers, loss of dopaminergic neurons in the substantia nigra of the brain should be investigated. This can be done by using tyrosine hydroxylase staining with or without a silver nucleolar staining technique (Healy-stoffel et al., 2012). Additionally, the presence of lewy bodies in the remaining neurons can be determined by immunohistochemical stainings such as the α -synuclein immunohistochemical staining (Stefanis, 2012) or the ubiquitin immunohistochemical staining (Chu et al., 2000). Also, expression of genes associated with PD, such as *α -synuclein*, *parkin (PARK)*, *phosphatase and tensin homolog (PTEN)-induced putative kinase 1 (PINK)* and *mitochondrial ribosomal protein S6 (MRPS6)* genes, can be measured in the brain (Lewis et al., 2012).

Apart from impairment of the stomach-brain axis resulting in systemic inflammation and dysregulation of the VIP expression, changes in the intestinal microbiome might also play a role in development of PD and/or influence its course. Several studies have indeed described a possible link between such changes and brain

disorders, including depression, autism, Alzheimer's disease and PD (Parracho et al., 2005; Bercik et al., 2011; Bravo et al., 2011; Zhang et al., 2015; Hu et al., 2016; Mulak et al., 2015). Changes in the gut microbiome may be caused by a broad range of triggers, including gastrointestinal infections (Ochoa-Reparaz and Kasper, 2014; Walters et al., 2014; Zhang et al., 2015). *H. pylori* infections have already been shown to affect the gastrointestinal microbiota in humans (Engstrand et al., 2013; Budzynski et al., 2014; He et al., 2016). Therefore, it might be recommended to evaluate the effect of an *H. suis* infection on the gut microbiome, for instance in experimentally infected mice and in *H. suis* infected human patients, before and after eradication of the infection.

Additionally, it might be recommended to evaluate whether the induced changes in brain homeostasis are directly due to the actual infection with *H. suis*, or rather to possible *H. suis*-induced changes in the gut microbiome. Intestinal microbiota transplantation from a *H. suis* infected mice to the intestinal tract of a control recipient in order to change the recipient's gut microbial composition might be useful here (Gupta et al., 2016). If it appears that the shift in the composition of the microbiota is associated with changes in brain homeostasis, intestinal microbiota transplantation from a healthy person to a patient suffering from a neurodegenerative disorder might be a potential therapeutic strategy. Indeed, intestinal microbial transplantation has already been suggested as a possible therapy for treatment of IBD, diarrhea, *Clostridium difficile* infections, etc. (Gupta et al., 2016).

Our results indicate that the administration of VIP might play a protective role in inflammation-induced disruption of the blood-CSF barrier integrity and brain homeostasis. The immunomodulatory and neuroprotective effect of VIP has already been suggested in the past (Kim et al., 2000; Delgado and Ganea, 2003; Delgado et al., 2004). Moreover, a protective effect of VIP in PD and Alzheimer's disease has been described (Delgado and Ganea, 2003; Song et al., 2012). Therefore, the administration of VIP might be another possible therapeutic strategy in patients suffering from PD or other neurodegenerative disorders.

References

- Abbott RD, Petrovitch H, White LR, Masaki KH, Tanner CM, Curb JD, Grandinetti A, Blanchette PL, Popper JS, Ross GW. 2001. Frequency of bowel movements and the future risk of Parkinson's disease. *Neurology* 5: 456-462.
- Agar S, Gould K. 2012. Clinical update on linezolid in the treatment of gram-positive bacterial infections. *Inf and drug resist* 5: 87-102.
- Aiba Y, Nakano Y, Koga Y, Takahashi K, Komatsu Y. 2015. A highly acid resistant novel strain of *Lactobacillus johnsonii* No. 1088 has antibacterial activity, including that against *Helicobacter pylori*, and inhibits gastrin-mediated acid production in mice. *MicrobiologyOpen* 4(3): 465-474.
- Alvarez-Arellano L, Maldonado-Bernal C. 2014. *Helicobacter pylori* and neurological diseases: married by the laws of inflammation. *World J Gastroenterol* 5(4): 400-404.
- Anlauf M, Schäfer MK, Eiden L, Weihe E. 2003. Chemical coding of the human gastrointestinal nervous system: cholinergic, VIPergic, and catecholaminergic phenotypes. *J Comp Neurol*. 459(1): 90-111.
- Annibale B, Di Giulio E, Caruana P, Lahner E, Capurso G, Bordi C, Delle Fave G. 2002. The long-term effects of cure of *Helicobacter pylori* infection on patients with atrophic body gastritis. *Alim Pharmacol and Ther* 16: 1723-1731.
- Antony PMA, Diederich NJ, Balling R. 2011. Parkinson's disease mouse models in translational research. *Mamm Genome*. 22(7-8): 401-419.
- Appelmek BJ, Negrini R, Moran AP, Kuipers EJ. 1997. Molecular mimicry between *Helicobacter pylori* and the host. *Trends Microbiol* 5: 70-73.
- Aranda V, Haire T, Nolan ME, Calarco JP, Rosenberg AZ, Fawcett JP, Pawson T, Muthuswamy SK. 2006. Par6-aPKC uncouples ErbB2 induced disruption of polarized epithelial organization from proliferation control. *Nat Cell Biol*.8(11): 1235-1245.
- Arnold IC, Zigova Z, Holden M, Lawley TD, Rad R, Dougan G, Falkow S, Bentley DS, Müller A. 2011. Comparative whole genome sequence analysis of the carcinogenic bacterial model pathogen *Helicobacter felis*. *Genome Biol Evol* 3: 302-308.
- Baele M, Van den Bulck K, Decostere A, Vandamme P, Hänninen ML, Ducatelle R, Haesebrouck F. 2004: Multiplex PCR assay for differentiation of *Helicobacter felis*, *H. bizzozeronii* and *H. salomonis*. *J Clin Microbiol*. 42: 1115-1122.
- Baele M, Decostere A, Vandamme P, Ceelen L, Hellemans A, Mast J, Chiers K, Ducatelle R, Haesebrouck F. 2008: Isolation and characterization of *Helicobacter suis* sp. nov. from pig stomach. *Int J Syst Evol Microbiol*. 58(Pt 6): 1350-8
- Bagheri N, Azadegan-Dehkordi F, Shirzad H, Rafiean-Kopaei M, Rahimian G, Razavi A. 2015. The biological functions of IL-17 in different clinical expressions of *Helicobacter pylori* infection. *Microbial Pathogenesis* 81: 33-38.
- Bercik P, De Giorgio R, Blennerhassett P, Verdú EF, Barbara G, Collins SM. 2002. Immune-mediated neural dysfunction in a murine model of chronic *Helicobacter pylori* infection. *Gastroenterol* 123: 1205-1215.
- Bercik P, Denou E, Collins J, Jackson W, Lu J, Jury J, Deng Y, Blennerhassett P, Macri J, McCoy KD, Verdu EF, Collins SM. 2011. The intestinal microbiota affect central levels of brain-derived neurotrophic factor and behavior in mice. *Gastroenterology* 141, 599-609, 609 e591-593.
- Bosschem I, Flahou B, Bakker J, Heuvelman E, Langermans JAM, De Bruyne E, Joosten M, Smet A, Ducatelle R, Haesebrouck F. 2016. Comparative virulence of *in vitro* cultures primate- and pig-associated *Helicobacter suis* strains in a BALB/c mouse and Mongolian gerbil model. *Helicobacter*, in press
- Bosschem I, Flahou B, Van Deun K, De Koker S, Volf J, Smet A, Ducatelle R, Devriendt B, Haesebrouck F. 2016. Species-specific immunity to *Helicobacter suis*. *Helicobacter*, in press.
- Bravo JA, Forsythe P, Chew MV, Escaravage E, Savignac HM, Dinan TG, Bienenstock J, Cryan JF. 2011. Ingestion of *Lactobacillus* strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proc Natl Acad Sci U S A* 108, 16050-16055
- Budzynski J, Klopocka M. 2014. Brain –gut axis in the pathogenesis of *Helicobacter pylori* infection. *World Journal Gastroenterol* 20(18): 5212-5225.
- Carethers JM. 2011. One colon lumen but two organs. *Gastroenterology* 141(2): 411-412.

- Caron TJ, Scott KJ, Fox JG, Hagen SJ. 2015. Tight junction disruption: *Helicobacter pylori* and dysregulation of the gastric mucosal barrier. *World J Gastroenterol*. 21(40): 11411–11427.
- Carvey PM, Zhao CH, Hendey B, Lum H, Trachtenberg J, Desai BS, Snyder J, Zhu YG, Ling ZD. 2005. 6-Hydroxydopamine-induced alterations in blood-brain barrier permeability. *Eur J Neurosci*. 22(5): 1158-1168.
- Cavaillon JM, Haeffner-Cavaillon N. 1986. Polymyxin-B inhibition of LPS-induced interleukin-1 secretion by human monocytes is dependent upon the LPS origin. *Mol Immunol*. 23(9): 965-9.
- Charlett A, Dobbs RJ, Weller C, Dobbs SM. 1997. Stasis in the gut: the source of xenobiotic in idiopathic parkinsonism. *Eur J Clin Pharmacol* 52 (Suppl): 168.
- Charlett A, Dobbs RJ, Purkiss AG, Wright, DJ, Peterson DW, Weller C, Dobbs SM. 1998. Cortisol is higher in parkinsonism and associated with gait deficit. *Acta Neurol Scand* 97: 77-85.
- Chen X, Lan X, Roche I, Liu R, Geiger JD. 2008. Caffeine protects against MPTP-induced blood-brain barrier dysfunction in mouse striatum, *J Neurochem*. 107(4): 1147–1157.
- Chen L, Yuan W, Chen Z, Wu S, Ge J, Chen J, Chen Z. 2015. Vasoactive intestinal peptide represses activation of tumor-associated macrophages in gastric cancer via regulation of TNF α , IL-6, IL-12 and iNos. *Internat Journal of Oncology* 47: 1360-1370.
- Chenoll E, Casinos B, Bataller E, Astals P, Echevarría J, Iglesias JR, Balbarie P, Ramón D, Genovés S. 2011. Novel probiotic *Bifidobacterium bifidum* CECT 7366 strain active against the pathogenic bacterium *Helicobacter pylori*. *Appl Environ Microbiol*. 2011 Feb;77(4): 1335-1343
- Chu CT, Caruso JL, Cummings TJ, Ervin J, Rosenberg C, Hulette CM. 2000. Ubiquitin immunochemistry as a diagnostic aid for community pathologists evaluating patients who have dementia. *Mod Pathol*. 13(4): 420-426.
- Conlin VS, Wu X, Nguyen C, Dai C, Vallance BA, Buchan AM, Boyer L, Jacobson K. 2009. Vasoactive intestinal peptide ameliorates intestinal barrier disruption associated with *Citrobacter rodentium*-induced colitis. *Am J Physiol Gastrointest Liver Physiol*. 297(4): G735-750
- Crabtree JE, Lindley IJ. 1994. Mucosal interleukin-8 and *Helicobacter pylori*-associated gastroduodenal disease. *Eur J Gastroenterol Hepatol*. 6(1): S33-38
- Cryan JF, Dinan TG. 2012. Mind altering microorganisms: the impact of the gut microbiota on the brain and behavior. *Nat Rev Neurosci*. 13(10): 701-12
- Csoti I, Jost WH, Reichmann H. 2016. Parkinson's disease between internal medicine and neurology. *J Neural Transm* 123: 3-17.
- De Bock M, Vandenbroucke RE, Decrock E, Culot M, Cecchelli R, Leybaert L. 2014. A new angle on blood-CNS interfaces: a role for connexins? *FEBS Lett* 588:1259-1270.
- Debongnie JC, Donnay M, Mairesse J. 1995. *Gastrospirillum hominis* (*Helicobacter heilmannii*): a cause of gastritis, sometimes transient, better diagnosed by touch of cytology. *Am J Gastroenterol*. 90(3): 411-6
- De Cooman L, Flahou B, Houf K, Smet A, Ducatelle R, Pasmans F, Haesebrouck F. 2013. Survival of *Helicobacter suis* bacteria in retail pig meat. *Int. J. Food Microbiol*. 166(1): 164-167.
- De Groote D, Haesebrouck F, van Doorn LJ, Vandamme P, Ducatelle R. 2001: Evaluation of a group-specific 16S ribosomal DNA-based PCR for detection of *Helicobacter bizzozeroni*, *Helicobacter felis* and *Helicobacter salomonis* in fresh and paraffin-embedded gastric biopsy specimens. *J Clin Microbiol* 39: 1197-1199.
- De Groote D, Van Doorn LJ, Van den Bulck K, Vandamme P, Vieth M, Stolte M, Debongnie JC, Burette A, Haesebrouck F, Ducatelle R. 2005: Detection of non-*Helicobacter pylori* *Helicobacter* species in "*Helicobacter heilmannii*"-infected humans. *Helicobacter* 10(5): 398-406.
- Delgado M, Ganea D. 2003. Vasoactive intestinal peptide prevents activated microglia-induced neurodegeneration under inflammatory conditions: potential therapeutic role in brain trauma. *FASEB J*. 2003 Oct;17(13): 1922-1924
- Demeestere D, Libert C, Vandenbroucke RE. 2015. Clinical implications of leukocyte infiltration at the choroid plexus in (neuro)inflammatory disorders. *Drug Discovery Today* 20: 928-941.
- De Vadder F, Plessier F, Gautier-Stein A, Mithieux G. 2015. Vasoactive intestinal peptide is a local mediator in a gut-brain neural axis activating intestinal gluconeogenesis. *Neurogastroenterol Motil*. 27(3): 443-448

- Dobbs RJ, Charlett A, Purkiss AG, Dobbs SM, Weller C, Peterson DW. 1999. Association of circulating TNF- α and IL-6 with ageing and parkinsonism. *Acta Neurol Scand* 100: 34-41.
- Dobbs SM, Dobbs RJ, Weller C, Charlett A. 2000. Link between *Helicobacter pylori* infection and idiopathic parkinsonism. *Medical Hypothesis* 55(2): 93-98.
- Dobbs RJ, Dobbs SM, Weller C, Bjarnason IT, Oxlade NL, Charlett A, Al-Janabi MA, Kerwin RW, Mahler RF, Price AB. 2005. Role of Chronic Infection and Inflammation in the Gastrointestinal Tract in the Aetiology and Pathogenesis of Idiopathic Parkinsonism. Part 1: eradication of *Helicobacter* in the cachexia of Idiopathic Parkinsonism. *Helicobacter* 10: 267–275.
- Dobbs RJ, Dobbs SM, Weller C, Charlett A, Bjarnason IT, Curry A, Ellis DS, Ibrahim MA, McCrossan MV, O'Donohue J, Owen RJ, Oxlade NL, Price AB, Sanderson JD, Sudhanva M, Williams J. 2008 *Helicobacter* hypothesis for IP: before and beyond. *Helicobacter* 13(5): 309-322.
- Dobbs SM, Dobbs RJ, Weller C, Charlett A, Bjarnason IT, Lawson AJ, Letley D, Harbin L, Price AB, Ibrahim MA, Oxlade NL, Bowthorpe J, Leckstroem D, Smee C, Plant JM, Peterson DW. 2010: Differential effect of *Helicobacter pylori* eradication on time trends in brady/hypokinesia and rigidity in Idiopathic Parkinsonism. *Helicobacter* 15: 279-294.
- Dobbs RJ, Charlett A, Dobbs SM, Weller C, Ibrahim MAA, Iguodala O, Smee C, Plant JM, Lawson AJ, Taylor D, Bjarnason I. 2012. Leukocyte-subset counts in idiopathic parkinsonism provide clues to a pathogenic pathway involving small intestinal bacterial overgrowth. A surveillance study. *Gut Pathogens* 4: 12.
- Dobbs SM, Charlett A, Dobbs RJ, Weller C, Iguodala O, Smee C, Lawson AJ, Taylor D, Bjarnason I. 2013: Antimicrobial surveillance in idiopathic parkinsonism : indication –specific improvement in hypokinesia following *Helicobacter pylori* eradication and non –specific effect of antimicrobials for other indications in worsening rigidity. *Helicobacter* 18: 187-196.
- Dobbs SM, Dobbs RJ, Weller C, Charlett A, Augustin A, Taylor D, Ibrahim MA, Bjarnason I. 2016. Peripheral aetiopathogenic drivers and mediators of Parkinson's disease and co-morbidities: role of gastrointestinal microbiota. *J. Neuroviol* 22(1): 22-32.
- Doherty NC, Tobias A, Watson S, Atherton JC. 2009. The effect of gut-signalling hormone, norepinephrine on the growth of the gastric pathogen *Helicobacter pylori*. *Helicobacter* 14: 223-230.
- Draeger S, Wüppenhorst N, Kist M, Glocker EO. 2015. Outcome of second- and third-line *Helicobacter pylori* eradication therapies based on antimicrobial susceptibility testing. *J Antimicrob Chemother.* 70(11): 3141-3145.
- Edwards LL, Quigley EM, Pfeiffer RF. 1992. Gastrointestinal dysfunction in Parkinson's disease. *Neurology* 42: 726-732.
- Engstrand L, Lindberg M. 2013. *Helicobacter pylori* and the gastric microbiota. *Best Pract Res Clin Gastroenterol.* 27: 39–45.
- Erin N, Türker S, Elpek O, Yildirim B. 2012. Differential changes in Substance P, VIP as well as neuropeptide Y levels in patients with gastritis or ulcer. *Peptides* 35: 218-224.
- Fedwick JP, Lapointe TK, Meddings JB, Sherman PM, Buret AG. 2005. *Helicobacter pylori* Activates Myosin Light-Chain Kinase To Disrupt Claudin-4 and Claudin-5 and Increase Epithelial Permeability. *Infect Immun.* 73(12): 7844–7852
- Ferrari CC, Tarelli R. 2011. Parkinson's disease and systemic inflammation. *Parkinson's Disease* doi:10.4061/2011/436813.
- Fiszer U, Fredrikson S, Czlonkowska A. 1996. Humoral response to hsp65 and hsp 70 in cerebrospinal fluid in Parkinson's disease. *J Neurol Sci* 139; 66-70.
- Flahou B, Haesebrouck F, Pasmans F, D'Herde K, Driessen A, Van Deun K, Smet A, Duchateau L, Chiers K, Ducatelle R. 2010. *Helicobacter suis* causes severe gastric pathology in mouse and Mongolian gerbil models of human gastric disease. *Plos one.* 5(11): e14083.
- Flahou B, Haesebrouck F, Chiers K, Van Deun K, De Smet L, Devreese B, Vandenberghe I, Favoreel H, Smet A, Pasmans F, D'Herde K, Ducatelle R. 2011. Gastric epithelial cell death caused by *Helicobacter suis* and *Helicobacter pylori* γ -glutamyl transpeptidase is mainly glutathione degradation-dependent. *Cell Microbiol* 13(12): 1933-1955
- Flahou B, Van Deun K, Pasmans F, Smet A, Volf J, Rychlik I, Ducatelle R, Haesebrouck F. 2012. The local immune response of mice after *Helicobacter suis* infection: strain differences and distinction with *H. pylori*. *Vet Research* 43: 75.
- Flahou B, Haesebrouck F, Smet A. 2016. Non-*Helicobacter pylori* *Helicobacter* infections in humans and animals. Book: *Helicobacter pylori* research: from bench to bedside, Springer Japan, in press.
- Fonseca ACC, Matias D, Garcia C, Amaral R, Geraldo LH, Freitas C, Lima FRS. 2014. The impact of microglial activation on blood-brain barrier in brain diseases. *Front Cell Neurosci.* 8: 362.

- Francavilla R, Lionetti E, Castellaneta SP, Magistà AM, Maurogiovanni G, Bucci N, De Canio A, Indrio F, Cavallo L, Ierardi E, Miniello VL . 2008. Inhibition of *Helicobacter pylori* infection in humans by *Lactobacillus reuteri* ATCC 55730 and effect on eradication therapy: a pilot study. *Helicobacter* 13: 127-134.
- Fukuda Y, Bamba H, Okui M, Tamura K, Tanida N, Satomi M, Shimoyama T, Nishigami T. 2001. *Helicobacter pylori* infection increases mucosal permeability of the stomach and the intestine. *Digestion* 63(1): 93-96.
- Godoy MCP, Tarelli R, Ferrari CC, Sarchi MI, Pitossi FJ. 2008. Central and systemic IL-1 exacerbates neurodegeneration and motor symptoms in a model of Parkinson's disease. *Brain* 131: 1880-1894.
- Gorell JM, Johnson CC, Rybicki BA, Peterson EL, Richardson RJ. 1998. The risk of Parkinson's disease with exposure to pesticides, farming, well water, and rural living. *Neurology*. 50(5): 1346-1350
- Guo H, Callaway JB, Ting JP-Y. 2015. Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nature medicine* 21 (7): 677-687.
- Gupta S, Allen-Versoe E, Petrof EO. 2016. Faecal microbiota transplantation: in perspective. *Therap Adv Gastroenterol*. 9(2): 229-239.
- Guttman JA , Finlay BB. 2009. Tight junctions as targets of infectious agents. *Biochimica et Biophysica Acta* 1788: 832-841.
- Haesebrouck F, Pasmans F, Flahou B, Chiers K, Baele M, Meyns T, Decostere A, Ducatelle R. 2009. Gastric helicobacters in domesticated animals and non-human primates and their significance for human health. *Clin Microbiol Rev* 22: 202-223.
- Hardy J, Cai H, Cookson MR, Gwinn-Hardy K, Singleton A. 2006. Genetics of Parkinson's disease and parkinsonism. *Ann Neurol* 60(4): 389-398.
- Hashim H, Azmin S, Razlan H Yahya NW, Tan HJ, Abdul Manaf MR, Ibrahim NM. 2014. Eradication of *Helicobacter pylori* Infection Improves Levodopa Action, Clinical Symptoms and Quality of Life in Patients with Parkinson's Disease. *PLoS One*. 9(11): e112330.
- Hasni S, Ippolito A, Illei GG. 2011. *Helicobacter pylori* and autoimmune diseases. *Oral Dis* 17(7): 621-627.
- He Q, Wang JP, Osato M, Lachman LB. 2002. Real-time quantitative PCR for detection of *Helicobacter pylori*. *J Clin Microbiol*. 40(10): 3720-3728.
- He C, Yang Z, Lu N. 2016. Imbalance of Gastrointestinal Microbiota in the Pathogenesis of *Helicobacter pylori*-Associated Diseases. *Helicobacter* 21 (5): 337–34.
- Healy-Stoffel M, Ahmad OS, Stanford JA, Levant B. 2012. A novel use of combined tyrosine hydroxylase and silver nucleolar staining to determine the effects of a unilateral intrastriatal lesion in the substantia nigra: a stereological study. *J. Neurosci Methods* 210 (1): 187-194.
- Hellemans A, Decostere A, Haesebrouck F, Ducatelle R. 2005. Evaluation of Antibiotic Treatment against "*Candidatus Helicobacter suis*" in a Mouse Model. *Antimicrob Agents Chemother*. 49(11): 4530–4535.
- Hirsch EC, Hunot S. 2009. Neuroinflammation in Parkinson's disease: a target for neuroprotection? *Lancet Neurol*. 8(4): 382-397.
- Hoffmann A, Bredno J, Wendland M, Derugin N, Ohara P, Wintermark M. 2011. High and Low Molecular Weight Fluorescein Isothiocyanate (FITC)-Dextran to Assess Blood-Brain Barrier Disruption: Technical Considerations. *Transl Stroke Res*. 2(1): 106–111.
- Hu X, Wang T, Jin F. 2016. Alzheimer's disease and gut microbiota. *Sci China Life Sci*, doi:10.1007/s11427-016-5083-9.
- Islek A, Yilmaz A, Elpek GO, Erin N. 2016. Childhood chronic gastritis and duodenitis: Role of altered sensory neuromediators. *World Journal Gastroenterol*. 22(37): 8349-8360.
- Ito T, Kobayashi D, Uchida K, Takemura T, Nagaoka S, Kobayashi I, Yokoyama T, Ishige I, Ishige Y, Ishida N, Furukawa A, Muraoka H, Ikeda S, Sekine M, Ando N, Suzuki Y, Yamada T, Suzuki T, Eishi Y. 2008. *Helicobacter pylori* invades the gastric mucosa and translocates to the gastric lymph nodes. *Lab Invest*. 88(6): 664-681.
- Joosten M, Flahou B, Meyns T, Smet A, Arts J, De Cooman L, Pasmans F, Ducatelle R, Haesebrouck F. 2013: Case report: *Helicobacter suis* infection in a pig veterinarian. *Helicobacter* 18: 392-396
- Joosten M, Blaecher C, Flahou B, Ducatelle R, Haesebrouck F, Smet A. 2013. Diversity in bacterium-host interactions within the species *Helicobacter heilmannii sensu stricto*. *Vet Res* 44: 65
- Kabir S. 2011. The role of interleukin-17 in the *Helicobacter pylori* induced infection and immunity. *Helicobacter* 16: 1-8.

- Kannarat GT, Boss JM, Tansey MG. 2013. The role of innate and adaptive immunity in Parkinson's disease. *J. Parkinson's Dis.* 3(4): 493-514.
- Kang JM, Kim M, Lee BH, Park HK, Jo HJ, Shin CM, Lee SH, Song IS. 2011. Risk factors for peptic ulcer bleeding in terms of *Helicobacter pylori*, NSAID's, and antiplatelet agents. *Scand J Gastroenterol* 46(11): 1295-1301.
- Kim WK, Kan Y, Ganea D, Hart RP, Gozes I, Jonakait GM. 2000. Vasoactive Intestinal Peptide and Pituitary Adenylate Cyclase-Activating Polypeptide Inhibit Tumor Necrosis Factor Production in Injured Spinal Cord and in Activated Microglia via a cAMP Dependent Pathway. *The Journal of Neuroscience* 20(10): 3622-3630
- Kim H, Lim JW, Kim KH. 2001. *Helicobacter pylori*-induced expression of interleukin-8 and Cyclooxygenase-2 in AGS gastric epithelial cells: mediation by nuclear factor- κ B. *Scand J Gastroenterol* 36: 706-716.
- Kivisto R, Linros J, Rossi M, Rautelin H, Hänninen ML. 2010: Characterization of multiple *Helicobacter bizzozeroni* isolates from a Finnish patient with severe dyspeptic symptoms and chronic active gastritis. *Helicobacter* 15: 58-66
- Kountouras J, Zavos C, Polyzos SA, Deretzi G, Vardaka E, Giartza-Taxidou E, Katsinelos E, Rapti D, Chatzopoulos D, Tzilves C. 2012. *Helicobacter pylori* infection and Parkinson's disease: apoptosis as an underlying common contributor. *Eur J Neurol* 19: e56.
- Kountouras J, Zavos C, Polyzos SA, Deretzi G. 2015. The gut-brain axis: interactions between *Helicobacter pylori* and enteric and central nervous systems. *Ann Gastroenterol.* 28(4): 506.
- Kortekaas R, Leenders KL, van Oostrom JC, Vaalburg W, Bart J, Willemsen AT, Hendrikse NH. 2005. Blood-brain barrier dysfunction in parkinsonian midbrain *in vivo*. *Ann Neurol.* 57(2): 176-179.
- Kusters and Kuipers, 2001: Antibiotic resistance of *Helicobacter pylori*. *Journal of applied microbial* 90: 1345-1445
- Kusters JG, Van Vliet AHM, Kuipers EJ. 2006. Pathogenesis of *Helicobacter pylori* infection. *Clin Microbiol Rev* 19: 449-490.
- Lee E, Burnett CA, Lulich N, Cameron LL, Sestito JP. 2002. Proportionate mortality of crop and livestock farmers in the United States, 1984-1993. *Am J Ind Med* 42(5): 410-420.
- Lee H, Pienaar IS. 2014. Disruption of the blood-brain barrier in Parkinson's disease: curse or route to cure? *Front Biosci (Landmark Ed)* 19: 272-280.
- Lewis PA, Mark R. Cookson MR. 2012. Gene expression in the Parkinson's disease brain. *Brain Res Bull.* 88(4): 302-312.
- Li XB, Chen HM, Lu H, Zheng Q, Chen XY, Peng YS, Ge ZZ, Liu WZ. 2009. Role of *Helicobacter pylori* infection on neuronal expression in the stomach and the spinal cord of a murine model. *J Dig Dis* 10: 286-292.
- Liang J, Ducatelle R, Pasmans F, Smet A, Haesebrouck F, Flahou B. 2013: Multilocus sequence typing of the porcine and human gastric pathogen *Helicobacter suis*. *J Clin Microbiol* 51: 920-926.
- Liang J, De Bruyne E, Ducatelle R, Smet A, Haesebrouck F, Flahou B. 2015. Purification of *Helicobacter suis* Strains From Biphasic Cultures by Single Colony Isolation: Influence on Strain Characteristics. *Helicobacter.* 20(3): 206-216
- Lindvall M, Alumets J, Edvinsson L, Fahrenkrug J, Håkanson R, Hånko J, Owman C, Schaffalitzky de Muckadell OB, Sundler F. 1978. Peptidergic (VIP) nerves in the mammalian choroid plexus. *Neurosci Lett.* 9(1): 77-82.
- Liu Q, Li Y, Zhao X, Yang X, Liu Q, Kong Q. 2015. Construction of *Escherichia coli* Mutant with Decreased Endotoxic Activity by Modifying Lipid A Structure. *Mar Drugs.* 13(6): 3388-406.
- Liu J, He L, Haesebrouck F, Gong Y, Flahou B, Cao Q, Zhang J. 2015. Prevalence of Coinfection with Gastric Non-*Helicobacter pylori* *Helicobacter* (NHPP) Species in *Helicobacter pylori*-infected Patients Suffering from Gastric Disease in Beijing, China. *Helicobacter.* 20(4): 284-290
- Liu C, Rossi M, Linden S, Medae P, Blaecher C, Bauwens E, Joosten M, Flahou B, Van den Broeck W, Ducatelle R, Haesebrouck F, Smet A. 2016. The *Helicobacter heilmannii* *hofE* and *hofF* Genes are Essential for Colonization of the Gastric Mucosa and Play a Role in IL-1 β -Induced Gastric MUC13 Expression. *Helicobacter* doi:10.1111/hel.12307
- Matsui H, Takahashi T, Murayama SY, Uchiyama I, Yamaguchi K, Shigenobu S, Matsumoto T, Kawakubo M, Horiuchi K, Ota H, Osaki T, Kamiya S, Smet A, Flahou B, Ducatelle R, Haesebrouck F, Takahashi S, Nakamura S, Nakamura M. 2014: Development of new PCR primers by comparative genomics for the detection of *Helicobacter suis* in gastric biopsy specimens. *Helicobacter* 19: 260-271

- Mayer EA, Knight R, Mazmanian SK, Cryan JF, Tillisch K. 2014. Gut microbes and the brain : Paradigm shift in neuroscience. *Journal of neuroscience* 34(46): 15490-15496.
- Medouakh L, Abdeslam AA, Bensoltane A. 2010. Antagonistic activity of *Lactobacillus* sp. against *Helicobacter pylori*. *Int J Micorbiol Res* 1(3): 80-86.
- Megraud F. 2004: *H. pylori* antibiotic resistance: prevalence, importance, and advances in testing. *Gut* 53: 1374-1384
- Megraud F, Lehours P. 2007: *Helicobacter pylori* detection and antimicrobial susceptibility testing. *Clin Microbiol Rev* 20: 280-322
- Mhyre TR, Boyd JT, Hamill RW, Maguire-Zeiss KA. 2012. Parkinson's disease. *Subcell Biochem* 65: 389-455.
- Miyaji H, Azuma T, Ito S, Abe Y, Ono H, Suto H, Ito Y, Yamazaki Y, Kohli Y, Kuriyama M. 1999. The effect of *Helicobacter pylori* eradication therapy on gastric antral myoelectrical activity and gastric emptying in patients with non-ulcer dyspepsia. *Aliment Pharmacol Ther* 13: 1303-1310.
- Montecucco C and Rappuoli R. 2001. Living dangerously: how *Helicobacter pylori* survives in the human stomach. *Nature Reviews Molecular Cell Biology* 2: 457-466
- Monzon H, Forne M, Esteve M, Rosinach M, Loras C, Espinos JC, Viver JM, Salas A, Fernandez-Banares F. 2013. *Helicobacter pylori* infection as a cause of iron deficiency anaemia of unknown origin. *World J Gastroenterol* 19(20): 4166-4171
- Mulak A, Bonaz B. 2015. Brain-gut-microbiota axis in Parkinson's disease. *World Journal Gastroenterol* 21(37): 10609-10620.
- Nakamura M, Murayama SY, Serizawa H, Sekiya Y, Eguchi M, Takahashi S, Nishikawa K, Takahashi T, Matsumoto T, Yamada H, Hibi T, Tsuchimoto K, Matsui H. 2007. "*Candidatus Helicobacter heilmannii*" from a Cynomolgus Monkey Induces Gastric Mucosa-Associated Lymphoid Tissue Lymphomas in C57BL/6 Mice". *Infect immun.* 75: 1214-1222.
- Ndip RN, MacKay WG, Farthing MJ, Weaver LT. 2003: Culturing *Helicobacter pylori* from clinical specimens: review of microbiological methods. *J Pediatr Gastroenterol Nutr.* 36: 616-622.
- Nielsen HH, Qiu J, Friis S, Wermuth L, Ritz B. 2012: Treatment for *Helicobacter pylori* infection and risk of Parkinson's disease in Denmark. *Eur J Neurol* 19: 864-869.
- Nilsson C, Lindvall-Axelsson M, Owman C. 1991. Simultaneous and continuous measurement of choroid plexus blood flow and cerebrospinal fluid production: effects of vasoactive intestinal polypeptide. *J Cereb Blood Flow Metab.* 11(5): 861-867.
- Ochoa-Reparaz, J. & Kasper, L. H. 2014 Gut microbiome and the risk factors in central nervous system autoimmunity. *FEBS Lett* 588, 4214-4222, doi:10.1016/j.febslet.2014.09.024.
- Odenbreit S, Wieland B, Haas R. 1996. Cloning and genetic characterization of *Helicobacter pylori* catalase and construction of a catalase-deficient mutant strain. *J Bacteriol.* 178(23): 6960-6967.
- O' Rourke JL, Solnick JV, Neilan BA, Seidel K, Hayter R, Hansen LM, Lee A. 2004: Description of '*Candidatus Helicobacter heilmannii*' based on DNA sequence analysis of the 16S rRNA and the urease genes. *J Pathol* 203: 896-903.
- Park JH, Seok SH, Baek MW, Lee HY, Kim DJ, Park JH. 2008: Gastric lesions and immune responses caused by long-term infections with *Helicobacter heilmannii* in C57BL/6 mice. *J Comp Pathol.* 139: 208-217
- Parracho, H. M., Bingham, M. O., Gibson, G. R. & McCartney, A. L. 2005. Differences between the gut microflora of children with autistic spectrum disorders and that of healthy children. *J Med Microbiol* 54, 987-991, doi:10.1099/jmm.0.46101-0.
- Peek RM JR, Fiske C and Wilson KT. 2010. Role of innate immunity in *Helicobacter pylori*-induced gastric malignancy. *Physiol Rev* 90: 831-858.
- Perry HV, Teeling J. 2013. Microglia and macrophages of the central nervous system: the contribution of microglia priming and systemic inflammation to chronic neurodegeneration. *Semin Immunopathol* 35: 601–612
- Pfeiffer RF. 2011. Gastrointestinal dysfunction in Parkinson's disease. *Parkinsonism Relat Disord* 17: 10-15.
- Pierantozzi M, Pietroiusti A, Sancesario G, Lunardi G, Fedele E, Giacomini P, Frasca S, Galante A, Marciari MG, Stanzione P. 2001. Reduced L-dopa absorption and increased clinical fluctuations in *Helicobacter pylori*-infected Parkinson's disease patients. *Neurol Sci* 22(1): 89-91.

- Pisani V, Stefani A, Pierantozzi M, Natoli S, Stanzione P, Franciotta D, Pisani A. 2012. Increased blood-cerebrospinal fluid transfer of albumin in advanced Parkinson's disease. *Journal of Neuroinflamm* 9: 188.
- Plaas M, Karis A, Innos J, Rebane E, Baekelandt V, Vaarmann A, Luuk H, Vasar E, Koks S. 2008. Alpha-synuclein A30P point-mutation generates age dependent nigrostriatal deficiency in mice. *Journal of Physiol and Pharmacol* 59 (2): 205-216.
- Poupard A, Emilie J. 1984. Autoimmunity in Parkinson's disease. In: Hassler RG, Christ JF, eds. *Parkinson specific motor and mental disorders. Role of the pallidum: pathophysiological biochemical and therapeutic aspects*. Adv Neurol 40. New York: Raven Press, 307-314.
- Radic M. 2014. Role of *Helicobacter pylori* infection in autoimmune systemic rheumatic diseases. *World J Gastroenterol*. 20(36): 12839-12846.
- Reale M, Iarlori C, Thomas A, Gambi D, Perfetti B, Nicola MDi, Onofri M. 2009. Peripheral cytokines profile in Parkinson's disease. *Brain Behaviour Immunity* 23: 55-63.
- Rees K, Stowe R, Patel S, Ives N, Breen K, Clarke CE, Ben-Shlomo Y. 2011. *Helicobacter pylori* eradication for Parkinson's disease. *Cochrane Database Syst Rev* doi10.1002/14651858.CD008453.pub2.
- Regnath T, Raecke O, Enninger A, Ignatius R. 2016. Increasing metronidazole and rifampicin resistance of *Helicobacter pylori* isolates obtained from children and adolescents between 2002 and 2015 in southwest Germany. *Helicobacter* doi: 10.1111/hel.12327
- Ren J, Bharti A, Raina D, Chen W, Ahmad R, Kufe D. 2006. MUC1 oncoprotein is targeted to mitochondria by heregulin-induced activation of c-Src and the molecular chaperone HSP90. *Oncogene* 25: 20-31
- Ricci C, Holton J, Vaira D. 2007: Diagnosis of *Helicobacter pylori*: invasive and non-invasive tests. *Best Pract Res Clin Gastroenterol*. 21: 299-3134
- Sanchez JE, Sáenz NG, Rincón MR, Martín IT, Sánchez EG, Martínez MJ. 2000. Susceptibility of *Helicobacter pylori* to mupirocin, oxazolidinones, quinupristin/dalfopristin and new quinolones. *Journal of Antimicrob Chemotherapy* 46: 283-285.
- Schiesling C, Kieper N, Seidel K, Krüger R. 2008. Review familial Parkinson's Disease-genetics, clinical phenotype and neuropathology in relation to the common sporadic form of the disease. *Neuropathol Appl Neurobiol*.34(3): 255-271
- Schott T, Kondadi PK, Hänninen ML, Rossi M. 2011. Comparative genomics of *Helicobacter pylori* and the human-derived *Helicobacter bizzozeronii* CIII-1 strain reveal the molecular basis of the zoonotic nature of non-pylori gastric *Helicobacter* infections in humans. *BMC Genomics*.12: 534.
- Schulz JD, Hawkes EL, Shaw CA. 2006. Cycad toxins, *Helicobacter pylori* and parkinsonism: cholesterol glucosides as the common denominator. *Med Hypotheses*. 66: 1222-1226.
- Sampson TR, Debelius JW, Thron T, Janssen S, Shastri GG, Ilhan ZE, Challis C, Schretter CE, Rocha S, Gradinaru V, Chesselet MF, Keshavarzian A, Shannon KM, Krajmalnik-Brown R, Wittung-Stafshede P, Knight R, Mazmanian SK. 2016. Gut microbiota regulate motor deficits and neuroinflammation in a model of Parkinson's disease. *Cell*. 167(6): 1469-1480.
- Semper RP, Mejías-Luque R, Groß C, Anderl F, Müller A, Vieth M, Busch DH, Prazeres da Costa C, Ruland J, Groß O, Gerhard M. 2014. *Helicobacter pylori*-induced IL-1 β secretion in innate immune cells is regulated by the NLRP3 inflammasome and requires the Cag pathogenicity island. *Journal of Immunol* 193:3566-3576.
- Sgouras D, Maragkoudakis P, Petraki K, Martinez-Gonzalez B, Eriotou E, Michopoulos S, Kalantzopoulos G, Tsakalidou E, Mentis A. 2004. *In vitro* and *in vivo* inhibition of *Helicobacter pylori* by *Lactobacillus casei* strain Shirota. *Appl and Environm Microbiol* 70(1): 518-526.
- Smet A, Van Nieuwerburgh F, Ledesma J, Flahou B, Deforce D, Ducatelle R, Haesebrouck F. 2013. Genome sequence of *Helicobacter heilmannii* sensu stricto ASB1 isolated from the gastric mucosa of a kitten with severe gastritis. *Genome Announc* 1:e00033-12.
- Smyk DS, Koutsoumpas AL, Mytilinaiou MG, Rigopoulou EI, Sakkas LI, Bogdanos DP. 2014. *Helicobacter pylori* and autoimmune disease: Cause or bystander. *World J Gastroenterol*. 20(3): 613-629.
- Solnick JV, Schauer DB. 2001. Emergence of diverse *Helicobacter* species in the pathogenesis of gastric and enterohepatic diseases. *Clin Microbiol Rev* 14(1): 59-97
- Solnick JV, Hansen LM, Canfield DR, Parsonnet J. 2001. Determination of the infectious dose of *Helicobacter pylori* during primary and secondary infection in rhesus monkeys. *Infection and Immunity* 69(11): 6887-6892.
- Stefanis L. 2012. α -Synuclein in Parkinson's Disease. *Cold Spring Harb Perspect Med*. 2(2): a009399.

- Stolp HB, Dziegielewska KM. 2009. Review: Role of developmental inflammation and blood-brain barrier dysfunction in neurodevelopmental and neurodegenerative disorders. *Neuropathology and applied Neurobiology* 35: 132-146.
- Stolte M, Kroher G, Meining A, Morgner A, Bayerdtirffer E, Bethke B. 1997. A comparison of *Helicobacter pylori* and *H. heilmannii* gastritis. A matched control study involving 404 patients. *Scand J Gastroenterol* 32: 28-33
- Strang RR. 1965. The association of gastrointestinal ulceration and Parkinson's disease. *Med J Austr* 52: 842-843.
- Sun YQ, Söderholm JD, Petersson F, Borch K. 2004. Long-standing gastric mucosal barrier dysfunction in *Helicobacter pylori*-induced gastritis in Mongolian gerbils. *Helicobacter* 9(3): 217-227.
- Tansey MG, McCoy MK, Frank-Cannon TC. 2007. Neuroinflammatory mechanisms in Parkinson's disease: Potential environmental triggers, pathways, and targets for early therapeutic intervention. *Exp Neurol* 208 (1): 1-25.
- Testerman TL and Morris J. 2014. Beyond the stomach: An updated view of *Helicobacter pylori* pathogenesis, diagnosis and treatment. *World J of Gastroenterol* 20(36): 12781-12808.
- Torok AM, Bouton AH, Goldberg JB. 2005. *Helicobacter pylori* induces interleukin-8 secretion by Toll-like receptor 2- and Toll-like receptor 5 dependent and independent pathways. *Infection and Immunity* 73 (3): 1523-1531.
- Thung I, Aramin H, Vavinskaya V, Gupta S, Park JY, Crowe SE, Valasek MA. 2016. Review article: the global emergence of *Helicobacter pylori* antibiotic resistance. *Aliment Pharmacol Ther* 43: 514-533.
- Valones MAA, Guimarães RL, Brandão LAC, de Souza PRE, Carvalho AAT, Crovela S. 2008. Principles and applications of polymerase chain reaction in medical diagnostic fields: a review. *Braz J Microbiol.* 40(1): 1-11.
- Vandenbroucke RE, Dejonckheere E, Van Lint P, Demeestere D, Van Wonterghem E, Vanlaere I, Puimège L, Van Hauwermeiren F, De Rycke R, Mc Guire C, Campeste C, López-Otin C, Matthys P, Leclercq G, Libert C. 2012. Matrix metalloprotease 8-dependent extracellular matrix cleavage at the blood-CSF barrier contributes to lethality during systemic inflammatory diseases. *Journal of Neuroscience* 32: 9805-9816.
- Van den Bulck K, Decostere A, Baele M, Driessen A, Debongnie JC, Burette A, Stolte M, Ducatelle R, Haesebrouck F. 2005. Identification of non-*Helicobacter pylori* spiral organisms in gastric samples from humans, dogs and cats. *J. Clin. Microbiol.* 42(5): 2256-2260.
- Van den Bulck K., Decostere A., Gruntar I., Baele M., Krt B., Ducatelle R., Haesebrouck F. 2005. *In vitro* antimicrobial susceptibility testing of *Helicobacter felis*, *Helicobacter bizzozeroni* and *Helicobacter salomonis*. *Antimicrob. Agents. Chemother.* 49: 2297-3000
- Vawter MP, Dillon-Carter O, Tourtellotte WW, Carvey P, Freed WJ. 1996. TGFbeta1 and TGFbeta2 concentrations are elevated in Parkinson's disease in ventricular cerebrospinal fluid. *Exp Neurol* 142 (2): 313-322.
- Vermeer PD, Einwalter LA, Moninger TO, Rokhlina T, Kern JA, Zabner J, Welsh MJ. 2003. Segregation of receptor and ligand regulates activation of epithelial growth factor receptor. *Nature.* 422(6929):322-326.
- Vermoote M, Vandekerckhove TTM, Flahou B, Pasmans F, Smet A, De Groote D, Van Crielinge W, Ducatelle R, Haesebrouck F. 2011. Genome sequence of *Helicobacter suis* supports its role in gastric pathology. *Vet Res* 42 (1): 51.
- Vermoote M, Pasmans F, Flahou B, Van Deun K, Ducatelle R, Haesebrouck F. 2011. Antimicrobial susceptibility pattern of *Helicobacter suis* strains. *Vet. Microbiol* 153: 339-342
- Vilaichone RK, Mahachai V, Tumwasorn S, Nunthapisud P, Kullavanijaya P. 2002. Inhibitory effect of *Lactobacillus acidophilus* on *Helicobacter pylori* in peptic ulcer patients: *in vitro* study. *J Med Assoc Thai.* 85 (1): S79-84.
- Villarán RF, Espinosa-Oliva AM, Sarmiento M, De Pablos RM, Argüelles S, Delgado-Cortés MJ, Sobrino V, Van Rooijen N, Venero JL, Herrera AJ, Cano J, Machado A. 2010. Ulcerative colitis exacerbates lipopolysaccharide-induced damage to the nigral dopaminergic system: potential risk factor in Parkinson's disease. *J Neurochem.* 114(6): 1687-700.
- Walters, W. A., Xu, Z. & Knight, R. 2014. Meta-analyses of human gut microbes associated with obesity and IBD. *FEBS Lett* 588, 4223-4233, doi:10.1016/j.febslet.2014.09.039.
- Wang Q, Liu Y, Zhou J. 2015. Neuroinflammation in Parkinson's disease and its potential as therapeutic target. *Transl Neurodegener.* 4: 19.
- Warren, J. R., and B. J. Marshall. 1983. Unidentified curved bacilli on gastric epithelium in active chronic gastritis. *Lancet* i: 1273-1275.
- White JR, Winter JA, Robinson K. 2015. Differential inflammatory response to *Helicobacter pylori* infection: etiology and clinical outcomes. *Journal of Inflammatory Research* 8: 137-147.

- Wirdefeldt K, Adami HO, Cole P, Trichopoulos D, Mandel J. 2011: Epidemiology and etiology of Parkinson's disease: a review of the evidence. *Eur J Epidemiol* 26: S1-S58.
- Wong F, Rayner-Hartley E, Byrne MF. 2014. Extraintestinal manifestations of *Helicobacter pylori*: a concise review. *World J of Gastroenterol* 20(34): 11950-11961.
- Wroblewski LE, Peek RM Jr. 2011. Targeted disruption of the epithelial-barrier by *Helicobacter pylori*. *Cell Commun Signal*. 9(1): 29.
- Wüppenhorst N, von Loewenich F, Hobmaier B, Vetter-Knoll M, Mohadjer S, Kist M. 2013. Culture of a gastric non-*Helicobacter pylori* *Helicobacter* from the stomach of a 14 year-old girl. *Helicobacter* 18: 1-5.
- Yagi BH, Zurenko GE. 1997. *In vitro* activity of linezolid and eperezolid, two novel oxazolidinone antimicrobial agents, against anaerobic bacteria. *Anaerobe* 3: 301-306.
- Yamanishi S, Iizumi T, Watanabe E, Shimizu M, Kamiya S, Nagata K, Kumagai Y, Fukunaga Y, Takahashi H. 2006. Implications for induction of autoimmunity via activation of B-1 cells by *Helicobacter pylori* urease. *Infect Immun* 74(1): 248-256.
- Zhang YJ, Li S, Gan RY, Zhou T, Xu DP, Li HB. 2015. Impacts of gut bacteria on human health and diseases. *Int J Mol Sci* 16, 7493-7519, doi:10.3390/ijms16047493

SUMMARY

Helicobacter pylori is the most prevalent *Helicobacter* species in humans. Infection with this agent has been associated with a broad range of gastric disorders, including gastritis, ulcers and cancer. Besides causing inflammation in the stomach, *H. pylori* infection has also been linked to extragastric manifestations, including Parkinson's Disease (PD). Patients with this neurodegenerative disorder frequently display gastrointestinal complaints but their symptoms cannot always be assigned to infection with *H. pylori*.

Gastric disease in humans has, however, also been associated with spiral-shaped non-*Helicobacter pylori* *Helicobacter* (NHPH) species, such as *H. suis*. This zoonotic agent, highly prevalent in pigs and non-human primates, is the most prevalent NHPH in humans and has been associated with gastritis, gastric and duodenal ulcers and mucosa-associated lymphoid tissue (MALT) lymphoma. The risk for developing MALT lymphoma is higher when infected with *H. suis* than *H. pylori*. Living in close contact with pigs or the consumption of raw or undercooked meat are considered to be risk factors for infection with this agent.

Compared to *H. pylori*, diagnosis of *H. suis* infections is extremely difficult due to the sparse and patchy distribution in the human stomach and the fastidious nature of this bacterium.

At the onset of this thesis, no information was available on the prevalence of *H. suis* in Parkinson's disease patients, nor was there any information found on how and to what extent an infection with this agent could contribute to the outcome of this neurodegenerative disorder. Additionally, the currently used methods to diagnose *H. suis* infections, lacked optimization in terms of sensitivity and specificity.

In our first study, we investigated the prevalence of *H. suis* infections in patients suffering from idiopathic parkinsonism (IP) and gastric complaints and compared it with the known frequency of *H. pylori* infection in this group. A control group of gastroenterology patients with no clinical parkinsonism was included for the purpose of comparison. Archived DNA extracts were examined for the presence of *H. suis* DNA using an *ureA*-based SYBR green RT-PCR assay. Results were confirmed by additional sequencing of the positive PCR amplicons. Additionally, multi-locus sequence typing (MLST) was performed to demonstrate that the *H. suis* DNA in the biopsy extracts was different from that of the *H. suis* lab strain used as positive control in the RT-PCR assay.

From the 60 patients with IP, 27% and 28% tested positive for *H. suis* and *H. pylori* respectively, whereas only 2% and 16% from the 256 control patients tested positive for *H. suis* and *H. pylori*, respectively. These results indicated that the relative risk of having an infection with *H. suis* was 10 times higher in PD patients compared to controls. Interestingly, from the 16 *H. suis* positive IP patients, 11 were from 19 with proven *H. pylori* eradication, 3 from 17 pre-*H. pylori* eradication and 2 from 24 *H. pylori* culture/PCR negative. Remarkably, the frequency of *H. suis* was the highest in the patients where *H. pylori* has been eradicated.

Partial MLST results for the *H. suis* DNA found in extracts from 6 IP patients and 1 control showed that the allele pattern in these samples differed from that of the DNA from the *in vitro* lab strain. This indicated that the results were not due to contamination.

In conclusion, this study highlighted that the prevalence of *H. suis* was higher in patients with IP compared to our control group and that it was even more exaggerated in those following *H. pylori* eradication therapy.

The brain is protected from alterations in systemic circulation by a series of specialized barriers, including the blood-brain barrier (BBB) and the blood cerebrospinal fluid barrier (blood-CSF barrier). Dysfunction of these barriers has been linked to multiple neurodegenerative disorders, including Parkinson's disease. Furthermore, systemic inflammation might contribute to the impairment of the BBB and the blood-CSF barrier. Since infections with *Helicobacter* have been suggested to induce systemic inflammation, it might be possible that a gastric *H. suis* infection could influence the integrity of the brain barriers and thereby contribute in the pathophysiology of neurodegenerative disorders. Therefore, **in the second study**, we evaluated the influence of an experimental infection with *H. suis* on the integrity of the stomach-brain axis in mice. To investigate this, C57BL/6 mice were inoculated intragastrically with viable *H. suis* and compared to a negative control group. At 4 weeks post-infection, animals were euthanized and samples from the stomach and brain were analyzed. An infection with *H. suis* induced gastric inflammation, characterized by the infiltration of mononuclear and polymorphonuclear cells in the mucosa and submucosa of the stomach. This finding was accompanied by the increased expression of interleukin (*IL*)-1 β , murine *IL*-8 homologs and increased protein levels of IL-1 β and IL-17. A significant increase in the gastric mucosal permeability was associated with aberrant expression of the tight junction protein zonula occludens (*ZO*)-1 and the transmembrane *Muc13* mucin. Furthermore, altered expression of the vasoactive intestinal peptide (*VIP*) and its receptor (*VIPR1*) was detected in the stomach of *H. suis*-infected mice. *VIP* has been shown to play a regulatory and protective role in barrier integrity, tight junction expression and mucin secretion in the gastrointestinal tract. These findings indicated that the loss of gastric-mucosal barrier integrity in *H. suis* infected mice might result in increased gastrointestinal leakage of cytokines, chemokines and bacterial compounds which could lead to systemic inflammation. Indeed, increased levels of IL-1 β and the murine IL-8 homologue, MIP1 α were observed in the serum of infected animals. Subsequently, the influence of *H. suis* on the integrity of the brain barriers (BBB and blood-CSF barrier) and brain homeostasis was investigated. A significant increase in the blood-cerebrospinal fluid (CSF) barrier permeability upon *H. suis* infection was seen. In contrast, blood-brain barrier permeability was unaffected at this acute stage of infection. Disturbance of the blood-CSF barrier was associated with an increased expression of the pro-inflammatory cytokine IL-1 β and the inducible nitric oxide synthase (*iNos*) and a decreased expression of the tight junction genes of claudin 1 (*clnd-1*) and *Occludin* (*Ocln*) in the choroid plexus epithelium that forms the blood-CSF barrier. Additionally, an increase of *IL*-6 expression in the hippocampus was observed upon infection with *H. suis*. These changes in cytokine expression in the brain were accompanied by the activation of microglia cells in *H. suis*-infected animals. Apart from the involvement of *VIP* in the maintenance of the gastrointestinal epithelial barrier, this neuropeptide also plays a role in brain homeostasis. Systemic administration of *VIP* indeed reversed the inflammation-induced disruption of the blood-CSF barrier integrity. In conclusion, we have identified the pivotal role of the choroid plexus in the communication between the gastrointestinal tract and the brain. Furthermore, our results indicate that disruption of the blood-CSF barrier is induced by *H. suis*-associated systemic inflammation and/or dysregulated *VIP* signaling.

Diagnosing gastric NHPH infections, in general, in humans is problematic. Non-invasive commercial tests are currently unavailable. Nowadays, diagnosis of infection with these fastidious pathogenic agents has been performed by the analysis of gastric biopsies using histological, molecular and microbiological techniques. To identify *H. suis* in the gastric biopsy samples, a *H. suis*-specific *ureA*-based SYBR green RT-PCR has been frequently used. For this molecular assay, additional sequencing of the positive amplicons is necessary, since cross-reactivity with host DNA and DNA from other gastric helicobacters occurred, questioning the specificity of this test. Attempts to directly and/or indirectly (after mouse-passage) isolate pure *H. suis* strains from human gastric biopsies remained unsuccessful. Direct isolation is hampered by the fastidious nature, the low colonization density and the patchy distribution of this pathogen, whereas indirect isolation is hampered by overgrowth with gastric contaminants, including *Lactobacillus* spp. In the **third study** of this thesis, we first developed a probe-based RT-PCR for the detection and quantitation of *H. suis* DNA in gastric biopsies. This RT-PCR was based on the amplification of the *H. suis*-specific *carR* gene. The probe detects a well-conserved 20bp long region within this gene. Validation of the assay revealed a high degree of specificity and sensitivity, high linear correlations (r^2 between 0.995-0.999) and high amplification efficiencies (>90%). In order to exclude cross-reaction with host DNA, gastric DNA extracts from human patients, pigs, mice and non-human primates were analyzed. In contrast to the *ureA*-based RT-PCR, no aspecific interactions with human, porcine, murine and non-human primate DNA, nor with DNA from other bacteria, including *Helicobacter* spp. and *Campylobacter* spp., were detected when using the *carR*-probe based RT-PCR.

Secondly, a mouse-passage-based protocol was further optimized for the isolation of low numbers of viable *H. suis* bacteria. Mice were inoculated with different numbers of viable *H. suis* bacteria (10^2 - 10^8). The animals were kept for 4 weeks to allow colonization and multiplication of *H. suis*. In order to inhibit the proliferation of gastric contaminants, including *Lactobacillus* spp., from the murine stomach, 5µg/ml linezolid was added to a biphasic medium consisting of *Brucella* agar with *Brucella* broth on top of it, both supplemented with vitox supplement, *Campylobacter*-selective supplement, amphotericin (5µg/ml), HCl (0.05%) and fetal bovine serum (20%). Using the optimized isolation protocol, we were able to re-isolate *H. suis* from the stomach of mice infected with at least 10^4 viable *H. suis* bacteria.

Finally, we also sequenced the genome of the porcine *H. suis* strain before and after mouse-passage in order to determine the influence of mouse-passage on the genome of *H. suis*. Our data shows that mouse-passage of *H. suis* does not result in major recombination events within the genome of this bacterium, suggesting that *H. suis* strains obtained after mouse-passage can be used for further characterization of this pathogen.

In conclusion, the methods described above can be implemented for detection or isolation of *H. suis* from human gastric biopsies.

Samenvatting

Helicobacter pylori is de meest voorkomende *Helicobacter* soort bij de mens. Deze kiem wordt geassocieerd met gastritis, peptische ulcera en maagkanker. Daarnaast worden infecties met *H. pylori* ook gelinkt aan andere aandoeningen, zoals de ziekte van Parkinson. Patiënten met de ziekte van Parkinson vertonen vaak maagklachten, maar hun symptomen kunnen niet altijd toegeschreven worden aan een infectie met deze pathogeen. Maagaandoeningen bij de mens kunnen ook veroorzaakt worden door infecties met niet-*H. pylori* *Helicobacter* species, zoals *H. suis* die van nature uit de maag van varkens en primaten koloniseert. Bij de mens veroorzaakt dit zoönotisch agens gastritis, peptische ulcera en lymfomen van het lymfoïd weefsel van de maagmucosa (MALT-lymfomen). Bovendien is het risico op de ontwikkeling van MALT-lymfomen groter bij een *H. suis* infectie dan bij een infectie met *H. pylori*. Mensen die in nauw contact leven met varkens of rauw/niet doorbakken varkensvlees consumeren, hebben meer kans om besmet te worden met *H. suis*.

De diagnose van infecties met *H. suis* bij de mens is moeilijker in vergelijking met *H. pylori* omwille van het zeer focale kolonisatiepatroon in de maag en de hoge moeilijkheidsgraad om dit agens *in vitro* te cultiveren. Bij de start van dit doctoraatsonderzoek was er nog geen informatie beschikbaar over het voorkomen van *H. suis* infecties bij patiënten met de ziekte van Parkinson. Het was eveneens nog onduidelijk of en in welke mate een infectie met deze pathogeen kan bijdragen tot het ziektebeeld van deze neurodegeneratieve aandoening. Bovendien bleek dat de aangewende technieken voor de detectie van *H. suis* infecties bij de mens niet optimaal waren.

In een **eerste studie** werd het voorkomen van *H. suis* infecties bestudeerd bij patiënten met de ziekte van Parkinson die maagklachten hadden en vervolgens vergeleken met de reeds gekende prevalentie van *H. pylori* binnen deze groep. Een controle groep (patiënten met maagklachten, maar zonder de ziekte van Parkinson) werd eveneens ingesloten ter vergelijking. Het DNA van de maagstalen werd onderzocht op de aanwezigheid van *H. suis* DNA aan de hand van een op het *ureA*-gen gebaseerde RT-PCR analyse en sequencerig van de positieve PCR amplicons. Multi-locus sequence typing (MLST) werd uitgevoerd om na te gaan of het *H. suis* DNA in de maagbiopten verschillend was van dat van de *H. suis* laboratorium stam die gebruikt werd als positieve controle.

In de groep van patiënten met de ziekte van Parkinson testte 27% positief voor *H. suis* en 28% voor *H. pylori*. In de controle groep werd *H. suis* slechts aangetoond in 2% van de patiënten, terwijl 16% van de patiënten geïnfecteerd waren met *H. pylori*. Deze resultaten tonen aan dat een *H. suis* infectie 10 maal frequenter voorkwam bij patiënten met de ziekte van Parkinson in vergelijking met de controle groep. Bovendien bleek dat de meeste patiënten met de ziekte van Parkinson die positief testten voor *H. suis*, een succesvolle *H. pylori* eradicatie therapie achter de rug hadden. Tot slot bleek uit de MLST resultaten dat het *H. suis* DNA in de maagbiopten verschillend was van dat van de laboratorium stam waardoor contaminatie kon uitgesloten worden.

Samengevat beschreven we in deze studie een opmerkelijk hoge prevalentie van *H. suis* infecties bij patiënten met de ziekte van Parkinson en voornamelijk bij de patiënten die voorafgaand een anti-*H. pylori* therapie hadden ondergaan.

De hersenen worden beschermd tegen gevaarlijke stoffen in de bloedsomloop door een aantal gespecialiseerde barrières, zoals de bloed-hersen barrière (BBB) en de bloed-cerebrospinaal vocht barrière (bloed-CSF barrière). Bij verstoring van deze barrières kunnen cytokines en chemokines vanuit de bloedbaan de hersenen binnendringen en inflammatie induceren. Er werd inderdaad reeds aangetoond dat systemische inflammatie een belangrijke rol speelt in de ontwikkeling van de ziekte van Parkinson. Bovendien doodt *H. suis* de epitheliale cellen in de maag en wordt de epitheelbarrière ter hoogte van de maag doorbroken, waardoor ontstekingsmediatoren ook in de circulatie kunnen terechtkomen. Daarom werd er tijdens **de tweede studie** van deze thesis nagegaan in welke mate en op welke manier een *H. suis* infectie de BBB en de bloed-CSF barrière beïnvloedt. Hiervoor werden C57BL/6 wildtype muizen intragastraal geïnoculeerd met levende *H. suis* bacteriën of met het opgroeimedium van deze kiem (negatieve controle). Vier weken na infectie werden de dieren geëuthanaseerd en stalen van de maag en de hersenen werden verzameld en geanalyseerd. Inflammatie ter hoogte van de maag werd gekenmerkt door de infiltratie van mononucleaire en polymorfonucleaire cellen in de mucosa en de submucosa van de maag en een verhoogde mRNA en/of eiwit expressie van interleukine (IL)-1 β , IL-17 en de chemokines KC en LIX, die bij muizen tot expressie worden gebracht en homoloog zijn aan IL-8 van mensen. Een verhoogde maagpermeabiliteit werd geassocieerd met veranderingen in mRNA en eiwit expressie van het transmembranair Muc13 mucine en het tight junction eiwit zonulae occludens (ZO)-1. Bij de met *H. suis* geïnfecteerde dieren werd tevens een onderdrukking van de expressie van het vasoactief intestinaal peptide (VIP) vastgesteld. Dit peptide speelt een belangrijke rol in het behoud van de maagmucosa integriteit. Een verstoorde mucosale maagbarrière kan leiden tot lekkage van cytokines, chemokines en bacteriële componenten vanuit de maag naar de bloedsomloop en kan dus resulteren in systemische inflammatie. In deze studie werden in het serum van de met *H. suis* geïnfecteerde muizen verhoogde titers gedetecteerd van IL-1 β en MIP1 α , een cytokine dat net als KC en LIX door muizen tot expressie wordt gebracht en homoloog is aan IL-8 van mensen. Vervolgens werd de impact van een *H. suis* infectie op de integriteit van de hersenbarrières onderzocht. Er werd een significante stijging in de bloed-CSF barrière permeabiliteit waargenomen. De permeabiliteit van de BBB, daarentegen, werd niet aangetast in dit acuut stadium van de infectie. De verstoring van de bloed-CSF barrière werd geassocieerd met een verhoogde expressie van *IL-1 β* en het induceerbare stikstof oxide synthase (*iNos*), alsook met een gedaalde expressie van de tight junctions claudine 1 (*clnd-1*) en occludine (*Ocln*) ter hoogte van het choroid plexus epitheel dat de bloed-CFS barrière vormt. Tevens werd in de hippocampus van de met *H. suis* geïnfecteerde dieren een verhoogde expressie waargenomen van het gen dat codeert voor IL-6. De veranderingen in de expressie van cytokines in de hersenen gingen eveneens gepaard met een activatie van de microgliacellen bij *H. suis* geïnfecteerde muizen. VIP zou niet enkel een rol spelen in het behoud van de integriteit van de maagmucosa, maar eveneens in het behoud van de homeostase in de hersenen. Vermits de expressie van VIP verstoord was in de maag van met *H. suis* geïnfecteerde dieren, wilden we in deze studie het effect van VIP op de hersenen nagaan. Hiervoor werden muizen intraperitoneaal geïnjecteerd met lipopolysaccharide, wat een ontstekingsreactie en een verstoring van bloed-CSF barrière veroorzaakt. Systemische toediening van VIP resulteerde in een herstel van de verstoorde bloed-CSF barrière.

Samengevat, werd in deze studie aangetoond dat de choroid plexus een centrale rol speelt in de communicatie tussen de maag en de hersenen. Bovendien wijzen onze resultaten erop dat een verstoorde bloed-CSF barrière het gevolg kan zijn van een door *H. suis*-geïnduceerde systemische inflammatie en/of een verstoring in de VIP huishouding.

De diagnose van infecties met niet-*H. pylori Helicobacter* species is problematisch en tot op heden zijn er nog geen commerciële testen beschikbaar. Voor het aantonen van *H. suis* DNA in stalen van de maag wordt er o.a. gebruik gemaakt van een SYBR green RT-PCR techniek gebaseerd op het *ureA* gen van dit agens. Bijkomende sequencerende van de positieve PCR amplicons is echter noodzakelijk omwille van het voorkomen van kruisreactiviteit met humaan DNA en DNA van andere gastrale *Helicobacter* species. Daarnaast is men er nog niet in geslaagd om *H. suis* te isoleren uit maagbiopten (rechtstreeks of via passage in een muismodel) en te cultiveren *in vitro*. In de **derde studie** werd dan ook eerst een nieuwe RT-PCR ontwikkeld voor de detectie en kwantificatie van *H. suis* DNA in maagbiopten. De RT-PCR was gebaseerd op de amplificatie van het *H. suis*-specifieke *carR* gen. Om amplicons te detecteren werd er, in plaats van het frequent gebruikte SYBR green, geopteerd voor een fluorescent gemerkte probe die een geconserveerde regio van 20 baseparen binnen het *carR* gen detecteert. Validatie van deze techniek toonde een hoge specificiteit en sensitiviteit aan, alsook hoge lineaire correlaties (r^2 tussen 0.995-0.999) en een hoge amplificatie efficiëntie (>90%). Er werd geen kruisreactie vastgesteld met DNA van andere bacteriën, humane patiënten, varkens, muizen en niet-humane primaten.

Vervolgens werd een bestaand protocol voor de isolatie van lage aantallen levende *H. suis* bacteriën uit de maag via passage in een muismodel verder geoptimaliseerd. Hiervoor werden muizen intragastraal geïnoculeerd met verschillende concentraties levende *H. suis* kiemen (10^8 - 10^2). De dieren werden gedurende 4 weken aangehouden om kolonisatie en vermeerdering van *H. suis* mogelijk te maken. Daarna werden ze geëuthanaseerd en de maag werd voorbehandeld met een 1% HCl oplossing gedurende 45 minuten. Deze voorbehandelde stalen werden dan geïnoculeerd in het bifasische opgroei-medium van *H. suis* waaraan 5µg/ml linezolid was toegevoegd. Dit antibioticum is nodig om te vermijden dat bacteriën die behoren tot de microbiota van de muizenmaag, waaronder *Lactobacillus* spp., het medium overwoekeren en isolatie van *H. suis* onmogelijk maken. Het opgroei-medium bestond uit *Brucella* agar met daarboven *Brucella* broth, beiden gesupplementeerd met Vitox, *Campylobacter*-selectief supplement, amphotericine (5µg/ml), HCl (0,05%) en geïnactiveerd foetaal kalfserum (20%). Door gebruik te maken van het geoptimaliseerde isolatie protocol slaagden we erin om *H. suis* te isoleren uit de maag van muizen die initieel werden geïnfecteerd met tenminste 10^4 levende *H. suis* kiemen.

Tot slot werd er ook nagegaan of de passage van *H. suis* in de muizenmaag een impact heeft op het genoom van deze bacterie. Hiervoor werd het genoom van de *H. suis* stam voor en na isolatie uit de muizenmaag gesequeneerd. Uit de genoomanalyses konden geen grote veranderingen in het genoom waargenomen worden. Samengevat kunnen we stellen dat de ontwikkelde en geoptimaliseerde diagnostische technieken gebruikt kunnen worden voor de detectie en/of isolatie van *H. suis* in humane maagbiopten.

CURRICULUM VITAE

Caroline Blaecher werd geboren op 18 januari 1988 te Brussel. Nadat ze haar secundair onderwijs aan het Jan van Ruusbroeckcollege succesvol had afgerond, startte ze in 2006 de studies diergeneeskunde aan de Universiteit Gent. In 2012 behaalde ze met onderscheiding het diploma van Master in de Diergeneeskunde.

Omwille van haar grote interesse in het wetenschappelijk onderzoek, startte ze onmiddellijk daarna een doctoraatsonderzoek bij de Vakgroep Pathologie, Bacteriologie en Pluimveeziekte aan de Faculteit Diergeneeskunde van de Universiteit Gent. Gedurende 4 jaar verrichtte ze, in het kader van een Geconcentreerde Onderzoeksactie (GOA) van de Universiteit Gent (Code GOA 01G00408) en de "Psychiatry Research Trust" (Registered Charity no. 284286) van Londen, onderzoek naar de mogelijke associatie tussen *Helicobacter suis* infecties en de ziekte van Parkinson en optimaliseerde ze diagnostische testen voor de detectie van deze infecties bij de mens. Dit doctoraatsonderzoek resulteerde in het huidige proefschrift.

Caroline Blaecher is auteur en medeauteur van verschillende publicaties in internationale wetenschappelijke tijdschriften. Ze nam eveneens actief deel aan nationale en internationale congressen en was daar meermaals spreker.

BIBLIOGRAPHY

Publications in international journals

- 1) **Blaecher C**, Bauwens E, Tay A, Peters F, Dobbs S, Dobbs J, Charlett A, Ducatelle R, Haesebrouck F, Smet A. A novel isolation protocol and probe-based RT-PCR for diagnosis of gastric infections with the zoonotic pathogen *Helicobacter suis*. *Helicobacter* 2016; doi: 10.1111/hel.12369.
- 2) Liu C, Rossi M, Linden S, Padra M, **Blaecher C**, Bauwens E, Joosten M, Flahou B, Van den Broucke W, Ducatelle R, Haesebrouck F, Smet A. The *Helicobacter heilmannii* HofE and HofF genes are essential for colonization of the gastric mucosa and play a role in IL-1 β -induced gastric MUC13 expression. *Helicobacter*.2016; 6: 504-522.
- 3) Liu C, Smet A, **Blaecher C**, Flahou B, Ducatelle R, Linden S, Haesebrouck F. Gastric *de novo* expression and spasmolytic polypeptide-expressing metaplasia during *Helicobacter heilmannii* infection. *Infect and Immun* 2014; 82(8): 3227-3239.
- 4) **Blaecher C***, Smet A*, Flahou B, Pasmans F, Ducatelle R, Taylor D, Weller C, Bjarnason I, Charlett A, Lawson AJ, Dobbs RJ, Dobbs SM, Haesebrouck F. Significantly higher frequency of *Helicobacter suis* in patients with idiopathic parkinsonism than in control patients. *Aliment Pharmacol Ther.* 2013; 38(11-12): 1347–1353. (* equal contribution).
- 5) Joosten M, **Blaecher C**, Flahou B, Ducatelle R, Haesebrouck F, Smet A. Diversity in bacterium-host interactions within the species *Helicobacter heilmannii sensu stricto*. *Vet Res.* 2013; 29: 44:65.

Congress abstracts

- 1) **Blaecher C**, Gorré N, Bauwens E, Flahou B, Ducatelle R, Libert C, Vandenbroucke RE, Haesebrouck F, Smet A. (2016). Influence of an experimental *Helicobacter suis* infection on the stomach-brain axis in mice. 12th International workshop on pathogenesis and host response in *Helicobacter* infections konventum-Helsingør, Denmark.
- 2) **Blaecher C**, Bauwens E, Dobbs S, Dobbs J, Ducatelle R, Haesebrouck F, Smet A. (2016). A novel isolation protocol and probe-based RT-PCR for diagnosis of gastric infections with the zoonotic pathogen *Helicobacter suis*. 12th International workshop on pathogenesis and host response in *Helicobacter* infections konventum-Helsingør, Denmark.
- 3) Bauwens E, **Blaecher C**, Gorré N, Ducatelle R, Libert C, Vandenbroucke RE, Haesebrouck F, Smet A. (2016). Impact of a *Helicobacter heilmannii* infection on the gut-brain axis and gut-microbiome. Program IRC Symposium 'Innate immunity and lymphoid homeostasis'-Ghent, Belgium.

- 4) Bauwens E, **Blaecher C**, Gorlé N, Ducatelle R, Libert C, Vandenbroucke RE, Haesebrouck F, Smet A. (2016). Impact of a *Helicobacter heilmannii* infection on the gut-brain axis and gut-microbiome. 12th International workshop on pathogenesis and host response in *Helicobacter* infections konventum-Helsingør, Denmark.
- 5) **Blaecher C**, Gorlé N, Bauwens E, Flahou B, Ducatelle R, Libert C, Vandenbroucke R, Haesebrouck F, Smet A. (2015). Influence of the stomach-brain axis in relation to an experimental *Helicobacter suis* infection in mice. XXVIIIth International workshop on *Helicobacter* and microbiota in inflammation and cancer-Nicosia, Cyprus.
- 6) Liu C, Rossi M, **Blaecher C**, Bauwens E, Flahou B, Ducatelle R, Haesebrouck F, Smet A. (2015). Involvement of the *Helicobacter heilmannii* locus encoding the HOF outer membrane proteins in binding to the gastric mucosa. XXVIIIth International workshop on *Helicobacter* and microbiota in inflammation and cancer-Nicosia, Cyprus.
- 7) Liu C, Smet A, **Blaecher C**, Flahou B, Ducatelle R, Linden S, Haesebrouck F. (2014). Gastric *de novo* Muc13 expression and spasmolytic polypeptide-expressing metaplasia during *Helicobacter heilmannii* infection. 11th International workshop on pathogenesis and host response in *Helicobacter* infections- Helsingør, Denmark.
- 8) Liu C, Smet A, **Blaecher C**, Flahou B, Ducatelle R, Linden S, Haesebrouck F. (2014). Gastric *de novo* Muc13 expression and spasmolytic polypeptide-expressing metaplasia during *Helicobacter heilmannii* infection. International symposium on mechanisms of innate immunity, cell death and inflammation (ICDI symposium)- Ghent, Belgium.
- 9) **Blaecher C**, Smet A, Flahou B, Pasmans F, Ducatelle R, Bjarnason I, Lawson AJ, Charlett A, Taylor D, Weller C, Dobbs RJ, Dobbs SM, Haesebrouck F. (2013). Parkinson's disease and *Helicobacter suis*: pigs, pork or transmission of human-adapted *H. suis*-like bacteria. Studiedag International Pig Veterinary Society (IPVS) Belgian Branch-Tervuren, Belgium.
- 10) Dobbs SM, Dobbs JR, Smet A, Weller C, Charlett A, Bjarnason I, Lawson AJ, Ibrahim MAA, Taylor D, **Blaecher C**, Pasmans F, Ducatelle R, Haesebrouck F. (2013). Clinical significance of *Helicobacter suis* in idiopathic parkinsonism. 26th International workshop on *Helicobacter* and related bacteria in chronic digestive inflammation and gastric cancer-Madrid, Spain.
- 11) Haesebrouck F, Smet A, Flahou B, **Blaecher C**, Pasmans F, Ducatelle R, Bjarnason I, Lawson AJ, Charlett A, Taylor D, Weller C, Dobbs RJ, Dobbs SM. (2013). Parkinson's Disease and *Helicobacter suis*: pigs, pork or transmission of human-adapted *H. suis*-like bacteria. 26th International workshop on *Helicobacter* and related bacteria in chronic digestive inflammation and gastric cancer-Madrid, Spain.

Dankwoord

“Teamwork is the secret that makes common people achieve uncommon results” is een quote van Onuhoa die voor mij weergeeft hoe een doctoraat tot stand komt. Meer dan 4 jaar geleden werd ik met open armen ontvangen op de vakgroep Pathologie, Bacteriologie en pluimveeziekten van de Faculteit Diergeneeskunde aan de Universiteit Gent. Doorheen de jaren kon ik steeds rekenen op steun, raad, hulp en begeleiding van de vele collega’s en in het bijzonder mijn promotoren.

Mijn eerste woord van dank is dan ook gericht aan mijn promotoren. **Prof. dr. Freddy Haesebrouck**, bedankt voor uw uitstekende begeleiding tijdens mijn doctoraat. Uw onuitputtelijke en oprechte interesse, toewijding, kritische aanpak en raadgevingen hebben voor een groot deel bijgedragen tot de finale versie van dit werk. Ik zou u ook willen bedanken voor de kans die u me gegeven heeft om dit doctoraat te starten, voor het vertrouwen en tot slot voor alle kennis die ik dankzij uw toedoen heb vergaard. **Prof. dr. Annemieke Smet**, heel erg bedankt voor de goede begeleiding, je brede waaier aan ideeën en de vele verbeteringen die dit doctoraat vorm hebben gegeven. Bedankt dat ik steeds bij jou terecht kon op momenten dat ik ‘vast’ zat of vragen had. Of ik nu positief of negatief nieuws bracht, ik kon steeds rekenen op jouw steun, vertrouwen en hulp. Door jouw doorzettingsvermogen, vastberadenheid en motivatie stimuleer je mensen om het beste van zichzelf te geven en te streven naar “beter”. Nogmaals bedankt voor alles en ik wens je nog tonnen veel succes in je toekomstige carrière!!!

I would also like to thank my promoters **Sylvia and John Dobbs** for all the help, support and confidence throughout these last 4 years. Thanks for the opportunity you have offered me to meet your patients, I really appreciated that a lot. Finally, I want to thank you both for the nice cooperation and your contribution to this thesis.

Vervolgens zou ik de leden van de begeleidings- en examencommissie **Prof. dr. R. Vandenbroucke**, **Prof. dr. R. Ducatelle**, **Prof. dr. B. De Winter** en **Prof. dr. S. Backert**, willen bedanken voor hun tijd en moeite om deze doctoraatsthesis grondig na te lezen. Jullie verbeteringen, opbouwende kritiek en suggesties zorgden ervoor dat dit proefschrift “af” is. Graag zou ik de promotor van de GOA, **Prof. dr. R. Ducatelle**, in het bijzonder willen bedanken. **Prof. dr. S. Backert**, thank you for travelling to Ghent to attend my PhD defense. Tot slot zou ik ook nog een woord van dank willen richten tot **Prof. dr. H. Favoreel**. Dank u wel om als voorzitter van de examencommissie alles in goede banen te leiden.

Zonder financiële steun was dit onderzoek nooit mogelijk geweest. Daarom zou ik graag de **Universiteit Gent** willen bedanken voor het toekennen van de GOA die financiële steun bood voor het uitvoeren van mijn doctoraatsonderzoek.

Furthermore, I would also like to thank the “**Psychiatry Research Trust**” (Registered Charity no. 284286) from London for their financial donations.

Bijkomend zou ik eveneens alle co-auteurs van buiten onze vakgroep willen bedanken voor de goede en aangename samenwerking en de bijdrage die ze aan mijn publicaties hebben geleverd: **Prof. D. Taylor, dr. C. Weller, dr. I. Bjarnason, dr. A. Charlett, dr. A. J. Lawson, dr. A. Tay, dr. F. Peters, dr. S. Balusu, Prof. dr. C. Libert en J. Vandewalle.**

Graag zou ik ook **Prof. dr. R. Vandenbroucke** en haar **onderzoeksteam**, en in het bijzonder **Nina Gorlé, Griet van Imschoot** en **Elien van Wontelghem**, willen bedanken voor hun bijdrage aan de tweede studie in dit proefschrift. Door de kennis en expertise van het *Helicobacter*-team en het 'brain-barrier'-team van Prof. dr. Vandenbroucke te combineren werden mooie resultaten en nieuwe inzichten bekomen. Dank jullie allemaal heel erg voor de leuke samenwerking, de organisatie, de uitleg, de hulp en 'last but not least' alle kennis.

Vervolgens zou ik graag mijn collega's willen bedanken voor alle hulp, de leuke werkomgeving en de onvergetelijke en mooie herinneringen die jullie me de afgelopen 4 jaar bezorgd hebben.

Bij de aanvang van deze doctoraatstudie werd ik opgenomen binnen het '*Helicobacter*'-team. Zij maakten me wegwijs in het labo, stelden me voor aan de rest van de collega's en stonden steeds klaar om te me te helpen. Hierdoor voelde ik me meteen thuis binnen deze dynamische onderzoeksgroep. **Eva**, ik weet niet goed waar ik voor jou moet beginnen. Ik denk niet dat het mogelijk is om in woorden weer te geven hoe dankbaar ik ben en hoeveel je de afgelopen twee jaar voor mij hebt betekend. Dank je voor al jouw hulp met mijn labowerk, voor alle leuke, hilarische momenten waar we samen om gelachen hebben, de steun die je altijd hebt gegeven, de leuke gesprekken, het vertrouwen, de eindeloze vriendschap en nog zoveel meer.... Bedankt voor alles, ik zal dit heel mijn leven meedragen! Verder wil ik je nog enorm veel succes wensen met het vervolg van je onderzoek. Ik duim heel erg hard voor jou. Met jouw motivatie, voeling met het onderzoek, je harde werkersmentaliteit en gedrevenheid ben ik er zeker van dat het in orde gaat komen! Je verdient het heel erg!!!! **Chloë**, ook jij bent twee jaar geleden begonnen aan onze vakgroep. Al van in het begin klikte het en dat is zo gebleven. Dank je voor de hulp, de leuke herinneringen, de vele leuke activiteiten die we samen gedaan hebben en de vele knuffels die op lastigere momenten veel steun boden. Met jouw gedrevenheid, toewijding en kennis zal je het ongetwijfeld ver schoppen. **Iris** en **Myrthe**, heel erg bedankt voor alle leuke babbels, de vele humoristische momenten, de koffiepauzetjes, de steun, de raad en de hulp waar ik bij jullie steeds voor terecht kon. Myrthe, jij mocht als eerste van ons het goede voorbeeld geven om je doctoraat af te leggen, Iris en ik zullen een voorbeeld aan je nemen. Ik wens jullie allebei enorm veel succes met jullie toekomstige carrière en nog zoveel meer. **Helena**, hoewel je nog maar net begonnen bent met je PhD, kan iedereen zien dat je van aanpakken weet, leergierig en enorm gemotiveerd bent. Ik wens je nog enorm veel succes met je doctoraat en al je toekomstplannen. Ik zal jullie allemaal heel erg missen. **Ellen** en **Hannah**, dank jullie voor de leuke momenten op het werk en daarbuiten. **Guangzhi** and **Cheng**, thank you both for the help with my experiments in the lab. I wish you both all the best! **Bram**, dank je wel voor je hulp, raad, verbeteringen en expertise tijdens mijn doctoraatonderzoek. Ik hoop dat alles naar behoren verloopt op je huidige job en wens je nog veel succes toe in je toekomstige carrière. **Lien** en **Miet**, ook jullie wens ik nog veel succes toe in de toekomst!

Graag zou ik kort ook een woord van dank willen richten tot **Prof dr. Pasmans, Prof. dr Van Immerseel** en **dr. Boyen** om mee in te staan voor de organisatie en de financiering van het labo. **Prof dr. Pasmans**, dank u wel voor uw bijdrage als promotor van het FOD project met betrekking tot *Helicobacter*.

Sofie DB en **Nathalie**, heel erg bedankt voor jullie expertise, geduld, ervaring, en al de hulp bij de experimenten. Ik was altijd heel erg blij en dankbaar als jullie een handje konden toesteken in mijn onderzoek. De momenten die we samen gependend hebben in de flows van het Salmonellalabo en bij de muisexperimenten in de stallen zullen me steeds bijblijven. **Serge** en **Arlette**, dank jullie wel voor al de hulp tijdens de routine en voor de leuke gesprekken die we gevoerd hebben terwijl ik de muizenbakjes aan het kuisen was. Serge, nog heel veel succes met de verbouwingen! Arlette, ik hoop dat je revalidatie vlot kan verlopen en dat je snel weer de natuur kan gaan verkennen! **Koen**, jij bent altijd het ‘zonnetje’ op het labo: als je vrolijk fluitend in de gang passeerde verscheen er spontaan een glimlach op mijn gezicht. Dank je voor al de administratieve zaken die je steeds met een lach voor mij hebt geregeld. **Jo**, dank je voor al de hulp bij de bestellingen en de snelle reactie als ik weer eens snel iets heel dringend moest hebben. **Gunther M.**, jij was er elke dag om me als eerste een ‘goeiemorgen’ toe te wensen. Je stond telkens weer klaar om alles te regelen als er weer eens muisjes aangekomen waren. **Connie**, hoeveel keer hebben we niet samen in de PCR labo’s gezeten... dank je voor alle leuke gesprekken die we daar gevoerd hebben. **Chana, Julie** en **Sarah**, het was steeds een plezier om met jullie samen in het labo te werken, jullie positieve ingesteldheid zal me steeds bijblijven. **Jackeline**, hoewel ik niet echt de tijd heb gehad om je goed te leren kennen, kan ik me jou enkel al glimlachend voorstellen en was je altijd te vinden voor een voor een fijne conversatie. **Christian, Delphine** en **Joachim**: Jullie wisten al hoe laat het was als ik bij jullie binnenkwam hé. “Ze is daar weer met haar muizenmaagjes” zullen jullie waarschijnlijk vaak gedacht hebben. Erg bedankt om steeds met een brede glimlach mijn staaltjes zo snel mogelijk in te bedden, te snijden en te kleuren. Het was een plezier om met jullie samen te werken! **Sarah L** en **Marjan**, dank jullie voor de snelle uitvoering van de immunohistochemische kleuringen en de fijne gesprekken. **Marleen**, zoals jij lopen er geen twee (ook niet je tweelingzus) op de aarde. Ik ben enorm blij dat ik je heb mogen leren kennen. Dank je wel voor de oneindige vriendschap, bezorgdheid, medeleven, hulp, advies en steun die je me de afgelopen 4 jaar hebt gegeven. Je ben echt goud waard en ik wens je nog veel geluk toe.

Karen, voor ons begon het allemaal met toertjes lopen rond de faculteit. Tijdens het lopen, hadden we natuurlijk veel tijd om te praten en daaruit ontstond een hechte vriendschap. Bedankt voor deze 4 mooie jaren, alle leuke momenten en de onvergetelijke herinneringen. Ik ga onze loopsessies heel erg missen. Nog veel succes met je doctoraat, ik duim heel erg voor jou. **Lieze**, de spring-in’t-veld van de vakgroep! Je hebt de gave om iedereen van het ene op het andere moment direct gelukkig te maken en te laten lachen. Je gedrevenheid, doorzettingsvermogen en enthousiasme zijn een voorbeeld voor iedereen! Bedankt voor alle leuke (soms knotsgekke) belevenissen. Nog veel succes verder met je PhD, jullie huisje, de diertjes, en nog zo veel meer. **Marc**, als ik jou tegenkwam, was mijn dag stante pede goed! Je wist me steeds te entertainen, op te beuren, te motiveren, aan het lachen te brengen, etc. Elke keer als je langs het raam van onze bureau passeerde werd ik begroet met een ‘zwaaitje en een brede glimlach’. Deze kleine, simpele dingen zullen me altijd bij blijven. Heel erg bedankt voor alles en ik wens je nog enorm veel geluk en voorspoed toe. **Roel**, ik denk niet dat er veel

mensen zo gedreven, gemotiveerd en gebeten zijn door het onderzoek als jij. Je was een voorbeeld voor ons allemaal! Je stond bovendien altijd klaar om mensen te helpen, zelf al moest je daarom je eigen werk uitstellen. Heel erg bedankt, nog veel succes in jullie huisje en ik wens je het allerbeste voor de toekomst. **Gunther**, al wat ik hierboven over Roel heb beschreven, is eveneens voor jou van toepassing. Door je brede kennis ben je één van de aanspreekpunten in het labo waar iedereen bij terecht kan. Jij gaat het ongetwijfeld nog ver brengen. Ik had ook het geluk dat ik je bureaugenoot kon zijn. Dank je wel voor alle leuke momenten! **An G**, of correcter Prof dr. Garmyn, dank je wel voor alle hulp, de steun en vooral de leuke tijd op onze bureau en daarbuiten! **Justine** en **Wolf**: na de herorganisatie van de plaatsen op de vakgroep zijn jullie onze bureaugenootjes geworden. Dank jullie wel voor de vele leuke momenten dat we samen hebben beleefd! Ik wens jullie allebei nog enorm veel succes in jullie verdere carrière. **Evelien** en **Jasmien** jullie zijn in oktober bij Eva en mij op de bureau komen zitten. Dank jullie wel voor de steun bij het schrijven van mijn thesis. Ik wens jullie nog veel geluk met jullie onderzoek. **Lien**, **Nele** en **Maxime**, dank jullie voor de vele onvergetelijke momenten dat we samen hebben beleefd op het labo maar ook buiten de werkuren! **Vanessa**, dank je wel voor alle goede raad en voor je hulp als ik een probleem had met PCR of RT-qPCR. Met jouw ervaring en kennis ben je quasi onmisbaar geworden voor het labo. **Evy**, we waren ongeveer gelijktijdig aan ons doctoraat aan het schrijven en ik heb daardoor veel steun en tips gekregen van jou. Dikke merci hiervoor! **Mark**, je hebt me enorm geholpen bij het ontwikkelen van de probe. Mede dankzij jou heb ik dat tot een goed einde kunnen brengen, waarvoor mijn oprechte dank!

Vanaf mijn eerste werkdag voelde ik me onmiddellijk thuis op onze vakgroep. Dit was grotendeels te wijten aan de vriendelijke, behulpzame en goedlachse mensen die er rondliepen en nog steeds rondlopen. Hoewel het steeds een komen en gaan is van personeel, vormen de collega's toch een hechte groep waarvan ik deel mocht uitmaken. **David**, **Jonas**, **Ilias**, **Elin**, **Alexander**, **Alex**, **Annatachja**, **Anneleen**, **Astra**, **Annelies**, **Ruth**, **Shaoiji**, **Silvana**, **Sofie G**, **Celine**, **Karolien**, **Dorien**, **Evelien**, **Fien**, **Filip**, **Deon**, **Julie**, **Keeley**, **Laura**, **Leen**, **Ilse**, **Tom**, **Lonneke**, **Melanie**, **Mojdeh**, **Pearl**, **Pudsa**, **Nathalie**, **Tijn**, **Gwij**, **Jonah** en alle professoren: bedankt voor alles!!!

Mijn volgend woord van dank zou ik graag richten naar mijn goede vrienden. **Leen**, **Jeroen** en **Liesbeth**. Jullie hebben van in het begin tot op het einde alles in verband mijn doctoraat kunnen meevolgen. Goede of slechte momenten, ik kon/kan steeds op jullie rekenen. Dank jullie wel voor jullie oprechte interesse, luisterend oor, geduld, raad en steun. Ook de deugddoende afleiding en de onvergetelijke, leuke momenten die we samen hebben beleefd zullen me steeds bijblijven. Jullie zijn vrienden voor het leven!!!

Tenslotte zou ik graag mijn familie willen bedanken voor hun eindeloos geluld, hun aanmoedigingen en hun onuitputtelijke steun. **Mama** en **Papa**, dank jullie wel voor alles wat jullie voor mij hebben gedaan en de kansen die jullie me hebben gegeven. De jongens en ik komen voor jullie steeds op de eerste plaats, waarvoor onze oprechte dank!!! We kunnen ons geen betere ouders wensen!!! **Christophe** en **Philippe**: als jullie 'grote' zus wil ik enkel het allerbeste voor jullie en dat is duidelijk wederzijds. Jullie staan altijd klaar voor mij en zijn er steeds op de momenten dat ik het nodig heb. Dank jullie wel voor alles broertjes! **Sanne** en **Julie**, dank jullie om er

voor mij te zijn en voor de goede band/vriendschap. Mijn broers hebben fantastische schoonzusjes voor me uitgekozen en daar ben ik enorm dankbaar voor!!! Ook aan de rest van mijn familie een welgemeende dank je wel!

